

Corrigendum

Corrigendum to “Prognostic Factors for COVID-19 Hospitalized Patients with Preexisting Type 2 Diabetes”

Yuanyuan Fu,¹ Ling Hu,² Hong-Wei Ren,² Yi Zuo,^{1,2} Shaoqiu Chen,¹ Qiu-Shi Zhang,² Chen Shao,² Yao Ma,² Lin Wu,² Jun-Jie Hao,² Chuan-Zhen Wang,² Zhanwei Wang,³ Richard Yanagihara,⁴ and Youping Deng¹

¹Department of Quantitative Health Sciences, John A. Burns School of Medicine, University of Hawaii at Manoa, Honolulu, HI, USA

²Tianyou Hospital, Affiliated to Wuhan University of Science and Technology, Wuhan, Hubei, China

³Cancer Epidemiology Program, University of Hawaii Cancer Center, University of Hawaii at Manoa, Honolulu, HI, USA

⁴Department of Pediatrics, John A. Burns School of Medicine, University of Hawaii at Manoa, Honolulu, HI, USA

Correspondence should be addressed to Youping Deng; dengy@hawaii.edu

Received 14 June 2022; Accepted 14 June 2022; Published 4 July 2022

Copyright © 2022 Yuanyuan Fu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Prognostic Factors for COVID-19 Hospitalized Patients with Preexisting Type 2 Diabetes” [1], the authors would like to update the Authors’ Contributions section to note the equal contribution of Yuanyuan Fu and Ling Hu. The updated section is as follows.

Authors’ Contributions

YD and LH conceived and supervised the study. LH, QSZ, CS, YM, LW, JJH, and CZW collected the epidemiological and clinical data. LH, YZ, SC, and HWR contributed to radiological figure interpretation. YF and ZW processed and conducted statistical data analyses. YF, ZW, and RY drafted the manuscript. All the authors reviewed and approved the final version for publication. YF and LH contributed equally to this study. YD and YF are responsible for the integrity of the data and the accuracy of the analyzed data.

References

- [1] Y. Fu, L. Hu, H.-W. Ren et al., “Prognostic factors for COVID-19 hospitalized patients with preexisting type 2 diabetes,” *International Journal of Endocrinology*, vol. 2022, Article ID 9322332, 13 pages, 2022.