

## Research Article

# Ecology of *Echinops giganteus* A. Rich. in Sub-Saharan Africa: Distribution, Ecoclimatic Niche, and Phytosociology

Patrick Njiméli Sonkoué <sup>1,2</sup> Manuella Kelly Djeuga Youga <sup>1</sup>  
Carine Pamela Kenfack <sup>1</sup> Armel Boris Takenne Wamba,<sup>1</sup>  
Geo Coppens D'eckenbrugge <sup>2</sup> Cécile Dubois <sup>2</sup> and Théophile Fonkou <sup>1</sup>

<sup>1</sup>Research Unit of Applied Botany, Department of Plant Biology, Faculty of Science, University of Dschang, P.O. Box 67, Dschang, Cameroon

<sup>2</sup>CIRAD, UMRAGAP, Montpellier, France

Correspondence should be addressed to Patrick Njiméli Sonkoué; [sonkouepatrick@gmail.com](mailto:sonkouepatrick@gmail.com)

Received 7 August 2022; Revised 4 January 2023; Accepted 6 January 2023; Published 23 January 2023

Academic Editor: Ram Chander Sihag

Copyright © 2023 Patrick Njiméli Sonkoué et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Echinops giganteus* A. Rich. is an aromatic and medicinal plant of the Asteraceae family exploited in Cameroon under the access and benefit sharing (ABS) standard. Despite its importance, little information exists on the ecology of *E. giganteus*. The aim of the present study was to contribute to a better understanding of its ecology for sustainable management in the Western Highlands of Cameroon. Occurrence data as well as stationary ecological information were collected in the field and from different databases. Bioclimatic data were extracted from the WorldClim database and processed using DIVA-GIS and Maxent software. The Braun-Blanquet quadrat method was used for the phytosociological study. Results showed that the distribution of *E. giganteus* in its wild state is restricted to sub-Saharan Africa. This distribution is likely conditioned by altitude (1000 m–2600 m), light, temperature, and rainfall. The bioclimatic variables that best explained this distribution were the mean annual temperature (Bio1: 38.8%) and the precipitation of the coldest quarter (Bio19: 24.9%), and their favorable ranges were between 2°C–32°C and 300 mm–1800 mm, respectively. *E. giganteus* is a heliophilic plant that prefers well-drained substrates and would not have a requirement for organic matter. The floristic analysis of the *E. giganteus* community identified 68 plant species in 59 genera and 28 botanical families, with the most represented family being the Asteraceae (49%). The average species richness per quadrat was 8 species, dominated by herbs. Species consistent with *E. giganteus* were *Aspilia africana* (Pers.) C. D. Adams and *Imperata cylindrica* (L.) P. Beauv. Chamaephytes and Phanerophytes were dominant among the biological types, while the phytogeographic types were dominated by Pantropical species (38.23%). The most represented diaspore types and modes of dissemination were pogonochores (35.85%) and anemochores (55.38%).

## 1. Introduction

The forests of the Congo Basin abound in exceptional biodiversity that constitutes a potential for the socioeconomic development of its member countries [1, 2]. However, very little is known about this biodiversity, posing a need for the regular identification and description of new species through ethnobotanical and forestry inventories [3, 4]. Some of these species are endemic and have strong socioeconomic potential that can contribute to the development of local

communities [5]. Unfortunately, many of these known genetic resources and associated traditional knowledge have been exploited for decades illegally and without a sustainable management strategy to supply industries [6]. Over the years, the phenomenon has grown, and the countries of origin, as well as the rural populations, find themselves victims of the illegal appropriation of their resources. Moreover, access to these resources has become a real concern for the countries that hold genetic resources because they are diminishing at a frightening rate [7]. In Cameroon,

forest ecosystems occupy about twenty-one million hectares with a floristic diversity of more than 8000 species [8]. Increasing exploitation of its resources by populations and industries has resulted in an annual loss of area in the order of 0.3 to 1% [9]. These losses pose a management and conservation problem with a risk of genetic erosion of the species [10]. Indeed, the overexploitation of certain resources has resulted in the progressive loss of biodiversity and the increased vulnerability of certain species such as *Prunus africana* (Hook.fil.) Kalkman, *Garcinia kola* Heckel, *Garcinia lucida* Vesque, *Siphonochilus aethiopicus* (Schweinf.) B. L. Burtt, and *Warburgia salutaris* (Bertol.fil.) Chiov. [11, 12].

Access to genetic resources and benefit sharing (ABS) represents the third objective of the 1992 Convention on Biological Diversity. It was made effective in 2010 by the Nagoya Protocol and standardizes the mechanism of exploitation of genetic resources. The implementation of this mechanism is an effective measure in the sustainable management of genetic resources in that it would facilitate the conservation of biodiversity as well as its sustainable use [13, 14].

In Cameroon, numerous bioprospecting activities are being carried out to identify genetic resources with high economic potential that can be valorized under the ABS principle. Several aromatic plants have therefore been identified and prioritized based on the requirements of industries, including *E. giganteus* [15–18]. Although there are intense efforts to identify potentially valuable plants, very few studies have focused on their ecology [19, 20]. A lack of ecological knowledge can hinder the conservation and sustainable use of the resource.

*E. giganteus* is an annual herbaceous plant with erect port habit belonging to the Asteraceae family. Because of its multiple uses, it is of great importance to human beings. In food, its roots are used as a spice. Medicinally, it is used for the treatment of fever, respiratory diseases, toothache, leucorrhoea, and earache. It is also attributed to having anti-inflammatory properties [21]. A study by Fankam et al. [22] showed that its extract had an antibacterial effect on *Mycobacterium tuberculosis*. Several other studies demonstrate its anticarcinogenic activity on several human cancer cell lines [23–25]. In agriculture, it is used as a pesticide for food preservation during storage [26]. It is prized by the cosmetics and perfumery industries for the quality of its essential oil. As such, it is exploited in the Western Cameroon Highlands according to ABS standards [18]. However, its exploitation requires the destruction of the whole plant, which increases its vulnerability. Despite this high potential, its many uses, and its vulnerability, very little information is available on its ecology.

The present study was aimed at contributing to the sustainable management of *E. giganteus* in its natural environment. The objectives were to (i) establish its current distribution map, (ii) determine the abiotic factors as well as the characteristics of its ecoclimatic niche, and (iii) make an inventory of companion species. This will facilitate a better understanding of its ecology and guide policy makers in making the right decisions for sustainable management of the resource.

## 2. Material and Methods

**2.1. Study Area.** The Western Highlands of Cameroon lie between 4°54" and 6°36" North latitude and 9°18" and 11°24" East longitude and cover an area of 3.1 million hectares. The demographic pressure in this region is high (168 inhabitants/km<sup>2</sup>), with a contrasting topography which can be differentiated into as follows: (i) the granite-gneissic plateau in the south, which rises to between 1200 and 1400 m in altitude with poly convex reliefs; (ii) modest mountains of 2100 m in altitude in the southwest; (iii) the basaltic plateau in the north, with a calmer topography between 1400 m and 1600 m in altitude; and (iv) the Noun plain along the eastern edge of the Bamoun Plateau [27]. Soils are highly variable, ranging from andosols that are generally fertile and rich in organic matter on the High Plateaus (2000–2700 m) to ferrallitic soils, deep and impoverished by intense cultivation in the valleys (1400–1500 m) [28]. The climate is of the "Cameroonian Highland" type, marked by two seasons: a dry season from mid-November to mid-March, and a rainy season from mid-March to mid-November. Average temperatures are low (19°C) and decrease as one ascends to peaks [29]. Heavy rainfall (1500–2500 mm) is witnessed in a monomodal pattern, with relative humidity ranging from 40 to 100% [30]. The vegetation is deeply marked by altitude: tree and shrub savannahs with *Annona senegalensis* Pers. and *Terminalia glaucescens* Planch. ex Benth., and grassy savannahs with *Aframomum latifolium* K. Schum. and *Imperata cylindrica*; remnants of the coastal forest; gallery forests and montane formations; and cultivated land. Agriculture is strongly practiced there, representing more than 25.2% of the farms in the whole country, with a degree of land development of about 86% of cultivable land.

**2.2. Occurrence and Station Data.** Occurrence data, as well as stationary ecological information, on *E. giganteus* were collected from the GBIF database, from the Cameroonian herbarium labels in Yaoundé and Limbé, and during field trips in the Western Highlands Cameroon. Google Earth Pro and Geonames tools allowed us to verify, correct, and/or assign geographic coordinates from the GBIF database based on geographic information associated with gazetteers [31]. The selected occurrence data were submitted to ArcMap 10.3.1 spatial analysis software to obtain the distribution map of *E. giganteus*.

**2.3. Climate Niche Characteristics.** The bands of bioclimatic variables were extracted from the WorldClim database (<https://www.worldclim.org/data/worldclim21.html>) at 2.5 minutes (~21 km<sup>2</sup>) ground resolution. The 19 variables used correspond to current climate conditions and cover the period 1970–2000. They were derived from interpolations of data from more than 20,000 weather stations covering the 5 continents. These variables were as follows: (Bio1) mean annual temperature; (Bio2) mean diurnal amplitude (mean of monthly temperatures (max temp-min temp)); (Bio3) isothermality (Bio2/Bio7); (Bio4) seasonality of temperature; (Bio5) maximum temperature of the hottest month; (Bio6)

minimum temperature of the coldest month; (Bio7) annual amplitude of temperature; (Bio8) mean temperature of the wettest quarter; (Bio9) mean temperature of the driest quarter; (Bio10) warmest quarter average temperature; (Bio11) coldest quarter average temperature; (Bio12) annual precipitation; (Bio13) wettest month precipitation; (Bio14) driest month precipitation; (Bio15) precipitation seasonality; (Bio16) wettest quarter precipitation; (Bio17) driest quarter precipitation; (Bio18) warmest quarter precipitation; and (Bio19) coldest quarter precipitation. The values of each bioclimatic variable were extracted at the different points of occurrence using DIVA-GIS software. The relative contribution of each bioclimatic variable to the distribution of *E. giganteus* was evaluated with the Jackknife test after running the Maxent software.

#### 2.4. Floristic Analysis

**2.4.1. Sampling Method.** Data were collected at the end of the rainy season (November 2021). Phytosociological surveys were carried out in quadrats each of 25 m<sup>2</sup> size following the sigmatist method of Braun-Blanquet (Figure 1) [32, 33]. The installation of the quadrats was conditioned by the presence of *E. giganteus*. Its density was evaluated by the sum of the stems of *E. giganteus* observed in all the quadrats and then extrapolated to the hectare. The minimum distance between the different quadrats was at least 100 m. For each quadrat, several pieces of information were collected, notably the geographic coordinates of the quadrat, the list of species, and the global cover rate in the quadrat. For each species surveyed per quadrat, an abundance-dominance coefficient [34] was assigned: (+) species present as isolated individuals (covering less than 1%), or an average cover (MR) of 0.5%; (1) species present as sparse individuals occupying less than 5% of the area, or an MR of 3%; (2) species present as abundant individuals, covering 5–25% of the area, or an MR of 15%; (3) species with 25–50% cover, or 37.5% MR; (4) species with 50–75% cover, or 62.5% MR; (5) species with 75–100% cover, or 87.5% MR.

Fertile samples (sample with flowers and or fruits) collected and constituted in the herbarium were identified in the Herbaria of Limbé and Yaoundé.

(1) **Floristic Diversity.** The analysis of *E. giganteus* stand data focused on the calculation of floristic parameters and several diversity indices. These included

- (i) Relative frequency (RF) is the ratio of the number of surveys where a given species is present to the total number of quadrats;
- (ii) Relative abundance (RA), which expresses the importance of the species in terms of numbers as the ratio of the number of individuals of that species to the total number of species;
- (iii) Alpha diversity ( $\alpha$ ) is composed of parameters such as species richness, Shannon index ( $H'$ ), Simpson diversity index, and Pielou equitability ( $E$ );




FIGURE 1: Flora sampling device.

- (1) Species richness ( $S$ ) defined as the number of species that a community contains;
- (2) Shannon diversity index ( $H'$ ) expressing the diversity of species within plant communities [35]. It is calculated according to the following formula:

$$H' = - \sum_{i=1}^s (P_i \times \text{Logn } P_i), \quad (1)$$

where  $i$  takes value from 1 to  $s$ ,  $s$  = number of species in that area,  $P_i = ni/N$ , = individuals of one particular  $i$ th species found in an area,  $N$  = total number of individuals found in that area,  $\text{Logn}$  = natural logarithm,  $H' = 0$  if all individuals in the stand belong to one and the same species,  $H'$  is also minimal if, in a stand, each species is represented by a single individual, except for one species, which is represented by all other individuals in the stand. The index is maximal ( $H_{\max}$ ) when all individuals are equally distributed over all species [36, 37].

- (3) Pielou equitability or regularity ( $E$ ) is the ratio between the effective diversity ( $H'$ ) and the theoretical maximum diversity ( $H_{\max}$ ). It reflects the degree of diversity achieved by a stand in relation to the theoretical maximum and the quality of organization of a plant community. Its value is 0 when only one species is present and 1 when all species have the same abundance [38, 39].

$$E = \frac{H'}{H_{\max}}. \quad (2)$$

- (4) Simpson's index ( $D$ ) expresses the probability that two individuals drawn at random from an infinite population belong to the same species [40]. This index will have a value of 0 to indicate the maximum diversity, and a value of 1 to indicate the minimum diversity.

$$D = 1 - \frac{\sum n(N-1)}{N(N-1)}, \quad (3)$$

where  $n$  = total number of a particular species and  $N$  = total number of all species.

- (iv) Degree of presence ( $P$ ) is the percentage of records in which the species was noted relative to the total number of records in the cluster. The scale of presence classes is represented by Roman numerals, in centesimal proportions, as follows: (V) constant species, present in 81 to 100% of the records; (IV) species present in 61 to 80% of the records; (III) species present in 41 to 60% of the records; (II) species present in 21 to 40% of the records; (I) accidental species, present in 1 to 20% of the records.

## (2) Data Analysis and Processing

- (i) Biological types: The classification of biological types was done following the method of Mouton [41] used by Just and Raunkiaer [42]. We distinguished (1) Phanerophytes (Ph): higher plants whose regeneration buds were located at more than 50 cm from the ground; (2) Chamaephytes (Ch): perennial woody or suffructive species, whose regeneration buds were located at 50 cm from the ground at the most; (3) Geophytes (G): buds were located below the soil surface; (4) Hydrophytes (Hy): aquatic plants whose persistent buds were located at the bottom of the water; (5) Hemicryptophytes (H): perennial plants whose buds were outcropping on the soil surface; (6) Therophytes (Th): these were the annual plants that formed their spores or seeds during a single life period.
- (ii) Phytogeographic types: These were classified following the world chorology of White [43] and Saadou [44]. These are as follows: African (A) species found only in Africa; Paletropical (Pal) species common to Africa, tropical Asia, and some islands of the Indian Ocean; Pantropical (Pan) species found in all tropical regions of the world; Cosmopolitan (Cos) species widely distributed on the surface of the Earth; Afro-Malagasy (AM) species common to Africa, Madagascar, Comoros, Mascarene Islands, and Seychelles; and Afro-Neotropical (AN) species common to Africa and tropical America.
- (iii) Morphological types of species: The morphological type is generally used by several phytogeographers to explain the physiognomic and ecological organization of the vegetation of a territory or a region.
- (iv) Types of leaf size of species: The size of the leaf usually indicates the degree of adaptation of the plant to the hygrometry and the sunshine of the environment. Just and Raunkiaer [42] proposed the categories of leaf size, which were completed by the notophyll class by Webb [45].
- (v) Types of diaspores and modes of dispersal of species: Types of diaspores, as well as modes of dispersal, provide information on the ability of species to disperse and regenerate. They have been categorized according to the morphological criteria set by Dansereau and Lems [46] and Gorel et al. [47].

## 3. Results and Discussion

**3.1. Current Distribution of *E. giganteus*.** A total of 64 occurrences data were retained for the distribution map of *E. giganteus*. This distribution is restricted to sub-Saharan Africa, between latitudes 15°00'00"S and 15°00'00"N and longitudes 0°0'0" and 45°0'00"E. It is distributed in six countries: Cameroon, the Democratic Republic of Congo, Ethiopia, Nigeria, Rwanda, and Tanzania (Figure 2). Their location corresponds to the equatorial, monsoon, and tropical savannah climatic zones. This result adds to the work of Menut et al. [48] on aromatic plants of Central Africa, who observed that the distribution of *E. giganteus* was endemic to Cameroon and Nigeria.

**3.2. Ecological Characteristics.** *E. giganteus* was observed in grassy savannahs, fallows, and cultivated fields, with associated herbaceous vegetation, altitudes between 1000 m and 2600 m, at the level of rounded summits, and on slopes. It was observed to grow preferentially in the open environments with strong exposure to light. Indeed, the plants of *E. giganteus* growing in the mid-shade presented a delay of growth characterized by a stunted vegetative part and very small flower heads when they had flowered. This observation could reveal the heliophilic character of the species.

As reported by Tematio et al. [28] and Tsozué et al. [49], the soils in Western Highlands of Cameroon are dominated by the typical haplohumox group. These are the soils with sandy-clay texture, poor nitrogen, low mineralized organic matter, and slightly acidic pH. It could therefore be inferred that *E. giganteus* would prefer well-drained substrates and has no requirement for organic matter. This low organic matter requirement could also be justified by the absence of humus observed on the substrate of *E. giganteus* groups.

**3.3. Climatic Constraints.** The characterization of the ecoclimatic space of *E. giganteus* provided information on the climatic constraints that influence its distribution. The Jackknife test from the MaxEnt niche model of *E. giganteus* showed that the individual bioclimatic variables that most explained its distribution were as follows: Bio8 (mean temperature of the wettest quarter) which were the environmental variables with the highest gain. On its own, it seemed to have the most useful information for the model. It was followed by Bio1 (average annual temperature) and Bio10 (average temperature of the warmest quarter). When omitted, the environmental variable that reduced the gain most was Bio19 (precipitation of the coldest quarter). It could therefore contain some information that the other variables did not. It is followed by the environmental variables Bio8 (average temperature of the wettest quarter) and Bio18 (precipitation of the warmest quarter) (Figure 3).

Figure 4 presents the relative contributions of environmental variables to the MaxEnt model with a value greater than 1. It shows that the environmental variables that




FIGURE 2: Distribution of *Echinops giganteus* in Africa.




FIGURE 3: Statistical result of the jackknife test.

contributed most to the model were mean annual temperature (Bio1: 38.8%), precipitation in the coldest quarter (Bio19: 24.9%), and mean temperature in the wettest quarter (Bio8: 14.8%). These variables are derivatives of temperature and precipitation, and contribute 78.5% to the distribution




FIGURE 4: Contribution of each environmental variable to the niche model of *E. giganteus*.

of *E. giganteus*. The distribution of *E. giganteus* is thus conditioned by precipitation and temperature. This suggests that relative humidity, which is closely related to both temperature and humidity, may be a significant factor conditioning the distribution of *E. giganteus* on the continent. It is widely recognized that the likelihood of a species successfully establishing a population in a new environment is an ecological process ultimately determined by environmental and nonenvironmental factors [50]. According to many researchers, the distribution of plants in tropical Africa is mainly conditioned by total rainfall, the duration of the dry season, and air and soil moisture [51, 52]. This is because water availability directly affects physiological processes related to plant germination and growth [53].

Based on the logistic function curves of the MaxEnt model (Figure 5), it was possible to determine the ranges of

values favorable to the establishment of *E. giganteus* for the 3 variables that conditioned its distribution. This shows that, for logistic output values greater than or equal to 0.5, the ranges of mean annual temperature (Bio1), precipitation of the coldest quarter (Bio19), and mean temperature of the wettest quarter (Bio8) between 2°C–32°C, 300 mm–1800 mm, and 4 mm–20.5 mm, respectively, are those offering the best conditions for the distribution of *E. giganteus*.

**3.4. Taxon Diversity and Diversity Indices.** The total 43 surveys carried out permitted to find 68 plant species divided into 59 genera and 28 botanical families according to the APG3 classification. The Asteraceae family (49%) was the most represented with 15 genera and 29 species, followed by the Poaceae family (24%) with 10 genera and 10 species, then the Fabaceae family (7.6%) with 7 genera and 8 species (Figure 6). The average species richness per quadrat was 8 species, for an average cover of 13.41. The most represented species in the quadrats were *Aspilia africana* and *Imperata cylindrica* with presence degrees of 93% and 86%, respectively (Figure 7).

This predominance of the Asteraceae and Poaceae families provides information on the characteristics of the vegetation type of African savannas [54]. It could be explained by their capacity to spontaneously colonize the environment once the living conditions become favorable [55]. In fact, they have a very high tillering capacity, and a high regrowth capacity after grazing and bushfires [56]. They are also resistant to climatic hazards and cryptogamic diseases [57].

The density of *E. giganteus* is estimated to be 2177 feet/hectare. The mean values of Shannon's index, Simpson's index, and Pielou's Equitability were, respectively, 1.35 bits, 0.649, and 0.541. These values not only show a low specific diversity on the one hand but also a certain regularity of the species distribution around *E. giganteus* on the other hand. The Shannon–Weaver index obtained in this study is lower than that found by Ngyete Nyikob Mbogue et al. [58]. Indeed, in a similar study in the Western Highlands of Cameroon, these researchers obtained a Shannon–Weaver index of 6.92.

### 3.5. Ecological Spectra

**3.5.1. Spectrum of Biological Types.** The analysis of the global biological spectrum of the inventoried species showed the dominance of Chamaephytes and Phanerophytes in the whole of the studied flora, with respective proportions of 41.2% and 32.4%. Therophytes were in third place with 11.8%. Hemicryptophytes are less represented with 7.4%, and Geophytes come last with 4.4% (Figure 8).

The dominance of Chamaephytes and Phanerophytes reflects the adaptive strategies of plants related to competition and stress tolerance. Indeed, a predominance of Chamaephytes is characteristic of vegetation subjected to stresses of various origins. They are better adapted to drought, are more

xerophilic and produce many seeds [59]. In this study, the water and bushfire were the most observed stresses in the area. Indeed, the landscape being of the mountainous type with well-drained ferrallitic soils, would prevent water retention [28]. In addition, this area is subject to bushfires due to agricultural activities. The predominance of Phanerophytes reflects the adaptive strategies of plants related to competition for available resources. A study based on the analysis of ecological spectra of the flora of savannas of the West Cameroon Highlands by Wouokoue et al. [56] reveals that the latter is dominated by Phanerophytes and Chamaephytes. Our result is similar to that of Ngyete Nyikob Mbogue et al. [58] who noted that in the Western Highlands of Cameroon Chamaephytes and Phanerophytes were the most represented biological types associated with *E. giganteus*.

**3.5.2. Spectrum of Phytogeographic Types.** The phytogeographic types showed a dominance of pantropical (38.2%), Afrotropical (14.7%), cosmopolitan (11.8%), and paleotropical (11.8%) species. There was a low proportion of Afro-American, Afro-Malagasy, and Cameroonian Ridge endemic species, each with 1.5% of the gross spectrum.

Phytogeographic types are good indicators of the dynamism or stability of plant communities. A predominance of widely distributed (pantropical, cosmopolitan, and paleotropical) and continentally distributed (Afrotropical) species is characteristic of heliophilic vegetation such as grassy savannas. It is also an indication of environmental disturbance [60]. This disturbance of the environment in this study could be due to the anthropogenic activities such as agriculture, grazing, and bushfires observed at the study site (Figure 9).

**3.5.3. Types of Diaspores and Dispersal Modes.** The types of diaspores, as well as the modes of dissemination, provide information on the capacity of groups of individuals associated with *E. giganteus* to colonize an environment and perpetuate the species.

Seven types of diaspores corresponding to four modes of dissemination were identified in the different surveys. The most dominant types of diaspores were pogonochores (35.85%) and Sclerochores (24.5%) (Figure 10), while the most dominant types of dissemination were anemochores (55.38%) and autochores (29.23%) (Figure 11).

The high proportion of anemochores (pogonochores and sclerochorees) is justified by the abundance of Poaceae and Asteraceae, which are the main dissemination strategies for savanna plants [56]. This is quite logical given the fact that the study was conducted in a savannah environment.

**3.5.4. Types of Leaf Size.** This study allowed for the identification of seven types of leaf size, namely leptophylls, macrophylls, megaphylls, mesophylls, microphylls, nanophylls, and notophylls (Figure 12). The dominant leaf size types were notophylls (25.8%), mesophylls (25.8%), and microphylls (21.2%). Megaphylls and leptophylls were the least represented, with 4.5% and 3% of species, respectively. No aphyllous species were observed.




FIGURE 5: MaxEnt model logistic function curves of the response of key climate factors. (a) Mean annual temperature (Bio1), (b) precipitation of the coldest quarter (Bio19), and (c) mean temperature of the wettest quarter (Bio8).




FIGURE 6: Diversity of families most represented in the surveys.

The size of the leaves gives information on the hygrometry and the sunshine in a general way. It represents a response to altitude, local climatic conditions, and the regional orographic gradient. The predominance of notophylls, mesophylls, and microphylls is characteristic of slope and high-altitude vegetation. This result is similar to that of [58].

3.5.5. *Morphological Types.* The morphological types of companion species to *E. giganteus* were represented by grasses (80.3%), shrubs (9.1%), trees (6.1%), and lianas (4.6%) (Figure 13). This high representation of herbaceous species provides information on the dynamic and regressive states of the ecosystem. The study area is a savannah ecosystem subject to regular grazing, bushfires, and agricultural




FIGURE 7: Relative frequency of the most represented species.




FIGURE 8: Proportion of biological types: (Ch) *Chamaephytes*, (Ge) *Geophytes*, (He) *Hemicryptophytes*, (Ph) *Phanerophytes*, and (Th) *Therophytes*.




FIGURE 9: Proportion of phytogeographic distribution types: (AA) African, (Afro-Trop) Afro-Tropical, (AM) Afro-Malagasy, (Cos) Cosmopolitan, (GC) Guineo-Congolese, (Mo) Montane, (Pal) Paleotropical, (Pan) Pantropical, (Plur-African) Pluriregional African, and (SZ) Sudan-Zambezi.

activities. It is considered a degradation stage with floristic diversity that is sometimes important but ephemeral. It is therefore dominated by annual species with high seed

production that are capable of colonizing the environment when living conditions are favorable. This would justify the predominance of Asteraceae and Poaceae.




FIGURE 10: Proportion of diaspores types: (Acan) Acanthochores, (Ballo) Ballochores, (Desmo) Desmochores, (Pogo) Pogonochores, (Ptero) Pterochores, (Sarco) Sarcochores, and (Scléro) Sclerochores.




FIGURE 11: Proportion of different modes of diaspores dissemination: (Ane) Anemochory, (Auto) Autochory, (Endozoo) Endozoochory, and (Epizoo) Epizoochory.




FIGURE 12: Proportion of the different types of leaf sizes: (Le) Leptophylls, (Ma) Macrophylls, (Me) Mesophylls, (Mg) Megaphylls, (Mi) Microphylls, (Na) Nanophylls, and (No) Notophylls.

#### 4. Conclusion

*E. giganteus* is an endemic species to some countries in sub-Saharan Africa. Its ecological requirements are conditioned by substrate, altitude, sunshine, rainfall, and temperature. The predominance of biological types, diaspores types, and phytogeographic types in association with it informs about the regressive state of the vegetation. This implies a possibility that the species may be vulnerable in the Western




FIGURE 13: Proportion of morphological types.

Highlands of Cameroon in the future. Cameroonian authorities and companies exploiting the resource will need to take measures for sustainable exploitation in accordance with ABS (access to genetic resources and benefit sharing) principles. The cultivation of the resource should be favored over its exploitation in its natural state. In future work, spatiotemporal modeling of the ecoclimatic niche of *E. giganteus* will allow us to predict the potential impact of climate change on its distribution and to predict areas potentially favorable to its cultivation.

#### Data Availability

Occurrence data as well as stationary ecological information on *E. giganteus* were collected from the GBIF database, from the Cameroonian herbarium labels in Yaoundé and Limbé, and during field trips in the Western Highlands of Cameroon. The bands of bioclimatic variables were extracted from the WorldClim database (<https://www.worldclim.org/data/worldclim21.html>) at 2.5 minutes (~21 km<sup>2</sup>) ground resolution.

#### Conflicts of Interest

The authors declare that they have no conflicts of interest.

#### Acknowledgments

The authors are grateful to the Sud Expert Plantes Développement Durable (SEP2D) program and to V. Mane Fils Company for supporting the fieldwork in Cameroon. The French Ministry for Foreign Affairs granted a mobility scholarship to Montpellier at CIRAD, where most of the climatic data analysis was carried out.

#### References

- [1] E. T. Dongfack, R. Ngoufo, and S. Ongolo, "Conservation de la biodiversité et développement local en périphérie du Parc National de Korup au Cameroun," *Revue Scientifique et Technique Forêt et Environnement Du Bassin Du Congo*, vol. 15, pp. 88–100, 2020.
- [2] H. Tchekoté, N. Z. L. Meva'a, J. H. Moudingo, and N. Djofang, "Diagnostic de la conservation pour une gestion durable de la biodiversité dans le Bakossi, Banyang-Mbo, Régions du Sud-Ouest et du Littoral au Cameroun," *Revue Scientifique et*

- Technique Forêt et Environnement Du Bassin Du Congo*, vol. 15, pp. 49–59, 2020.
- [3] L. Zemagho, O. Lachenaud, and B. Sonké, “Four new species of Sabicea (Rubiaceae) from tropical Africa, with additional notes on the genus,” *Candollea*, vol. 73, no. 2, pp. 277–293, 2018.
- [4] M. Cheek and J. M. Onana, “The endemic plant species of Mt Kupe, Cameroon with a new Critically Endangered cloud-forest tree species, *Vepris zapfackii* (Rutaceae),” *Kew Bulletin*, vol. 76, no. 4, pp. 721–734, 2021.
- [5] M. I. Peya, *Impact de la crise écologique mondiale sur la biodiversité dans le bassin versant du Congo de 2013 à juin 2020*, 2020.
- [6] V. Boisvert, “Bioprospection et biopiraterie: le visage de Janus d’une activité méconnue,” *Cahier Du Gemdev*, vol. 30, pp. 123–136, 2005.
- [7] S. Brunel, “La biodiversité, un enjeu international,” in *Les missions militaires au service de la biodiversité*, pp. 21–44, EDP Sciences, Les Ulis, France, 2021.
- [8] R. Letouzey, “Notice de la carte phytogéographique du Cameroun au 1:500.000,” 1985, [https://scholar.google.com/scholar?hl=fr&as\\_sdt=0%2C5&q=%29.+Notice+de+la+carte+phytogéographique+du+Cameroun+au+1%2F500+000&btnG=](https://scholar.google.com/scholar?hl=fr&as_sdt=0%2C5&q=%29.+Notice+de+la+carte+phytogéographique+du+Cameroun+au+1%2F500+000&btnG=).
- [9] H. Bikie, O. Ndoye, and S. Wd, “L’Impact de la crise économique sur les systèmes agricoles et le changement du couvert forestier dans la zone forestière humide du Cameroun,” *Center For International Forestry Research*, vol. 27, p. 21, 2000.
- [10] B. Sonké, *Etudes floristiques et structurales des forêts de la réserve de faune du Dja (Cameroun)*, Université Libre de Bruxelles, Brussels, Belgium, 1998, <https://doi.org/565559%20565559-1001-08-07-2014%2014-17-49%20abbyy11.pdf>.
- [11] N. M. Guedje, B. A. Nkongmeneck, and J. Lejoly, “Composition floristique et structure des formations à *Garcinia lucida* dans la région de Bipindi-Akom II (Sud-Cameroun),” *Acta Botanica Gallica*, vol. 149, no. 2, pp. 157–178, 2002.
- [12] R. Kutalek and A. Prinz, “African medicinal plants,” *Handbook of Medicinal Plants*, CBS publishers, New Delhi, India, 2020.
- [13] T. Greiber, S. P. Moreno, M. Åhrén et al., “Guide explicatif du Protocole de Nagoya sur l’accès et le partage des avantages,” in *UICN – Droit et politique de l’environnement* no. 83, International Union for Conservation of Nature, Gland, Switzerland, 2014.
- [14] R. Z. Bastos, O. D. Canto, K. Galy, and I. Vestris, “Le régime international sur l’accès aux ressources génétiques au prisme de l’entrée en vigueur du Protocole de Nagoya,” *Revista de Direito Internacional*, vol. 13, no. 2, p. 16, 2016.
- [15] H. Gueaba Mbuntcha, H. Paulin Dogmo Fogang, V. Woguem et al., “Chemical characterization and antibacterial efficacy of essential oils of three lamiaceae species growing in Cameroon,” *Journal of Diseases and Medicinal Plants*, vol. 7, no. 1, pp. 14–21, 2021.
- [16] M. K. D. Youga, P. S. Njiméli, C. P. Kenfack, J. B. W. Taffo, W. N. Tacham, and T. Fonkou, “Knowledge and traditional uses of some aromatic and cosmetic plants species in the western highlands of Cameroon,” *Open Journal of Applied Sciences*, vol. 12, no. 10, pp. 1698–1718, 2022.
- [17] T. Fonkou, M. K. Djeuga Youga, H. Mbuntcha et al., “Valorisation des plantes aromatiques et cosmétiques suivant le principe d’accès et de partage des avantages, Bioprospection de la région de l’Ouest au Cameroun,” in *Biodiversité des écosystèmes intertropicaux. Connaissance, gestion durable et valorisation*, P. Jean-Pierre, A.-C. Stéphanie, B. Claire et al., Eds., p. 14, 2022, <https://www.editions.ird.fr/produit/662/9782709929394/biodiversite-des-ecosystemes-intertropicaux>, 2021st edition.
- [18] M. T. Mahop, “The post Nagoya Protocol ABS regime in Cameroon: exploring the extent to which ongoing policy, regulatory developments and ABS practices uphold the obligations of the Protocol,” *Global Transformations in the Use of Biodiversity for Research and Development*, Springer, Cham, Switzerland, pp. 207–236, 2022.
- [19] C. W. Fennell, K. L. Lindsey, L. J. McGaw et al., “Assessing African medicinal plants for efficacy and safety: pharmacological screening and toxicology,” *Journal of Ethnopharmacology*, vol. 94, no. 2–3, pp. 205–217, 2004.
- [20] T. Ticktin, “The ecological implications of harvesting non-timber forest products,” *Journal of Applied Ecology*, vol. 41, no. 1, pp. 11–21, 2004.
- [21] H. Bitew and A. Hymete, “The genus *Echinops*: phytochemistry and biological activities: a review,” *Frontiers in Pharmacology*, vol. 10, pp. 1234–1329, 2019.
- [22] A. G. Fankam, V. Kuete, I. K. Voukeng, J. R. Kuate, and J. M. Pages, “Antibacterial activities of selected Cameroonian spices and their synergistic effects with antibiotics against multidrug-resistant phenotypes,” *BMC Complementary and Alternative Medicine*, vol. 11, no. 1, p. 104, 2011.
- [23] V. Kuete, L. P. Sandjo, B. Wiench, and T. Efferth, “Cytotoxicity and modes of action of four Cameroonian dietary spices ethno-medically used to treat Cancers: *Echinops giganteus*, *Xylopiya aethiopyca*, *Imperata cylindrica* and *Piper capense*,” *Journal of Ethnopharmacology*, vol. 149, no. 1, pp. 245–253, 2013.
- [24] L. P. Sandjo, V. Kuete, X. N. Siwe, H. M. P. Poumale, and T. Efferth, “Cytotoxicity of an unprecedented brominated oleanolide and a new furoceramide from the Cameroonian spice, *Echinops giganteus*,” *Natural Product Research*, vol. 30, no. 22, pp. 2529–2537, 2016.
- [25] A. Somaida, I. Tariq, G. Ambreen et al., “Potent cytotoxicity of four cameroonian plant extracts on different cancer cell lines,” *Pharmaceuticals*, vol. 13, no. 11, pp. 357–419, 2020.
- [26] L. S. T. Ngamo and T. Hance, “Diversité des ravageurs des denrées et méthodes alternatives de lutte en milieu tropical,” *Tropicicultura*, vol. 25, no. 4, pp. 215–220, 2007.
- [27] M. A. Manga, A. Akoa, D. Abolo et al., “Structure et composition floristiques des agroforêts à base de caféiers arabica (*coffea arabica* l.) dans les hauts plateaux de l’ouest du cameroun,” *International Journal of Brain and Cognitive Sciences*, vol. 7, no. 4, p. 1474, 2014.
- [28] P. Tematio, L. Kengni, D. Bitom et al., “Soils and their distribution on bambouto volcanic mountain, west cameroon highland, central Africa,” *Journal of African Earth Sciences*, vol. 39, no. 3–5, pp. 447–457, 2004.
- [29] J. R. Fogaing, E. C. Ndonmou, M. Kuete Fogang, M. L. Avana-Tientcheu, and M. Tsalefac, “Potentiel de stockage de carbone des agro forêts du versant oriental des monts Bamboutos dans les Hautes Terres de l’ Ouest du Cameroun dans les Hautes Terres de l’ Ouest du Cameroun,” *Revue Scientifique et Technique Forêt et Environnement Du Bassin Du Congo*, vol. 16, p. 15, 2021.
- [30] A. Tsobeng, C. Mbosso, B. Takoutsing, E. Asaah, and A. Degrande, “Farmer perception about climate change adaptation and mitigation in rain forest and western highlands agro-ecological zones in Cameroon,” in *Proceedings of the World Agroforestry Centre (ICRAF-WCA)*, pp. 14–17, Nairobi, Kenya, August 2009.

- [31] G. Coppens d'Eeckenbrugge and J. M. Lacape, "Distribution and differentiation of wild, feral, and cultivated populations of perennial upland cotton (*Gossypium hirsutum* L.) in Mesoamerica and the Caribbean," *PLoS One*, vol. 9, no. 9, Article ID e107458, 2014.
- [32] J. Mangambu Mokoso, V. Noiha Noumi, L. Zapfack, and E. B. Sonké, "Etude phytosociologique du groupement à *Piper capensis* (RD Congo)," *International Journal of Environmental Studies*, vol. 67, no. 3, pp. 417–430, 2010.
- [33] M. Kuma, Z. Achiso, A. Chinasho, D. Yaya, and S. Tessema, "Floristics and diversity of invasive alien plant species in humbo district, south Ethiopia," *International Journal of Ecology*, vol. 2021, Article ID 6999846, 7 pages, 2021.
- [34] J. Braun-Blanquet, "Parc national suisse, Zerne et Bernina," *Bulletin de la Société Botanique de France*, vol. 98, no. 10, pp. 54–58, 1951.
- [35] C. E. Shannon and W. Weaver, "The theory of mathematical communication," *International Business*, vol. 131, 1949.
- [36] A. Lahed, S. Makhlof, S. Karim, G. Omar, and D. Salaheddine, "Quelques Aspects sur l'avifaune dans Deux Palmeraies du Sahara Septentrional (Algérie)," *Algerian Journal of Arid Environment*, vol. 3, no. 1, pp. 59–67, 2013.
- [37] A. Laurec, J.-C. Le Guen, and S. Frontier, "Collecte de l'information pour une gestion rationnelle des stocks halieutiques," *Stratégies d'échantillonnage en écologie*, vol. 17, pp. 385–415, 1983.
- [38] J. Blondel, "Biogéographie écologie masson," 1979, [https://scholar.google.com/scholar?hl=fr&as\\_sdt=0%2C5&q=Blondel%2C+J.+%281979%29+%3A+Biogéographie+écologie.+Masson%2C+Paris%2C+173p&btnG=](https://scholar.google.com/scholar?hl=fr&as_sdt=0%2C5&q=Blondel%2C+J.+%281979%29+%3A+Biogéographie+écologie.+Masson%2C+Paris%2C+173p&btnG=).
- [39] B. D. Ripley, "Spectral analysis and the analysis of pattern in plant communities," *Journal of Ecology*, vol. 66, no. 3, p. 965, 1978.
- [40] E. H. Simpson, "Measurement of diversity," *Nature*, vol. 163, no. 4148, p. 688, 1949.
- [41] J. A. Mouton, "Les types biologiques foliaires de Raunkiaer. Etat actuel de la question," *Bulletin de la Société Botanique de France*, vol. 113, no. 2, pp. 28–36, 1966.
- [42] T. Just and C. Raunkiaer, "The life forms of plants and statistical plant geography," *The American Midland Naturalist*, vol. 15, no. 6, p. 786, 1934.
- [43] F. White, "The vegetation of Africa," vol. 20, 1983, <https://www.cabdirect.org/cabdirect/abstract/19840692540>.
- [44] M. Saadou, *La végétation des milieux drainés nigériens à l'est du fleuve Niger*, Université Niamey, Niamey, Niger, 1990, [https://scholar.google.com/scholar?hl=fr&as\\_sdt=0%2C5&q=Saadou%2C+M.+%281990%29.+La+végétation+des+milieux+drainés+nigériens+à+l'est+du+fleuve+Niger.+Thèse+de+doctorat%2C+Université+Niamey%2C+Niger%2C+393p&btnG=](https://scholar.google.com/scholar?hl=fr&as_sdt=0%2C5&q=Saadou%2C+M.+%281990%29.+La+végétation+des+milieux+drainés+nigériens+à+l'est+du+fleuve+Niger.+Thèse+de+doctorat%2C+Université+Niamey%2C+Niger%2C+393p&btnG=).
- [45] L. J. Webb, "A physiognomic classification of Australian rain forests," *Journal of Ecology*, vol. 47, no. 3, p. 551, 1959.
- [46] P. Dansereau and K. Lems, "The grading of dispersal types in plant communities and their ecological significance," *Cabdirect.Org*, vol. 71, pp. 1–52, 1957.
- [47] A. P. Gorel, J. Duminil, J. L. Doucet, and A. Fayolle, "Ecological niche divergence associated with species and populations differentiation in *Erythrophleum* (Fabaceae, Caesalpinioideae)," *Plant Ecology and Evolution*, vol. 152, no. 1, pp. 41–52, 2019.
- [48] C. Menut, G. Lamaty, P. Weyerstahl, H. Marschall, I. Seelmann, and P. H. Amvam Zollo, "Aromatic plants of tropical Central Africa. Part XXXI. Tricyclic sesquiterpenes from the root essential oil of *Echinops giganteus* var. *lelyi* C. D. Adams," *Flavour and Fragrance Journal*, vol. 12, no. 6, pp. 415–421, 1997.
- [49] D. Tsozué, J. P. Nghonda, P. Tematio, and S. D. Basga, "Changes in soil properties and soil organic carbon stocks along an elevation gradient at Mount Bambouto, Central Africa," *Catena*, vol. 175, pp. 251–262, 2019.
- [50] R. Baker, R. Cannon, P. Bartlett, and I. Barker, "Novel strategies for assessing and managing the risks posed by invasive alien species to global crop production and biodiversity," *Annals of Applied Biology*, vol. 146, no. 2, pp. 177–191, 2005.
- [51] C. A. Adomou, B. Sinsin, and L. J. G. Maese, "Phytosociological and chorological approaches to phytogeography: a meso-scale study in Benin aridsteppe," *Systematics and Geography of Plants*, vol. 76, no. 2, p. 25, 2006.
- [52] A. Laouali, A. I. Salamatou, M. M. Boubacar, and M. Ali, "Impact potentiel du changement climatique sur la dynamique des peuplements de *Prosopis africana* (G. et Perr.) Taub. à l'horizon 2050 au Niger," *Journal of Applied Biosciences*, vol. 155, pp. 16049–16059, 2020.
- [53] M. Aoubouazza, "Estimation des besoins en eau du Cèdre à Ras El Ma et à Boutrouba (Moyen Atlas Central tabulaire)," *Revue Marocaine Des Sciences Agronomiques et Vétérinaires*, vol. 6, no. 1, p. 13, 2019.
- [54] A. Mahamane, M. Saadou, M. B. Danjimo et al., "Biodiversité végétale Au Niger: état des connaissances actuelles," *Université de Lomé: Accueil*, vol. 18, pp. 81–93, 2009.
- [55] A. Benaradj, H. Boucherit, O. Hasnaoui, and B. Babali, "Phytoécologie de la formation steppique à Drinn (*Stipagrostis pungens*) dans la région de Naâma (ouest Algérien) ESF1," *Flora Mediterranea*, vol. 31, pp. 159–171, 2021.
- [56] T. J. B. Wouokoue, A. Tientcheu, M. Louise et al., "Savannas highlands of Cameroon: floristic composition, functional traits and conservation status," *Asian Journal of Research in Botany*, vol. 4, no. 4, pp. 81–99, 2020.
- [57] G. S. Kouadja, A. C. Kouame, K. E. Kouadio, N. C. Kouassi, and B. J. Kouao, "Caractéristiques Nutritionnelles D'un Pâturage de Jachère Dans la Savane de Korhogo, au Nord de la Côte d'Ivoire," *European Scientific Journal*, vol. 17, no. 37, p. 14, 2021.
- [58] C. Ngyete Nyikob Mbogue, A. Mendi Grace, W. T. J. Baudoin, and N. F. Jong, "Phytosociology of *Echinops giganteus* in the western highland of Cameroon," *European Scientific Journal*, vol. 16, no. 36, pp. 345–360, 2020.
- [59] N. Bouazza, "Analyse de la diversité phytoécologique du chêne zeen (*Quercus faginea* subsp. *tlemcenensis* (DC): cas de la réserve de Moutas," Doctoral Dissertation, Université Djillali Liabes De Sidi Bel Abbes, Sidi Bel Abbès, Algeria, 2021, <http://hdl.handle.net/123456789/3287>.
- [60] B. Sinsin, "Formes de vie et diversité spécifique des associations de forêts claires du nord du Bénin," *Systematics and Geography of Plants*, vol. 71, no. 2, pp. 873–888, 2001.