Research Article

Exceptional Enlargement of the Mitochondrial Genome Results from Distinct Causes in Different Rain Frogs (Anura: Brevicipitidae: *Breviceps*)

Keitaro Hemmi,¹ Ryosuke Kakehashi,^{1,2} Chiaki Kambayashi,¹ Louis Du Preez,^{3,4} Leslie Minter,³ Nobuaki Furuno,¹ and Atsushi Kurabayashi^{1,2,3}

¹Amphibian Research Center, Hiroshima University, Hiroshima 739-8526, Japan

²Faculty of Bio-Science, Nagahama Institute of Bio-Science and Technology, Shiga 526-0829, Japan

³Unit for Environmental Sciences and Management, North-West University, Potchefstroom 2520, South Africa

⁴South African Institute for Aquatic Biodiversity, Somerset Street, Grahamstown 6139, South Africa

Correspondence should be addressed to Atsushi Kurabayashi; kuraba@nagahama-i-bio.ac.jp

Received 15 September 2019; Accepted 11 December 2019; Published 23 January 2020

Academic Editor: Hieronim Jakubowski

Copyright © 2020 Keitaro Hemmi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The mitochondrial (mt) genome of the bushveld rain frog (*Breviceps adspersus*, Brevicipitidae, Afrobatrachia) is the largest (28.8 kbp) among the vertebrates investigated to date. The major cause of genome size enlargement in this species is the duplication of multiple genomic regions. To investigate the evolutionary lineage, timing, and process of mt genome enlargement, we sequenced the complete mtDNAs of two congeneric rain frogs, *B. mossambicus* and *B. poweri*. The mt genomic organization, gene content, and gene arrangements of these two rain frogs are very similar to each other but differ from those of *B. adspersus*. The *B. mossambicus* mt genome (22.5 kbp) does not differ significantly from that of most other afrobatrachians. In contrast, the *B. poweri* mtDNA (28.1 kbp) is considerably larger: currently the second largest among vertebrates, after *B. adspersus*. The main causes of genome enlargement differ among *Breviceps* species. Unusual elongation (12.5 kbp) of the control region (CR), a single major noncoding region of the vertebrate mt genome, is responsible for the extremely large mt genome in *B. poweri*. Based on the current *Breviceps* phylogeny and estimated divergence age, it can be concluded that the genome enlargements occurred independently in each species lineage within relatively short periods. Furthermore, a high nucleotide substitution rate and relaxation of selective pressures, which are considered to be involved in changes in genome size, were also detected in afrobatrachian lineages. Our results suggest that these factors were not direct causes but may have indirectly affected mt genome enlargements in *Breviceps*.

1. Introduction

Mitochondrial (mt) genomes of multicellular animals (metazoans) are generally closed-circular and double-stranded DNA molecules approximately 14–20 kbp in length [1–3]. However, genome size ranges from 6 to 48 kbp [2, 4], and linear and fragmented mtDNAs have been found in approximately 9000 animals investigated thus far [5–7]. In most metazoans, this small organelle genome encodes a typical set of 37 genes: 13 protein genes, involved in the electron

transport system of respiration, two rRNA genes (*rrns*), and 22 tRNA genes (*trns*). In addition, animal mt genomes contain one long noncoding region, harboring several sequence elements related to mt genomic transcription and replication, named the control region (CR) or alternatively the D-loop region [2].

The mt genomic structure of metazoans, particularly vertebrates, tends to be conserved among closely related taxa. The same mt gene content and gene arrangement (synteny) are observed from fishes to mammals (e.g., [8, 9]). This

genomic organization would have arisen in a common ancestor of vertebrates and has been maintained in a wide range of vertebrate taxa over 400 million years [10, 11].

However, in some vertebrate taxa, increases in gene content due to gene duplication and rearrangements of mt gene positions are often observed (e.g., [8, 12]). In particular, a greater degree of structural change in mt genomes has been reported for amphibians, especially modern anurans (neobatrachians) (e.g., [13, 14]). Among anurans, the members of the basal group (a paraphyletic group generally called the “Archaeobatrachia”) retain the typical ancestral (i.e., symplexiomorphic) mt genome organization of vertebrates [15, 16]. In contrast, most of the neobatrachians share the derived positions of three *trns* translocated from their original locations (LTPF-*trn* cluster) [15, 17]. The Neobatrachia contains two superfamilies, Hyloidea and Ranoidea. The latter includes three large groups, Microhylidae, Natatanura, and Afrobatrachia [18, 19]. Of these, microhylid frogs retain the mt genomic structure of typical neobatrachian, while in natatanuran and afrobatrachian taxa, mt genomic rearrangements accompanying duplications and translocations of mt genes and the CR are often observed [13, 14, 16, 20, 21]. In particular, the mt genomes of afrobatrachians show large-scale structural changes.

The Afrobatrachia is a group of frogs, endemic to Africa, which currently consists of 422 described species in four families: Arthroleptidae, Brevicipitidae, Hemisotidae, and Hyperoliidae [22]. We have sequenced the complete mt genomes of four afrobatrachians representing all four afrobatrachian families [23]. All of these frogs tend to have large mt genomes, exceeding 20 kbp, and three of them have duplicated mt genes and/or CRs (excluding *Hemisus*). In particular, the mt genome of *Breviceps adspersus* (Brevicipitidae) has a highly reconstructed mt genome with many gene rearrangements and many duplicated gene regions. Consequently, the size of the *B. adspersus* mt genome is the largest among those vertebrates for which mtDNA has been sequenced to date (also the 13th largest among metazoans) [23].

To investigate the evolutionary origin and significance of the unusually large mt genome of *B. adspersus*, we analyzed the mt genomes of two additional *Breviceps* species, *B. mossambicus* and *B. poweri*. It has been suggested that the duplication of mt genes and the CRs, which results in an increase in genome size, are the result of nonadaptive evolution which, in insects, is correlated with an acceleration of nucleotide substitution rate [24] and a relaxation of purifying selective pressure, leading to a reduction in functional constraints that purge slightly deleterious mutations [25]. Thus, we investigated the changes in the substitution rate and selection pressure among afrobatrachian mt genomes and discuss the correlation between mt genome size and the change in the substitution rate and selective pressure.

2. Materials and Methods

2.1. Specimens Used. In this study, we used four frog specimens: one Mozambique rain frog, *Breviceps mossambicus* (Peters, 1854), and three Power’s rain frogs, *B. poweri* (Par-

ker, 1934) [26, 27]. These frogs were obtained via the pet trade; thus, their collection sites are unknown. The frog specimens were stored in 95.5% ethanol as part of AK’s personal amphibian collection. The voucher numbers are 15-004, 15-008, and 15-010 for *B. poweri* and 14-001 for *B. mossambicus*.

2.2. Molecular Experiments. We extracted and purified the total DNA from the liver tissue of each fixed specimen with a DNeasy Blood & Tissue Kit (QIAGEN K. K., Tokyo, Japan) or using phenol/chloroform extraction with “DNA sui-sui” extraction buffer (Rizo Inc., Tsukuba, Japan) and ethanol precipitation methods [28].

From the purified total DNA, the whole mtDNA was amplified by PCR and sequenced for *B. poweri* (voucher 15-004) and *B. mossambicus*. The PCR amplification and sequencing procedures from Kurabayashi et al. [10] were followed. Specifically, for each specimen, we amplified 10 overlapping fragments containing the entire mt genome using the long and accurate (LA) PCR method with LA-Taq (Takara Bio Inc., Shiga, Japan) according to the manufacturer’s instructions. These fragments were purified using the gel extraction method with a GenElute Agarose Spin Column (Sigma-Aldrich Japan Inc., Tokyo, Japan). The purified PCR fragments were sequenced using the primer walking method with an ABI 3130xl automated DNA sequencer (Applied Biosystems, Foster City, CA, USA). In this study, we used a total of 117 PCR primers for mtDNA amplification and/or sequencing, and 108 of them were newly designed during this study. All the primers used are listed in Supplementary Table S1.

2.3. NGS. The PCR fragments containing the CRs were very long, and these fragments harbored many direct repeat sequences that could not be read by the primer walking method (see Supplementary Fig. S1). Thus, these fragments were sequenced using single-molecule real-time (SMRT) sequencing with the PacBio RS II next-generation sequencer (NGS), which allows exceptionally long read sequencing (max length per read > 40 kbp) [29]. We also applied the multiplex-amplicon approach (see [29]). Briefly, we amplified the CR fragments from three *B. poweri* and one *B. mossambicus* specimens (approx. 13 and 8 kbp, respectively; see Supplementary Fig. S2) with the primers having distinct 3’ tag sequences for each specimen (consequently, the PCR fragments amplified from the same specimen have the same specific tag sequences, and thus, the PCR results were sortable from the mass NGS output-multiplex-amplicon method). Approximately 5 µg of the gel-purified CR fragments of two *B. poweri* (vouchers 15-004 and 15-010) and one *B. mossambicus* was used for the library construction for NGS. We outsourced the library construction and SMRT sequencing to CoMIT (Center of Medical Innovation and Translational Research) of Osaka University. The first SMRT run allowed for the determination of the entire CR sequences of *B. mossambicus* and one *B. poweri* specimen (15-010). However, we could not obtain enough sequence reads for another *B. poweri* specimen (15-004). Thus, for this specimen, we made two internal PCR primers (named bfCSB_Fow1 and bfCSB_Rev2, Supplementary Table S1) based on

the resultant CR sequence of *B. poweri* (15-010). Using these primers, two fragments of the 5' and 3' sides of the CR (approx. 4 and 9 kbp, respectively) were separately amplified from *B. poweri* (15-004) and the fragments were sequenced by another SMRT run. The assembled sequences of each CR fragment were reconstructed from the RS II outputs using the Long Amplicon Analysis program implemented in the SMRT Link analysis system [29].

The assembled whole mtDNA sequences of *B. mossambicus* and *B. poweri* (15-004) and the CR sequence of *B. poweri* (15-010) were deposited in the International Nucleotide Sequence Database Collaboration (INSDC) under the accession numbers LC498571, LC498572, and LC498573, respectively.

2.4. Phylogenetic Analyses and Divergence Time Estimation.

We performed phylogenetic tree reconstructions and a divergence time estimation by adding the sequence data obtained in this study to the dataset of Kurabayashi and Sumida [23], and the analytical methods used by Irisarri et al. [15] were followed. The previous dataset includes not only mt gene sequences but also nuclear genes. The nuclear data were also used in this study. Mitochondrial sequence data of afrobatrachians reported by Zhang et al. [16] were not used here because of a lack of genome size information and sequences of some mt genes. The genes used and their accession numbers are listed in Supplementary Table S2.

First, we aligned the sequences of each of the 13 protein genes, two *rrns*, and 22 *trns*, separately. The protein and RNA genes were aligned using the TranslatorX program with the default setting [30] and MAFFT with the L-INS-i option [31], respectively. The sequences of *trns* were manually aligned using their secondary structures as a guide. Ambiguously aligned sites were deleted using the Gblocks program ver. 0.91b with the default parameters [32]. The final alignment dataset consisted of 21,063 bp (consisting of 13,938 and 7125 mt and nuclear gene sequences, respectively) from 49 operational taxonomic units (OTUs). The alignment data used are provided in Supplementary Data S1.

We used the following sequences in the phylogenetic analyses: two mt *rrns*, 22 mt *trns*, and the 1st and 2nd codons of the 13 mt and nine nuclear protein genes (total 15,093 bp). We did not use the 3rd codon positions of the protein genes in the phylogenetic analyses (and divergence time estimation) because it is known that their fast substitution rates could distort the reconstruction of deep anuran phylogenies [33, 34]. We also applied a partitioning strategy in the phylogenetic analyses; i.e., the concatenated sequence data were partitioned into statistically suitable partitions and a distinct nucleotide substitution model was applied for each sequence partition. The PartitionFinder program ver. 1.1.1 [35] was used to estimate the best partitioning scheme using Akaike's information criterion (AIC) [36]. Consequently, a setting with 17 distinct partitions was suggested as the best partitioning scheme and this partitioning scheme was used in both the maximum likelihood (ML) and Bayesian inference (BI) analyses. An independent general time reversible+gamma distribution (GTR+G) substitution model for each of the 17

partitions was applied in the ML analysis. For the BI analysis, the best substitution model was estimated for each partition using the Kakusan4 program [37]. The detailed partitioning scheme and the suggested substitution models in BI are summarized in Supplementary Data S1.

We performed phylogenetic reconstructions using ML and BI methods. RAXML ver. 8.2.12 and MrBayes ver. 3.2.6 software packages were used for the ML and BI analyses, respectively [38, 39]. The rapid hill climbing algorithm (implemented in RAXML) with the starting tree option of 100 randomized parsimonious trees was applied for the ML analysis. For the BI analysis, 10 million generations of four Markov chains (MCs) were run with one sampling per every 1000 generations and the 1st 10% samples were discarded as burn-in. The convergence of the posterior distribution of model parameters (all parameters reached >200) was checked using Tracer ver. 1.5 [40]. The supports for the internal branches of reconstructed trees were evaluated using bootstrap percentages (BPs) calculated by 1000 pseudoreplications and Bayesian postprobabilities (BPPs) in ML and BI analyses, respectively.

The divergence times of anurans were estimated using a Bayesian dating method with the BEAST ver. 2.5.2 program package [41]. In the estimation, the amphibian phylogenies recovered from both ML and BI analyses were used as the topology constraint (Figure 1). The sauropsid monophyly (lizard+bird), not recovered by our ML and BI analyses, was *a priori* assumed in this dating analysis as in previous studies [15, 23]. The same data partitioning scheme and substitution models used in the BI analysis were also applied here. We used the Yule process [42] to describe cladogenesis. The final MCs were run twice for 100 million generations with one sampling per every 10,000 generations, and the 1st one million generations were discarded as burn-in. The posterior distributions of model parameters were checked in the same way as the above BI analysis. Following Irisarri et al. [15], we applied seven (lower age boundaries) calibration points as follows: (A) Sauropsida-Synapsida split: >312 million years ago (Ma), (B) Archosauromorpha-Lepidosauromorpha split: >260 Ma, (C) Cryptobranchidae-Hynobiidae split: >146 Ma, (D) Anura-Caudata split: >249 Ma, (E) most recent common ancestor (MCA) of Discoglossoidea: >161 Ma, (F) MCA of Pipoidea: >146 Ma, and (G) Calyptocephalella-Lechriodius split: >53 Ma. These were used as prior boundaries for divergence time estimation.

2.5. Relative Rate Test. We compared the relative rates of nucleotide substitutions of mt genes (all mt genes, all mt protein-coding genes, all *rrns*, and all *trns*) among afrobatrachian lineages using relative rate tests (RRTs) [43] with the RRTree program [44]. The Kimura two-parameter substitution model [45] was used for the estimation of genetic distances. In this analysis, we used the gene data of 24 neobatrachians. Noncompared lineages were used as the outgroups in each comparison (e.g., when we compared *Hemisus* and the three *Breviceps* species, the lineages of the remaining 20 neobatrachians were regarded as the outgroups). The lineages for each comparison are shown in Table 1.

TABLE 1: Comparisons of the relative rates of nucleotide substitutions among afrobatrachian-related lineages. The lineages with faster substitution rates are shown in bold.

Compared genes	Compared lineages		Relative substitution rates		Probability	Significance**
	Lineage 1	Lineage 2	Lineage 1	Lineage 2		
Microhylids vs. Afrobatrachians						
All 37 mt genes	Microhylids	Afrobatrachians	0.340	0.380	1.0×10^{-7}	**
All 13 protein genes	Microhylids	Afrobatrachians	0.391	0.428	1.0×10^{-7}	**
All rRNA genes	Microhylids	Afrobatrachians	0.165	0.204	1.0×10^{-7}	**
All tRNA genes	Microhylids	Afrobatrachians	0.225	0.286	1.0×10^{-7}	**
Laurentobatrachia (Arthroleptidae+Hyperoliidae) vs. Xenosyneunitanura (Hemisotidae+Brevicipitidae)						
All 37 mt genes	Laurentobatrachians	Xenosyneunitanurans	0.343	0.361	2.1×10^{-5}	**
All 13 protein genes	Laurentobatrachians	Xenosyneunitanurans	0.389	0.408	2.6×10^{-4}	**
All rRNA genes	Laurentobatrachians	Xenosyneunitanurans	0.178	0.192	0.098	
All tRNA genes	Laurentobatrachians	Xenosyneunitanurans	0.248	0.267	0.114	
<i>Hemiscus</i> vs. <i>Breviceps</i> (Hemisotidae vs. Brevicipitidae)						
All 37 mt genes	<i>Hemiscus</i>	<i>Breviceps</i>	0.369	0.358	0.058	
All 13 protein genes	<i>Hemiscus</i>	<i>Breviceps</i>	0.413	0.404	0.207	
All rRNA genes	<i>Hemiscus</i>	<i>Breviceps</i>	0.203	0.184	0.083	
All tRNA genes	<i>Hemiscus</i>	<i>Breviceps</i>	0.279	0.266	0.470	
<i>Breviceps mossambicus</i> vs. <i>B. adspersus</i>						
All 37 mt genes	<i>B. mossambicus</i>	<i>B. adspersus</i>	0.351	0.351	0.993	
All 13 protein genes	<i>B. mossambicus</i>	<i>B. adspersus</i>	0.393	0.394	0.906	
All rRNA genes	<i>B. mossambicus</i>	<i>B. adspersus</i>	0.185	0.190	0.514	
All tRNA genes	<i>B. mossambicus</i>	<i>B. adspersus</i>	0.279	0.266	0.237	
<i>Breviceps mossambicus</i> vs. <i>B. poweri</i>						
All 37 mt genes	<i>B. mossambicus</i>	<i>B. poweri</i>	0.351	0.375	4.6×10^{-6}	**
All 13 protein genes	<i>B. mossambicus</i>	<i>B. poweri</i>	0.393	0.423	5.9×10^{-6}	**
All rRNA genes	<i>B. mossambicus</i>	<i>B. poweri</i>	0.185	0.193	0.311	
All tRNA genes	<i>B. mossambicus</i>	<i>B. poweri</i>	0.279	0.287	0.551	
<i>Breviceps adspersus</i> vs. <i>B. poweri</i>						
All 37 mt genes	<i>B. adspersus</i>	<i>B. poweri</i>	0.351	0.375	3.7×10^{-6}	**
All 13 protein genes	<i>B. adspersus</i>	<i>B. poweri</i>	0.394	0.423	6.5×10^{-6}	**
All rRNA genes	<i>B. adspersus</i>	<i>B. poweri</i>	0.190	0.193	0.695	
All tRNA genes	<i>B. adspersus</i>	<i>B. poweri</i>	0.266	0.287	0.087	

**<0.001.

3. Results and Discussion

3.1. Huge mt Genomes of Rain Frogs. We determined the entire mt DNA sequences of two rain frogs, *B. mossambicus* and *B. poweri*. These mt genomes are 22,553 and 28,059 bp in length, which are very large for a vertebrate mt genome (typical size range is 16–17 kbp [1]). Although the genome size of *B. mossambicus* is similar to those of other afrobatrachian frogs (e.g., marbled reed frog (*Hyperolius marmoratus*): 22,595 bp and hairy frog (*Trichobatrachus robustus*): 21,418 bp; see Figure 1), the *B. poweri* mt genome was the second largest among the vertebrates investigated thus far (according to NCBI organelle genome resources [7] as of August 2019). The third largest vertebrate mt genome is

25,972 bp for the prickly gecko (*Heteronotia binoei*) [50]. Therefore, the mt genome of *B. poweri* is closer in size to that of *B. adspersus*, which possesses the largest known mt genome of vertebrates (28,757 bp).

In contrast to the large genome sizes, the mt gene content of both *B. mossambicus* and *B. poweri* is similar to that of many other vertebrates (Figure 1), containing the set of 37 single genes (13 proteins, two rRNA, and 22 tRNA genes) and single long and short noncoding regions commonly found in vertebrate mt genomes. The long noncoding region is referred to as CR and contains putative sequence elements involved in DNA replication and RNA transcription (e.g., terminate associate sequence (TAS) and conserved sequence blocks 1–3 (CSB1–3); see Supplementary Fig. S1 and [51]),

TABLE 2: Branch models used to estimate the dN/dS ratio (ω) of ranoid lineages.

Model	Constraint of the model	$-\ln L^*$ of the model	LRT** vs.			
			Model 0	1	2	3
0	Constant ω	All neobatrachian branches have single ω	148549.3	—	—	—
1	Two ω (Afrobatrachia)	Afrobatrachian branches have one unique ω	148536.2	3.2×10^{-7}	—	—
2	Two ω (Ranoidea)	Ranoid branches have one unique ω	148539.0	5.5×10^{-6}	NC***	—
3	Eleven ω	All afrobatrachian branches have distinct ω	148515.9	5.1×10^{-10}	1.3×10^{-5}	1.4×10^{-6}
4	Thirty-one ω	All ranoid branches have distinct ω	148443.6	9.6×10^{-29}	2.7×10^{-24}	2.6×10^{-25}
					6.6×10^{-21}	

* $-\ln L$: minus log likelihood. **LRT: P value of the likelihood ratio test. ***Models 1 and 2 have the same degrees of freedom (3) and cannot be compared by LRT. However, model 2 had a higher $-\ln L$.

and a short noncoding region is known as the *light*-strand replication origin (O_L). The presence of a small pseudogene, *trnS*(AGY), between NADH dehydrogenase subunits 4 and 5 (*nd4* and *nd5*) is the singular exception in gene content (Figure 1).

The gene content of *B. mossambicus* and *B. poweri* is not similar to that of the congeneric *B. adspersus* mt genome, which has many duplicated and triplicated gene regions and duplicated CRs, making it the largest known mt genome among vertebrates (Figure 1). Instead, the cause of the mt genome enlargement in *B. mossambicus* and *B. poweri* is the unusual expansions of the control regions. The lengths of their CRs are 6,618 and 12,537 bps, respectively, although the typical CR lengths in vertebrates are 1–2 kbp [1]. The long CRs of *B. poweri* and *B. mossambicus* result from the occurrence of multiple and long-tandem repeats (Supplementary Fig. S1). The *B. mossambicus* CR contains four distinct direct tandem repeats, and these repetitive sequences, totaling 4228 bp in length, occupy 63.9% of the CR. The CR of *B. poweri* (individual 4) has six distinct repeat sequence groups. Two of them are quite long (see Supplementary Fig. S1): one repeat group consists of 1,150 and 1,151 bp, nearly complete, repeat sequences (repeat 1: totaling 2301 bp), while another is composed of 23 units of 233–405 bp incomplete direct repeat sequences (repeat 3: totaling 7,339 bp). Consequently, the total length of the six repeat sequences in the *B. poweri* CR is 10,625 bp (occupying 84.7% of the CR nucleotides). The long CR over 10 kbp in length is quite rare and has never been completely sequenced in other vertebrate taxa (approximately 12 kbp CR occurred in a Malagasy frog (*Gephyromantis pseudoasper*) [20]).

To determine whether the unusually long CR is specific to the individual frogs used or is a common characteristic of the frog species, we checked the CRs of two additional *B. poweri* individuals. The PCR fragments having similar lengths (approx. 13 kbp, including the whole CR and 5' and 3' flanking gene regions) are commonly amplified in all three individuals (Supplementary Fig. S2). We also sequenced the CR of one of the two additional individuals. Although the CR lengths differ by 627 bp between the *B. poweri* individuals (12,537 bp vs. 11,910 bp), the CR sequences of the two *B. poweri* individuals are quite similar,

excluding the number of repeat units. Therefore, the very long CR, with numerous repetitive sequences, seems to be a common feature of this rain frog species.

It is almost impossible to precisely sequence long repetitive regions exceeding 10 kbp, such as those of the *B. poweri* CR by conventional sequencing methods (i.e., primer walking and construction of deletion mutants [20]) or by using the NGS technique with a short-read sequencing strategy [52]. In this study, a long-read strategy with PacBio RS II allowed us to relatively easily sequence such long repetitive sequences. These results demonstrate the superiority and necessity of long-read sequencing in analyzing the repetitive sequences occasionally found in both organelles and nuclear genomes.

3.2. Phylogeny, Timing, and Distinct Causes of mt Genome Enlargement in These Congeneric Rain Frogs. To infer the phylogenetic lineages and evolutionary times of the mt genome enlargement of afrobatrachians, including *Brevicipes*, we performed molecular phylogenetic analyses and divergence time estimation. The ML and BI methods reconstructed the same tree topology, and the maximum log likelihood ($\ln L$) of the resultant ML and the average $\ln L$ of the BI tree were -224667.83 and -221797.17, respectively. The time tree of amphibians having the ML/BI tree topology is shown in Figure 2. The major anuran phylogenies and the split ages agreed completely or to a substantial degree with those from previous studies: e.g., the monophyly of neobatrachians and paraphyly of archaeobatrachians with respect to neobatrachians (most recent common ancestor of Neobatrachia = 162.3 Ma), monophyly of ranoids (MCA = 127.0 Ma), and the three major clades of ranoids (Natatanura, Microhylidae, and Afrobatrachia, which started to split from 127.0 Ma) (e.g., [15, 16, 19, 53]).

The resultant phylogenies of afrobatrachian taxa completely match those of recent studies [16, 23, 54] (Figure 3), excluding the Afrobatrachia+Microhylidae clade not supported in Feng et al. [19]. Specifically, (1) Afrobatrachia is monophyletic, (2) Microhylidae becomes the sister taxon of Afrobatrachia, (3) Brevicipitidae forms a clade with Hemisotidae (this clade is called Xenosyneunitanura

FIGURE 2: Time tree of anurans. A phylogenetic tree reflecting the divergence ages estimated using a Bayesian relaxed dating method with the 15,093 bp nucleotide data. The tree topology of amphibians is the same as that of the resultant ML and BI trees. Bold branches indicate the lineages leading to the extant anurans. Horizontal blue bars on each node indicate 95% credibility intervals of estimated divergence age. The bootstrap probability (BP) of ML and Bayesian postprobabilities (BPP) are also shown on the right side of each node (BP value/*, ** > 95 and 99 BPPs), and the calibration points used in the dating analysis are indicated on the corresponding nodes (A to G). The scale of the horizontal axis is in million years.

[18]), (4) *Breviceps* is monophyletic and *B. poweri* is the basal taxon of the three species sampled in this genus, and (5) *B. mossambicus* and *B. adpersus* have a closer affinity within this sample. The estimated divergence ages of the corresponding nodes are as follows: (1) 118.1 Ma for the divergence between Afrobatrachia and Microhylidae, (2) 107.9 Ma for the most recent common ancestor of afrobatrachians, (3) 87.4 Ma for the split of Brevicipitidae and Hemisotidae, (4) 47.1 Ma for the divergence of *B. poweri* from the lineage of *B. mossambicus*+*B. adpersus*, and (5) 34.0 Ma for the split of *B. mossambicus* and *B. adpersus*. The divergence ages estimated here largely agree with those from recent studies [16, 23]. However, for the *Breviceps* divergences, younger ages have been esti-

mated by Nielsen et al. [54] (20 and 18 Ma for the divergences of 4 and 5, respectively). They used a total of 24 *Breviceps* taxa and applied some young calibration points within afrobatrachian lineages in their dating analyses. These differences in the data analysis could account for the differences in estimated divergence ages between our study and that of Nielsen et al. [54].

As described above, the mt genomes of microhylid frogs, the sister group of Afrobatrachia, exhibit average genome sizes for vertebrates (16.7–17.2 kbp) [55, 56]. In contrast, the afrobatrachian mt genomes exceed 20 kbp for all six species examined. Thus, the mt genome enlargement appears to be an evolutionary trend that has arisen in the lineages leading to the living afrobatrachians after the split from the

FIGURE 3: Changes in the dN/dS ratio (ω) among ranoid lineages. The estimated ω values of neobatrachian branches are shown (based on model 4 in Table 2). The tree topology is the same as those of the ML and BI trees reconstructed in this study. The constant ω (0.053) of nonranoid neobatrachian lineages was regarded as the background value. The estimated ω is shown on each ranoid branch. A high ω indicates the relaxation of purifying pressure. The branches for which ω values are lower (<0.053) and 1.5 times higher (>0.08) than the background are shown in blue and red colors, respectively. The frog taxa having mtDNAs exceeding >20 kbp and 28 kbp are also highlighted by orange and red colors, respectively.

microhylid lineage (Figures 2 and 3). In particular, the mt genome sizes of the two *Breviceps* species, *B. adspersus* and *B. poweri*, are over 28 kbp, making them as the largest and second largest known mt genomes of vertebrates. With regard to the interspecific phylogeny of *Breviceps*, these two species are not monophyletic; i.e., *B. adspersus* is closer to *B. mossambicus* than to *B. poweri* (Figure 1). Overall, our results indicate that the huge mt genomes arose in two independent rain frog lineages within relatively short periods (>47 and >30 million years for the lineages leading to *B. poweri* and *B. adspersus*, respectively). Furthermore, it is noteworthy that the physical causes of mt genome enlarge-

ment differ between these two rain frogs. The duplication of multiple gene/CRs and the accumulation of repetitive sequences are the main causes of mt genome enlargement in *B. adspersus* and *B. poweri*, respectively.

There are several examples of mt genome enlargement shared by congeneric species. For example, Malagasy poison frogs (*Mantella* spp.) commonly have >22 kbp mt genomes enlarged by the duplication of genes and CRs [20, 25], and two *Scapharca* ark shells exhibit >46 kbp mt genomes, mainly caused by elongations (>30 kbp) of noncoding sequences [4, 57]. However, there has been no similar example to that of *Breviceps* where the mt genome enlargement occurs

in independent congeneric lineages, from distinct causes. Thus, the mt genomic structure is highly variable in this frog taxon.

3.3. Gene Rearrangements and Evolutionary Inference. Although a highly rearranged mt genome is present in *B. adspersus* [23], the mt gene content and their arrangements of the congeneric *B. poweri* and *B. mossambicus* do not deviate largely from the typical mt genomes of vertebrates (Figure 1). The latter *Breviceps* species have almost the same gene arrangement, with the exception of two small translocations of *trnN* and light-strand replication origin (O_L) within the $WN-O_L-ACY$ *trn* cluster (Figure 1). Because the order of *trnsWN-O_L* found in *B. poweri* (and also in *B. adspersus*) is a primitive (plesiomorphic) gene order commonly shared by other afrobatrachians (Figure 1) [16, 23], the $O_L-trnsNW$ order of *B. mossambicus* appears to be newly emerged in the lineage leading to this species.

The *B. poweri* and *B. mossambicus* mt gene arrangements are similar to those of the mt gene orders of neobatrachians, especially *Hemisus*. Excluding the two minor *trn* translocations that occurred within the same *trn* clusters (specifically $O_L-trnsNW$ in *B. mossambicus* and *trnsPLTF* in *Hemisus*), the translocation of *trnsHS* is the only distinctive difference between *Breviceps* and *Hemisus*. Although the *trnsHS* is located between *nd4* and *nd5* in most neobatrachians, including *Hemisus*, this *trn* block lies between the cytochrome apoenzyme *b* gene (*cytb*) and CR in *B. poweri* and *B. mossambicus*. The novel *cytb-trnsHS-CR* arrangement could have arisen in a common ancestor of *Breviceps* after the split with other brevicipitid genera, for the following reasons: (1) the original *nd4-trnsHS-nd5* arrangement remains in another brevicipitid, *Callulina kreffti*, mt genome [16] and (2) the pseudogene of *trnS* is found at its original position between *nd4* and *nd5* in *B. poweri* and *B. mossambicus* (Figure 1).

The gene rearrangements in animal mt genomes are considered to reflect animal evolution (e.g., [23, 58–60]), and the rearranged gene orders found in this study can be regarded as novel phylogenetic markers for brevicipitid taxa. Specifically, the $O_L-trnsNW$ can be considered characteristic of some members of the *B. mossambicus* group [54], while the *cytb-trnsHS-CR* can be regarded as a synapomorphy of *Breviceps*.

In the *B. adspersus* mt genome, the *trnsHS* block is further translocated and positioned within the triplicated *trns-LTPF-12Srrn-trnV-16Srrn* cluster. In addition, duplications and translocations of the *trnsWN-OL* cluster and CR were also found in this species (Figure 1) [23]. The detailed phylogenetic lineage and evolutionary period of these large genomic modifications have not been clarified. This study shows that the mt genomes of *B. poweri* and *B. mossambicus* are not markedly rearranged from the typical neobatrachian type, although *B. adspersus* branches between these species' lineages (Figure 1 and see [54]). Therefore, it can be concluded that the unique mt genome of *B. adspersus* evolved only in the lineage leading to this species. Large mt genomic modifications resulting in stepwise gene rearrangements along with multiple lineage splits have been reported in Malagasy frogs (Mantellidae) [20]. However, in *Breviceps*, large-scale genomic changes occurred in a single species lineage

in a relatively short period (>47 Ma in this study and >20 Ma according to [54]), suggesting that the genomic structure of vertebrate mtDNA could harbor higher structural variability than is generally believed (e.g., [9]).

3.4. Substitution Rates and Selective Pressure on the Afrobatrachian mt Genomes and Their Correlations with Genome Size Expansion. As mentioned above, the duplication of genes/CRs and the occurrence of numerous repetitive sequences in the CR are responsible for the huge mt genomes of *B. adspersus* and *B. poweri*, respectively. Most of these events occurred in tandem, except for the nontandem duplications of CR and *trnS*(AGY) (Figure 1), which could have been caused by a nontandem duplication mechanism [20, 23]. In animal mt genomes, it is believed that both duplicated gene regions and tandem repeats in the CR have emerged because of errors in mtDNA replication, such as imprecise replication termination and strand slippage of the nascent DNA strand (e.g., [61, 62]). Furthermore, it has been suggested that accelerated nucleotide changes lead to the frequent tandem duplications via frequent substitutions at the initiation and termination points of mtDNA replication [24]. To investigate the correlation between genome size and nucleotide substitution rates, we compared the relative substitution rates of mt genes (four categories: all 37 genes, 13 protein genes, two *rrns*, and 22 *trns*) among afrobatrachian-related taxa using the RRT (Table 1).

The RRT showed that the substitution rates of afrobatrachian mt genes commonly having large genome sizes (>20 kbp) are significantly faster than those of the sister taxon, the microhylids, with normal genome sizes in all compared categories. Furthermore, the substitution rates of xenosyneunitanurans (Brevicipitidae+Hemisotidae), including the longest mt genomic species, are significantly higher than those of another afrobatrachian group, the laurentobatrachians (Arthroleptidae+Hyperoliidae) for all mt genes and protein genes (but are not significant for *rrns* and *trns*). These results suggest a correlation between the substitution rate and mt genome size. However, the substitution rates of the *Hemisus* mtDNA with a 20 kbp genome size are not significant but are faster than those of *Breviceps* species in all four compared categories. Among *Breviceps* species, the *B. poweri* mt genes show faster substitution rates compared to those of *B. mossambicus* and *B. adspersus*, although the latter species has the largest mt genome among vertebrates. From these results, it is concluded that mt genomes with large genome sizes also tend to have fast nucleotide substitution rates but the latter factor is not a direct cause of genome enlargement in afrobatrachian frogs.

It has been shown that animal mt genomes are subject to a strong purifying pressure that suppresses mutations leading to functional changes because the functions of mt coding genes are essential for respiration (e.g., [63]). Nevertheless, it has been reported that the purifying pressure of mt genomes is relaxed in neobatrachians, especially in the ranoid lineages [15, 23]. A relaxation of purifying selective pressure, leading to a reduction in functional

constraints that purge slightly deleterious mutations, has been suggested as one of the causes of mt genome enlargement [25]. In general, the ratio of nonsynonymous and synonymous substitutions (dN/dS ratio, ω) is useful to understanding the conditions for selection of the genes: positive selection ($\omega > 1$), neutral evolution ($\omega = 1$), and purifying selection ($\omega < 1$). Here, we estimate the dN/dS ratio using the branch model to investigate the changes in selective pressures in the ranoid lineages. In this analysis, we calculated the ω value(s) of the five branch models and compared the resultant log likelihood values among these models (Table 2). The LRT showed that model 4, with variable ω for all neobatrachian branches, is the best fitted among the models tested ($P = 6.6 \times 10^{-21}$ vs. model 3 with the 2nd largest likelihood value).

Figure 3 indicates the estimated ω values for the ranoid lineages under the best branch model (model 4). In all neobatrachian lineages, the estimated ω values are less than 1 (0.039–0.096) and confirm that the mt genomes of neobatrachian frogs are under strong purifying pressure. Compared to the background ω value (0.053) of the non-ranoid neobatrachian lineage, the ω values are high in 25 of 31 ranoid branches (the branches shown in black and red in Figure 3), indicating that the purifying selection has been relaxed in these lineages. Largely relaxed branches mainly correspond to the ancestral lineages, specifically common ancestral lineages leading to *Breviceps* (i.e., ranoids, afrobatrachians, and microhylids, afrobatrachians, and xenosyneunitanurans: $\omega = 0.096$, 0.090, 0.89, and 0.89, respectively). In contrast, the ω values of the branches leading to the extant frogs (i.e., terminal branches) tend to be lower. On the other hand, the ω values of six out of 16 terminal branches are less than the background ω value, indicating that the purifying selection increased again in these lineages.

Among anurans, the mt genomes with >20 kbp size have only been reported from ranoid taxa (see [7]) and obvious increases in the dN/dS ratio are found in most ranoid lineages. These facts appear to support the idea that the sizes of the mt genome can change concomitantly with lower selection strengths [25]. However, it is remarkable that reincreased purifying pressures are found in species lineages leading to *Mantella* and *Hoplobatrachus*, having >20 kbp mt genomes ($\omega = 0.051$ and 0.052, respectively). Furthermore, selective pressures of the species lineages of *B. adspersus* and *B. poweri* ($\omega = 0.071$ and 0.068, respectively), i.e., the exact lineages with increased genome size, are not as relaxed as those of their ancestral lineages. Consequently, our findings suggest that low selective pressure does not directly cause the huge mt genomes as does substitution rate (see the above). Rather, the existence of relaxed purifying selection in the ancestral lineages seems to have an indirect effect on the mt genome size. This indirect correlation might be caused by the accelerated accumulation of deleterious mutations under low selective pressure. Significantly, the mutations occur at the DNA replication-related sites which could induce numerous repetitive regions, leading to large mt genomes, as pointed out in Shao et al. [24].

4. Conclusion

In this study, we show that *B. poweri* has the second largest known mt genome among the vertebrates after its congeneric *B. adspersus*. We also found that the unusually large mt genomes did not arise in a common ancestor of these *Breviceps* species but rather that the genome enlargement occurred independently in each species lineage within relatively short periods. Consistent with this evolutionary inference, the causes of the genome enlargement differed between these species. At present, 19 nominal species are known in *Breviceps* [22], but only three mt genomes have been analyzed. It is remarkable that two of these mt genomes are the largest known among the vertebrates. Also, the mt genomic structures differ significantly between these congeneric species, suggesting high variability in the mt genomic structures in this genus. Future investigations of *Breviceps* mt genomes may provide new insights into the hidden diversity of vertebrate mt genomes.

Data Availability

The data used to support the findings of this study are available from the INSDC under the accession numbers LC498571, LC498572, and LC498573 and Supplementary Data S1.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Acknowledgments

We are indebted to CoMIT Omics Center of Osaka University for the PacBio RS II sequencing. We would like to thank Ms. I. Takemura, Nagahama Institute of Bio-Science and Technology, for her helpful assistance in the data analyses in the SMRT Link program. This study was supported by JSPS KAKENHI grants to AK: Grant-in-Aids for Challenging Exploratory Research (#26560437) and for Scientific Research (B) (#17H04608).

Supplementary Materials

Supplementary Table S1: list of primers used in this study. Supplementary Table S2: details on OTUs and gene data used. Supplementary Data S1: alignment data, partitioning strategy, and substitution models used in phylogenetic reconstructions and divergence time estimation. Supplementary Fig. S1: control regions of *Breviceps poweri* and *B. mossambicus*. Supplementary Fig. S2: PCR fragments of *Breviceps poweri* and *B. mossambicus* control regions. (Supplementary Materials)

References

- [1] D. R. Wolstenholme, "Animal mitochondrial DNA: structure and evolution," in *Mitochondrial Genomes*, D. R.

- Wolstenholme and K. W. Jeon, Eds., pp. 173–216, Academic Press, New York, 1992.
- [2] F. Pereira, J. Carneiro, and B. van Asch, “A guide for mitochondrial DNA analysis in non-human forensic Investigations~!2010-01-07~!2010-04-02~!2010-05-17~!,” *The Open Forensic Science Journal*, vol. 3, no. 2, pp. 33–44, 2010.
 - [3] D. V. Lavrov and W. Pett, “Animal mitochondrial DNA as we do not know it: mt-genome organization and evolution in nonbilaterian lineages,” *Genome Biology and Evolution*, vol. 8, no. 9, pp. 2896–2913, 2016.
 - [4] Y. Hou, B. Wu, Z.-H. Liu et al., “Complete mitochondrial genome of ark shell *Scapharca subcrenata*,” *Mitochondrial DNA Part A*, vol. 27, no. 2, pp. 939–940, 2016.
 - [5] E. Kayal, B. Bentlage, A. G. Collins, M. Kayal, S. Pirro, and D. V. Lavro, “Evolution of linear mitochondrial genomes in medusozoan cnidarians,” *Genome Biology and Evolution*, vol. 4, no. 1, pp. 1–12, 2011.
 - [6] F. Song, H. Li, G. H. Liu et al., “Mitochondrial genome fragmentation unites the parasitic lice of eutherian mammals,” *Systematic Biology*, vol. 68, no. 3, pp. 430–440, 2019.
 - [7] NCBI organelle genome resources Available from <http://www.ncbi.nlm.nih.gov/genome/browse!/organelles/>.
 - [8] J. L. Boore, “Animal mitochondrial genomes,” *Nucleic Acids Research*, vol. 27, no. 8, pp. 1767–1780, 1999.
 - [9] J. Warzecha, A. Fornal, M. Oczkowicz, and M. Bugno-Poniewierska, “A molecular characteristic of the Anatidae mitochondrial control region – a review,” *Annals of Animal Science*, vol. 18, no. 1, pp. 3–15, 2018.
 - [10] A. Kurabayashi, C. Usuki, N. Mikami et al., “Complete nucleotide sequence of the mitochondrial genome of a Malagasy poison frog *Mantella madagascariensis*: evolutionary implications on mitochondrial genomes of higher anuran groups,” *Molecular Phylogenetics and Evolution*, vol. 39, no. 1, pp. 223–236, 2006.
 - [11] S. Kumar, G. Stecher, M. Suleski, and S. B. Hedges, “TimeTree: a resource for timelines, timetrees, and divergence times,” *Molecular Biology and Evolution*, vol. 34, no. 7, pp. 1812–1819, 2017.
 - [12] R. L. Mueller and J. L. Boore, “Molecular mechanisms of extensive mitochondrial gene rearrangement in plethodontid salamanders,” *Molecular Biology and Evolution*, vol. 22, no. 10, pp. 2104–2112, 2005.
 - [13] A. Kurabayashi, N. Yoshikawa, N. Sato et al., “Complete mitochondrial DNA sequence of the endangered frog *Odorrana ishikawae* (family Ranidae) and unexpected diversity of mt gene arrangements in ranids,” *Molecular Phylogenetics and Evolution*, vol. 56, no. 2, pp. 543–553, 2010.
 - [14] R. Kakehashi, A. Kurabayashi, S. Oumi, S. Katsuren, M. Hoso, and M. Sumida, “Mitochondrial genomes of Japanese *Babina* frogs (Ranidae, Anura): unique gene arrangements and the phylogenetic position of genus *Babina*,” *Genes & Genetic Systems*, vol. 88, no. 1, pp. 59–67, 2013.
 - [15] I. Irisarri, D. S. Mauro, F. Abascal, A. Ohler, M. Vences, and R. Zardoya, “The origin of modern frogs (Neobatrachia) was accompanied by acceleration in mitochondrial and nuclear substitution rates,” *BMC Genomics*, vol. 13, no. 1, p. 626, 2012.
 - [16] P. Zhang, D. Liang, R.-L. Mao, D. M. Hillis, D. B. Wake, and D. C. Cannatella, “Efficient sequencing of anuran mtDNAs and a mitogenomic exploration of the phylogeny and evolution of frogs,” *Molecular Biology and Evolution*, vol. 30, no. 8, pp. 1899–1915, 2013.
 - [17] M. Sumida, Y. Kanamori, H. Kaneda et al., “Complete nucleotide sequence and gene rearrangement of the mitochondrial genome of the Japanese pond frog *Rana nigromaculata*,” *Genes & Genetic Systems*, vol. 76, no. 5, pp. 311–325, 2001.
 - [18] D. R. Frost, T. Grant, J. Faivovich et al., “The amphibian tree of life,” *Bulletin of the American Museum of Natural History*, vol. 297, pp. 1–291, 2006.
 - [19] Y. J. Feng, D. C. Blackburn, D. Liang et al., “Phylogenomics reveals rapid, simultaneous diversification of three major clades of Gondwanan frogs at the Cretaceous–Paleogene boundary,” *Proceedings of the National Academy of Sciences*, vol. 114, no. 29, pp. E5864–E5870, 2017.
 - [20] A. Kurabayashi, M. Sumida, H. Yonekawa, F. Glaw, M. Vences, and M. Hasegawa, “Phylogeny, recombination, and mechanisms of stepwise mitochondrial genome reorganization in mantellid frogs from Madagascar,” *Molecular Biology and Evolution*, vol. 25, no. 5, pp. 874–891, 2008.
 - [21] Y. Xia, Y. Zheng, I. Miura, P. B. Y. Wong, R. W. Murphy, and X. Zeng, “The evolution of mitochondrial genomes in modern frogs (Neobatrachia): nonadaptive evolution of mitochondrial genome reorganization,” *BMC Genomics*, vol. 15, no. 1, p. 691, 2014.
 - [22] D. R. Frost, *Amphibian species of the world: an online reference. Version 6.0*, American Museum of Natural History, New York, USA, 2019.
 - [23] A. Kurabayashi and M. Sumida, “Afrobatrachian mitochondrial genomes: genome reorganization, gene rearrangement mechanisms, and evolutionary trends of duplicated and rearranged genes,” *BMC Genomics*, vol. 14, no. 1, p. 633, 2013.
 - [24] R. Shao, M. Dowton, A. Murrell, and S. C. Barker, “Rates of gene rearrangement and nucleotide substitution are correlated in the mitochondrial genomes of insects,” *Molecular Biology and Evolution*, vol. 20, no. 10, pp. 1612–1619, 2003.
 - [25] B. Boussau, J. M. Brown, and M. K. Fujita, “Nonadaptive evolution of mitochondrial genome size,” *Evolution*, vol. 65, no. 9, pp. 2706–2711, 2011.
 - [26] W. C. H. Peters, “Diagnosen neuer Batrachier, welche zusammen mit der früher (24. Juli und 17. August) gegebenen Übersicht der Schlangen und Eidechsen mitgeteilt werden,” *Bericht über die zur Bekanntmachung geeigneten Verhandlungen der königlich-preussischen Akademie der Wissenschaften zu Berlin*, vol. 1854, pp. 614–628, 1854.
 - [27] H. W. Parker, *A monograph of the frogs of the family Microhylidae*, Trustees of the British Museum, London, 1934.
 - [28] J. Sambrook and D. W. Russell, *Molecular Cloning: A Laboratory Manual*, vol. 1, Cold Spring Harbor Laboratory Press, New York, 3rd Edition edition, 2001.
 - [29] Pacific Bioscience, “SMRT Link user guide (v5.1.0),” 2018, Available from <https://http://www.pacb.com/documentation/smart-link-user-guide-v5-1-0/>.
 - [30] F. Abascal, R. Zardoya, and M. J. Telford, “TranslatorX: multiple alignment of nucleotide sequences guided by amino acid translations,” *Nucleic Acids Research*, vol. 8, pp. 7–13, 2010.
 - [31] K. Katoh and H. Toh, “Recent developments in the MAFFT multiple sequence alignment program,” *Briefings in Bioinformatics*, vol. 9, no. 4, pp. 286–298, 2008.
 - [32] J. Castresana, “Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis,” *Molecular Biology and Evolution*, vol. 17, no. 4, pp. 540–552, 2000.
 - [33] S. Hoegg, M. Vences, H. Brinkmann, and A. Meyer, “Phylogeny and comparative substitution rates of frogs inferred from

- sequences of three nuclear genes," *Molecular Biology and Evolution*, vol. 21, no. 7, pp. 1188–1200, 2004.
- [34] C. Gissi, M. D. San, G. Pesole, and R. Zardoya, "Mitochondrial phylogeny of Anura (Amphibia): a case study of congruent phylogenetic reconstruction using amino acid and nucleotide characters," *Gene*, vol. 366, no. 2, pp. 228–237, 2006.
 - [35] R. Lanfear, B. Calcott, S. Y. W. Ho, and S. Guindon, "Partition-Finder: combined selection of partitioning schemes and substitution models for phylogenetic analyses," *Molecular Biology and Evolution*, vol. 29, no. 6, pp. 1695–1701, 2012.
 - [36] H. Akaike, "Information theory and an extension of the maximum likelihood principle," *Proceedings of the 2nd International Symposium on Information Theory*, B. N. Petrov and C. F. Budapest, Eds., , pp. 267–281, Akademiai Kiado, Budapest, 1973.
 - [37] A. S. Tanabe, "Kakusan4 and Aminosan: two programs for comparing nonpartitioned, proportional, and separate models for combined molecular phylogenetic analyses of multilocus sequence data," *Molecular Ecology Resources*, vol. 11, no. 5, pp. 914–921, 2011.
 - [38] A. Stamatakis, "RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies," *Bioinformatics*, vol. 30, no. 9, pp. 1312–1313, 2014.
 - [39] F. Ronquist, M. Teslenko, P. van der Mark et al., "MRBAYES 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space," *Systematic Biology*, vol. 61, no. 3, pp. 539–542, 2012.
 - [40] A. Rambaut and A. J. Drummond, 2009, Tracer v. 1.5 available from <http://tree.bio.ed.ac.uk/software/tracer/>.
 - [41] R. Bouckaert, T. G. Vaughan, J. Barido-Sottani et al., "BEAST 2.5: an advanced software platform for Bayesian evolutionary analysis," *PLoS Computational Biology*, vol. 15, no. 4, p. e1006650, 2019.
 - [42] G. U. Yule, "A mathematical theory of evolution, based on the conclusions of Dr. J. C. Willis, F.R.S," *Philosophical Transactions of the Royal Society B: Biological Sciences*, vol. 213, no. 402–410, pp. 21–87, 1925.
 - [43] P. Li and J. Bousquet, "Relative rate test for nucleotide substitutions between two lineages," *Molecular Biology and Evolution*, vol. 9, pp. 1185–1189, 1992.
 - [44] R. M. Robinson and D. Huchon, "RRTree: relative-rate tests between groups of sequences on a phylogenetic tree," *Bioinformatics*, vol. 16, no. 3, pp. 296–297, 2000.
 - [45] M. Kimura, "A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences," *Journal of Molecular Evolution*, vol. 16, no. 2, pp. 111–120, 1980.
 - [46] Z. Yang, *Computational molecular evolution*, Oxford University Press, New York, 2006.
 - [47] E. Buschiazzi, C. Ritland, J. Bohlmann, and K. Ritland, "Slow but not low: genomic comparisons reveal slower evolutionary rate and higher dN/dS in conifers compared to angiosperms," *BMC Evolutionary Biology*, vol. 12, no. 1, p. 8, 2012.
 - [48] Z. Yang, "PAML: a program package for phylogenetic analysis by maximum likelihood," *Computer Applications in the Biosciences*, vol. 13, no. 5, pp. 555–556, 1997.
 - [49] Z. Yang, "PAML 4: phylogenetic analysis by maximum likelihood," *Molecular Biology and Evolution*, vol. 24, no. 8, pp. 1586–1591, 2007.
 - [50] M. K. Fujita, J. L. Boore, and C. Moritz, "Multiple origins and rapid evolution of duplicated mitochondrial genes in parthenogenetic geckos (*Heteronotia binoei*; Squamata, Gekkonidae)," *Molecular Biology and Evolution*, vol. 24, no. 12, pp. 2775–2786, 2007.
 - [51] E. Jemt, Ö. Persson, Y. Shi et al., "Regulation of DNA replication at the end of the mitochondrial D-loop involves the helicase TWINKLE and a conserved sequence element," *Nucleic Acids Research*, vol. 43, no. 19, pp. 9262–9275, 2015.
 - [52] M. Mahmoud, M. Zywicki, T. Twardowski, and W. M. Karlowski, "Efficiency of PacBio long read correction by 2nd generation Illumina sequencing," *Genomics*, vol. 111, no. 1, pp. 43–49, 2019.
 - [53] R. A. Pyron and J. J. Wiens, "A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of extant frogs, salamanders, and caecilians," *Molecular Phylogenetics and Evolution*, vol. 61, no. 2, pp. 543–583, 2011.
 - [54] S. V. Nielsen, R. Savel, S. R. Daniels, W. Conradie, M. P. Heinicke, and B. P. Noonan, "Multilocus phylogenetics in a widespread African anuran lineage (*Brevicipitidae: Breviceps*) reveals patterns of diversity reflecting geoclimatic change," *Journal of Biogeography*, vol. 45, no. 9, pp. 2067–2079, 2018.
 - [55] P. Zhang, H. Zhou, Y. Q. Chen, Y. F. Liu, and L. H. Qu, "Mitogenomic perspectives on the origin and phylogeny of living amphibians," *Systematic Biology*, vol. 54, no. 3, pp. 391–400, 2005.
 - [56] D. S. Hwang and J. S. Lee, "Complete mitochondrial genome of the boreal digging frog *Kaloula borealis* (Anura, Microhylidae)," *Mitochondrial DNA*, vol. 23, no. 4, pp. 301–302, 2012.
 - [57] Y. G. Liu, T. Kurokawa, M. Sekino, T. Tanabe, and K. Watanabe, "Complete mitochondrial DNA sequence of the ark shell *Scapharca broughtonii* : An ultra-large metazoan mitochondrial genome," *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, vol. 8, no. 1, pp. 72–81, 2013.
 - [58] J. L. Boore, T. M. Collins, D. Stanton, L. L. Daehler, and W. M. Brown, "Deducing the pattern of arthropod phylogeny from mitochondrial DNA rearrangements," *Nature*, vol. 376, no. 6536, pp. 163–165, 1995.
 - [59] J. L. Boore and W. M. Brown, "Big trees from little genomes: mitochondrial gene order as a phylogenetic tool," *Current Opinion in Genetics and Development*, vol. 8, no. 6, pp. 668–674, 1998.
 - [60] J. Y. Zhang, L.-P. Zhang, D.-N. Yu, K. B. Storey, and R.-Q. Zheng, "Complete mitochondrial genomes of *Nanorana taihangnica* and *N. yunnanensis* (Anura: Dicroglossidae) with novel gene arrangements and phylogenetic relationship of Dicroglossidae," *BMC Evolutionary Biology*, vol. 18, no. 1, p. 26, 2018.
 - [61] C. Moritz and W. Brown, "Tandem duplications in animal mitochondrial DNAs: variation in incidence and gene content among lizards," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 84, no. 20, pp. 7183–7187, 1987.
 - [62] Y. Xia, Y. Zheng, R. W. Murphy, and X. Zeng, "Intraspecific rearrangement of mitochondrial genome suggests the prevalence of the tandem duplication-random loss (TDLR) mechanism in *Quasipaa boulengeri*," *BMC genomics*, vol. 17, no. 1, p. 965, 2016.
 - [63] J.-Y. Li, C. Zeng, G.-Y. Yan, and L.-S. He, "Characterization of the mitochondrial genome of an ancient amphipod *Halice* sp. MT-2017 (Pardaliscidae) from 10,908 m in the Mariana Trench," *Scientific Reports*, vol. 9, no. 1, p. 2610, 2019.