

Research Article

Awareness of Hypertension, Hypercholesterolemia, and Diabetes Mellitus and Associated Characteristics in Russian Adults

Filip Sahatqija,¹ Monica Hunsberger,¹ Sarah Cook,² Kamila Kholmatova,^{3,4} Marina Shapkina,⁵ Sofia Malyutina,^{5,6} and Alexander V. Kudryavtsev^{3,4}

¹Shalgrenska Academy, University of Gothenburg, Gothenburg 41390, Sweden

²School of Public Health, Faculty of Medicine, Imperial College London, London SW7 2AZ, UK

³International Research Competence Centre, Northern State Medical University, Arkhangelsk 163069, Russia

⁴Department of Community Medicine, UiT The Arctic University of Norway, Tromsø N-9037, Norway

⁵Research Institute of Internal and Preventive Medicine, Branch of Institute of Cytology and Genetics,

Siberian Branch of the Russian Academy of Sciences, Novosibirsk 630089, Russia

⁶Department of Therapy, Haematology and Transfusiology, Novosibirsk State Medical University, Novosibirsk 630090, Russia

Correspondence should be addressed to Monica Hunsberger; monica.hunsberger@gu.se

Received 30 November 2023; Revised 19 February 2024; Accepted 2 March 2024; Published 25 March 2024

Academic Editor: Massimo Salvetti

Copyright © 2024 Filip Sahatqija et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Russia has higher cardiovascular disease (CVD) mortality compared to other European countries. The major CVD risk factors are age, male sex, and three conditions, namely hypertension, hypercholesterolemia, and diabetes mellitus (DM). This study aimed to assess awareness of these three conditions among Russian adults (N= 3803) and the associated socio-demographic, lifestyle, and health characteristics. We used cross-sectional data from a randomly drawn population-based sample of Russians aged 35–69 years, who participated in the Know Your Heart (KYH) study conducted in Arkhangelsk and Novosibirsk between 2015–2018. Participants' self-reported awareness of hypertension, hypercholesterolemia, and DM was assessed against the measures at the KYH health check (blood pressure, cholesterol, HbA1c and/or use of medication for each condition). Prevalence estimates for the awareness were age- and sex-standardized to the Standard European Population. Socio-demographic, lifestyle, and health-related correlates of the awareness were investigated using logistic regression modelling. Among participants with hypertension (N= 2206), hypercholesterolemia (N= 3171), and DM (N= 329) recorded at a health check, 79%, 45%, and 61% self-reported these conditions, respectively. Higher awareness of hypercholesterolemia and hypertension was associated with older age, female sex, nonsmoking status, obesity, and history of CVD diagnoses. Low household income and history of CVD diagnoses were associated with being aware of DM. The awareness rates of hypertension were relatively high, whereas awareness rates of hypercholesterolemia and DM were asasociated or DM. The awareness rates of hypertension could be improved in Russia through increasing the awareness of the risk factors.

1. Introduction

Cardiovascular diseases (CVDs), a cluster of disorders of heart and blood vessels, are the leading cause of death globally [1]. However, many countries have reduced CVD incidence and mortality through the successful implementation of prevention programmes and the introduction of surgical treatment strategies [2]. Eastern European countries fall behind other European countries in reducing CVD mortality rates [3]. In 2019, the lowest age-standardized CVD mortality rate of 190 per 100,000 population was registered in France [4]. For comparison, Russia had a CVD mortality rate of 574 per 100,000 population in 2019 [5], which was the third highest in Europe after Ukraine and Bulgaria [6]. Despite existing research, the causes of the high CVD mortality rates in Russia are still not extensively explained [7, 8]. Hypertension, hypercholesterolemia, and diabetes mellitus (DM) are the major CVD risk factors [9, 10]. At the early stages, these conditions are often latent due to their asymptomatic or mild manifestations [11–13], and individuals may be unaware of these conditions until they progress to exhibiting symptoms or complications [14]. Public health interventions aimed at raising awareness and early diagnosis of modifiable risk factors are acknowledged tools to reduce CVD morbidity and mortality [14]. To be effective, campaigns must be population-specific and culturally appropriate to target the high-risk groups [15].

Previous research has reported associations between the awareness of hypertension, DM, and hypercholesterolemia and sociodemographic factors (age, sex, education level, employment status, place of residence, and income) [3, 16, 17]. Other authors also described associations of the awareness with behavioural factors such as physical activity, smoking, alcohol consumption, obesity, and previous experience with cardiovascular events or comorbidities [18–20].

Awareness of hypertension in Russia was previously estimated to be 68-80% in males and 76-86% in females [3, 21–24]. Factors associated with hypertension awareness were older age [21–23], female sex [3, 21, 23], higher level of education among males [21], and urban residence [21]. The HAPIEE study (2002–2005) showed relatively low awareness of diabetes (30.8% in males and 45.2% in females) and hypercholesterolemia (11.2% in males and 15.3% in females) in Russia compared with Poland, Lithuania, and the Czech Republic [3]. Several studies assessed the control of DM and high cholesterol [25–27], but we could not find Russian population-based studies focusing behavioral and health characteristics associated with the awareness.

This study aimed to assess awareness of hypertension, hypercholesterolemia, and DM in Russian adults with objective confirmation of these conditions and to investigate socio-demographic, lifestyle, and health-related correlates of awareness.

2. Materials and Methods

2.1. Study Design. The study is based on data from the "Know Your Heart" (KYH) cross-sectional study, conducted from 2015–2018 in two Russian cities, Arkhangelsk and Novosibirsk, with a general population sample aged 35–69 years.

2.2. Recruitment of Participants. The study's sampling frame was drawn from the databases of the Arkhangelsk and Novosibirsk regional health insurance funds (lists of depersonalized addresses of residents with mandatory health insurance supplemented by age and sex). Random addresses were selected for home visits, stratified by age and sex, targeting to recruit equal numbers of participants by sex and 5-year age groups. Trained interviewers visited the selected addresses and invited household members of the predefined age (± 2 years) and sex to participate in the study, one person per household. After the ascertainment of informed consent, a participant underwent a baseline interview at home. After the interview, participants were invited to a health check at a polyclinic. The health check comprised a physical examination of cardiovascular health and a medical interview, which included the collection of self-reported data on health status and medication use. Details of the KYH study rationale, design, sampling, and data collection procedures have been published previously [28].

The response for the baseline interview was 51.0% (68.2% in Arkhangelsk, 41.4% in Novosibirsk) of the total invitees. This reflects the proportion of all eligible participants of the relevant age and sex who were approached, invited, and agreed to participate in the study. After initial participation and interview, there was 96% health check attendance in Arkhangelsk and 83% in Novosibirsk, with an average 16-day gap to baseline interviews.

2.3. Data Collection. Systolic and diastolic blood pressure (SBP and DBP) were measured at the health check using an OMRON 705 IT automatic blood pressure monitor (OMRON Healthcare). The measurements were performed in a seated position three times with two-minute intervals between. The mean values of the second and third measurements were used in the analysis.

Blood samples were taken at the health check to assess cardiometabolic parameters, including total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and glycated haemoglobin (HbA1c). Participants were asked to fast for at least four hours prior to the health check. The blood samples were aliquoted, frozen, and stored. Throughout the data collection, the samples were shipped on dry ice to a laboratory in Moscow and stored at -80° C. All laboratory analyses were performed in one batch at the end of the study. Levels of TC (mmol/L) and LDL-C (mmol/L) were assessed in blood serum using enzymatic color tests, and HbA1c (%) levels were assessed in the whole blood using immuno-turbidimetric tests (AU 680; Chemistry System Beckman Coulter).

Participants were requested to bring their current medications to the health check, and 27% of them did so. They were asked to show their currently used medications or if they did not bring them, to list them. Commercial names for up to seven medications per participant, as well as doses and frequencies were recorded and subsequently classified at the end of the study using the international WHO anatomical therapeutic chemical (ATC) classification system version 2016 [29].

Self-reports of hypertension, hypercholesterolemia, and DM were collected with three questions asked in the medical interview: "Have you ever been told by a doctor or nurse that you have.... high blood pressure?," "high cholesterol?," or "diabetes mellitus?" Responses like "do not know" were considered as negative answers.

2.4. Ascertaining the Presence of Risk Factors. Hypertension was defined as SBP >140 mmHg and/or DBP >90 mmHg at the health check and/or reported daily intake of antihypertensive medication (ATC classes C02, C03, C07, C08, or C09). Hypercholesterolemia was ascertained if a participant had total cholesterol \geq 5.2 mmol/L and/or LDL cholesterol of >3.0 mmol/L and/or reported daily intake of lipid lowering medication (ATC class C10). Diabetes was defined as HbA1c \geq 6.5% and/or self-reported the intake of antidiabetic medication (ATC class A10).

2.5. Assessing Awareness of Risk Factors. Awareness of hypertension, hypercholesterolemia, or DM was considered present if a participant had the condition(s) ascertained at the health check and self-reported positively to "have you ever been told by a doctor or nurse that you have" the conditions under investigation.

2.6. Associated Characteristics. To describe sociodemographic correlates of the awareness of risk factors, we used the following variables from baseline interview: age (years), sex (male or female), completed higher education (yes or no), and participant's self-reported financial constrains indicating household income (low, middle, and high). Low income was defined as having financial difficulties to buy food or clothes; middle income—having enough money for food and clothes but experiencing constrains in buying large domestic appliances or a new car; high income—reporting no difficulties to buy a large new car but constraints to buy a flat or house, or no financial constraints at all.

Smoking data were recorded as self-reported daily smoking, and participants were divided into never, former, and current smokers. Data on alcohol consumption were collected using the Alcohol Use Disorders Identification Test (AUDIT) [30]. An AUDIT score ≥ 8 was defined as hazardous drinking.

Height (cm) was measured at the health check using the Seca® 217 stadiometer (Seca Ltd., Hamburg, Germany). Weight (kg) was measured using TANITA BC 418 body composition analyser (Tanita Corp., Tokyo, Japan). Body mass index (BMI) was calculated as weight in kilograms divided by squared height in meters. Obesity was defined as a BMI \geq 30 kg/m².

Data on the history of CVD diagnoses were obtained by asking the participants about having ever been diagnosed with angina, stroke, myocardial infarction, atrial fibrillation, or heart failure.

2.7. Sample. A total of 5089 men and women had the baseline interview as a part of the KYH study (Figure 1). Out of them, 547 failed to attend the health check, and 38 of the attendees did not provide the consent for sharing the collected data with third parties. Therefore, based on the data access application, the first author received the anonymized data of 4504 KYH participants, 2362 from Arkhangelsk and 2142 from Novosibirsk, limited to the variables used in this study. Another 701 participants were excluded from analyses for the following reasons: 27 were older than 69 years by the time of the health check, and 674 had missing or unusable data on one or more variables of interest. Correspondingly, 3803 KYH participants were included in the analyses. Participants with missing answers (n = 149) to the questions



*27 were older than 69 years at the time of health check, 674 had missing data on one or more variables of interest.

FIGURE 1: Flowchart describing the selection of study participants.

about ever being told by a doctor or nurse about having high blood pressure (n = 7), hypercholesterolemia (n = 114), or diabetes (n = 28) were treated as negative responses.

2.8. Statistical Analysis. The crude prevalence of hypertension, hypercholesterolemia, and DM, and the proportions of people aware of having these risk factors were presented with 95% confidence intervals (CI). Awareness estimates were also presented age- and sex-standardized [31]. Univariable and multivariable binary logistic regressions were used to assess associations between socio-demographic, lifestyle, and health characteristics with the awareness of each of the three risk factors. Finally, we estimated proportions (95% CI) of participants taking medication for any of the investigated three conditions, depending on their awareness of having these risk factors. Statistical analyses were performed using STATA V.17 (StataCorp, TX, USA).

2.9. *Ethical Approval.* The KYH study complied with the Declaration of Helsinki and was approved by the ethics committees of the London School of Hygiene & Tropical Medicine (approval number 8808, date 247/02/2015), Novosibirsk State Medical University (approval number 75, approval received 21/05/2015), the Institute of Preventative Medicine (no approval number; approval received 26/12/2014), Novosibirsk and the Northern State Medical University, Arkhangelsk (approval number 01/01-15, received 27/01/2015).

3. Results

The mean age of participants was 53.9 ± 9.7 years. The sample included 1512 residents of Novosibirsk and 2291 of Arkhangelsk. Fifty-eight percent had ascertained

Characteristics	Total sample $N = 3803$	Males N = 1591 Abs (%)	Females $N = 2212$
City of residence			
Årkhangelsk	2291 (60.2)	957 (60.1)	1334 (60.3)
Novosibirsk	1512 (39.8)	634 (39.9)	878 (39.7)
Age (years)			
35-44	825 (21.7)	314 (19.7)	511 (23.1)
45-54	1051 (27.6)	451 (28.4)	600 (27.1)
55-64	1251 (32.9)	542 (34.0)	709 (32.1)
65–69	676 (17.8)	284 (17.9)	392 (17.7)
Higher (university) education	1430 (37.6)	563 (35.4)	867 (39.2)
Household income ^a			
Low	720 (18.9)	263 (16.5)	457 (20.7)
Middle	2862 (75.3)	1219 (76.6)	1643 (74.3)
High	221 (5.8)	109 (6.9)	112 (5.0)
Smoking status			
Never	1886 (49.6)	404 (25.4)	1482 (67.0)
Former	951 (25.0)	586 (36.8)	365 (16.5)
Current	966 (25.4)	601 (37.8)	365 (16.5)
Hazardous drinking ^b	435 (11.4)	393 (24.7)	42 (1.9)
Obesity, BMI ≥30	1207 (31.7)	417 (26.2)	790 (35.7)
Self-reported CVD ^c	941 (24.7)	379 (23.8)	562 (25.4)
Ascertained risk factors ^d			
Hypertension	2206 (58.0)	1017 (63.9)	1189 (53.8)
Hypercholesterolemia	3171 (83.4)	1325 (83.3)	1846 (83.5)
Diabetes mellitus	329 (8.7)	122 (7.7)	207 (9.4)
Self-reported risk-factors			
Hypertension	2224 (58.5)	914 (57.5)	1310 (59.2)
Hypercholesterolemia	1686 (44.3)	563 (35.4)	1123 (50.8)
Diabetes mellitus	298 (7.8)	105 (6.6)	193 (8.7)
Medication use			
Antihypertensives ^e	1547 (40.7)	590 (37.1)	957 (43.3)
Lipid modifying agents ^f	367 (9.6)	162 (10.2)	205 (9.3)
Antidiabetics ^g	239 (6.3)	74 (4.5)	165 (7.5)

TABLE 1: Characteristics of the total study population and stratified by sex.

^aLow income was defined as difficulties to buy food or clothes, middle–large domestic appliances or new car, high-flat, house, or having no financial constraints; ^bAlcohol Use Disorders Identification Test (AUDIT) score ≥ 8 ; ^cEver diagnosed with angina, stroke, myocardial infarction, atrial fibrillation, or heart failure; ^dAscertained at the health check by objective measurements and/or data on medication use; ^eCodes C02, C03, C07, C08, C09 according to WHO anatomical therapeutic chemical classification (ATC); ^fCode C10 according to ATC; ^gCode A10 according to ATC.

hypertension, 83.4% had hypercholesterolemia, and 8.7% had DM (Table 1). Age- and sex-standardized to ESP2013, the prevalence of hypertension, hypercholesterolemia, and DM were 53.5%, 81.6%, and 7.1%, respectively. There were no significant differences between Novosibirsk and Arkhangelsk (Table 3).

3.1. Awareness. Age- and sex-standardized prevalence of the awareness of hypertension was 79.3% among all participants with hypertension, significantly higher in Arkhangelsk (81.4%) compared to Novosibirsk (74.7%) (Table 2). The age-standardized prevalence was higher in females compared to males in the total sample (84.0% vs. 74.6%), and there were comparable differences between sexes in both sites taken separately.

The age- and sex-standardized prevalence of the awareness of hypercholesterolemia, the most prevalent of the three risk factors under study, was 44.7% among all participants with this risk factor. Like with hypertension, the awareness of hypercholesterolemia was higher in Arkhangelsk (46.5%) than in Novosibirsk (41.6%), and the age-standardized estimates were higher in females compared to males in the total sample (51.4% vs. 38.0%), with similar differences observed in both sites.

For DM, the least prevalent risk factor under study, ageand sex-standardized prevalence of the awareness in the total sample of diabetic participants was 61.2%. There were no significant differences between the sites in the total sample, but females with diabetes in Novosibirsk had substantially higher awareness prevalence (81.0%) compared to their Arkhangelsk counterparts (51.0%). Comparisons of the agestandardized estimates between males and females showed no differences.

3.2. Associated Characteristics. In both univariable and multivariable regression analyses, the odds of being aware of having hypertension were higher among 55–69-year-old participants compared to 35–44-year-old participants, in females compared to males, in obese participants compared to the non-obese, and in those who self-reported CVD

		lotal study pop	ulation		Novosibirsk study	population		Arkhangelsk study	population
		Crude	Standardized		Crude	Standardized		Crude	Standardized
	N^{a}	estimates	estimates ^b	N^{a}	estimates	estimates ^a	N^{a}	estimates	estimates ^a
		Percent	(95% CI)		Percent	(95% CI)		Percent	(95% CI)
Total sample									
Hypertension	2206	83.6 (82.0; 85.1)	79.3 (76.9; 81.5)	890	81.4 (78.7; 83.8)	74.7 (70.4; 78.5)	1316	85.2 (83.2; 87.0)	$81.4 \ (78.4; \ 84.0)^{\dagger}$
Hypercholesterolemia	3171	50.3 (48.6; 52.1)	44.7 (42.9; 46.5)	1288	49.4(46.7; 52.1)	41.6 (38.9; 44.4)	1883	51.0 (48.7; 53.2)	$46.5 (44.2; 48.8)^{\dagger}$
Diabetes mellitus	329	73.9 (68.8; 78.3)	61.2 (53.4; 68.4)	148	77.0 (69.5; 83.1)	69.1 (60.5; 76.5)	181	71.3 (64.2; 77.4)	58.3 (50.2; 66.1)
Males									
Hypertension	1017	77.1 (74.4; 79.6)	74.6 (71.2; 77.7)	396	72.0 (67.3; 76.2)	65.5 (58.9; 71.5)	621	80.4 (77.0; 83.3)	$79.0(75.0; 82.4)^{\dagger}$
Hypercholesteremia	1325	40.2 (37.5; 42.8)	38.0(35.4;40.8)	532	38.5 (34.5; 42.8)	34.0 (29.9; 38.3)	793	41.2 (37.9; 44.7)	40.3 (36.8; 43.8)
Diabetes mellitus	122	72.1 (63.4; 79.4)	60.5 $(51.1; 69.1)$	58	70.7 (57.7; 81.0)	57.2 (43.5; 69.9)	64	73.4 (61.2; 82.9)	65.7 (55.2; 74.9)
Females									
Hypertension	1189	89.2 (87.3; 90.9)	$84.0 \ (80.6; \ 86.9)^{\ddagger}$	494	88.9 (85.8; 91.4)	$83.8 (78.2; 88.3)^{\ddagger}$	695	89.5 (87.0; 91.6)	83.8 (79.2; 87.5) [‡]
Hypercholesterolemia	1846	57.6(55.4; 59.9)	$51.4 (49.0; 53.7)^{\ddagger}$	756	57.0 (53.5; 60.5)	$49.3 \ (45.7; \ 52.9)^{\ddagger}$	1090	58.1 (55.1; 61.0)	$52.7 (49.7; 55.7)^{\ddagger}$
Diabetes mellitus	207	74.9 (68.5; 80.4)	$61.8 \ (49.3; 73.0)$	90	81.1 (71.6; 88.0)	81.0(70.4; 88.4)	117	70.1 (61.1; 77.7)	$51.0 (38.5; 63.3)^{\dagger}$

TABLE 2: Prevalence of the awareness of hypertension, hypercholesterolemia, and diabetes mellitus among study participants with corresponding risk factors in the total sample and with

International Journal of Hypertension

diagnoses compared to those who did not (Figure 2). Highincome participants had reduced odds of awareness compared to those with middle incomes, but only in the univariate analysis. This was not observed after mutual adjustment for other confounding variables. Current smokers consistently demonstrated the reduced odds of the awareness of hypertension in both analyses.

Univariable and multivariable regressions showed increased odds of being aware of having hypercholesterolemia among those age 45–69 years compared to 35–44 years, in females compared to males, in obese compared to nonobese, and if having self-reported CVD diagnoses. Smokers and hazardous drinkers demonstrated the reduced odds of the awareness in univariable analyses, but only smoking sustained the association after all the covariates were mutually adjusted in the multivariable model.

The odds of the awareness of having DM were increased in participants with low income compared to the middleincome group in both univariable and multivariable analyses, while those in the high-income group had the reduced odds of DM only in the adjusted analysis. In both analyses, the odds of being aware were also increased in participants who self-reported CVDs compared to those who did not. The exact odds ratios with respective confidence intervals are provided in supplementary file 1.

3.3. Risk Factor Control by Awareness Status. The proportion of participants with hypertension, hypercholesterolemia, and diabetes mellitus who self-reported relevant medication treatment was 71.1%, 11.6%, and 72.6%, respectively. These proportions showed disparities depending on participant's awareness status. Among the participants who were aware of their condition, a substantially higher proportion received a relevant medication to control hypertension (80.3%), hypercholesterolemia (20.4%), and DM (87.2%) compared to participants who were unaware (18.0%, 2.6%, and 31.4%, respectively) (Figure 3).

4. Discussion

Our study demonstrated that the standardized prevalence estimates for awareness of hypertension, hypercholesterolemia, and DM in Russian adults with corresponding conditions were 79.3%, 44.7%, and 61.2%, respectively. Depending on the risk factor considered, the awareness had independent positive associations with older age, female sex, low income, obesity, prior CVD diagnoses, and negative association with current smoking.

4.1. Prevalence of Awareness. With standardization, the prevalence of hypertension was 53.5% in the study population, and the overall awareness proportion in participants with this risk factor (79.3%) was higher compared to the other two risk factors under study. This awareness proportion was comparable to the latest published estimates in Russia (68% in males, 86% in females) [3], and it was higher than the estimates for France at 37.5% [15], Luxemburg at

40% [32], Denmark at 40% [33], Portugal at 46% [34], Italy at 56% [35], Czech Republic at 67% [36], Spain at 64% [37], Sweden at 65% [38], Poland at 67% [39], and England at 65% [40]. However, awareness was lower than in the Netherlands at 80% [41], Germany at 80% [42], and Greece at 90% [43].

The standardized prevalence of hypercholesterolemia (81.6%) was the highest in the study population among the investigated risk factors, but the awareness proportion was the lowest (44.7%). At the same time, it was higher than the other estimates for Russia (11% in males, 21% in females) [3] and the findings in Luxembourg (15%) [32], comparable to the estimates for the Czech Republic (40%) [3] and Italy (43%) [35], but lower than for Poland (51%) [44].

DM had the lowest standardized prevalence (7.1%) among the three studied risk factors, with an awareness proportion of 61.2%. The latter was within the range of earlier published estimates for Russia varying from 31% and 45% for men and females, respectively [3], to the total estimate of 73% [45], and it was close to the proportion of diabetics treated with sugar-lowering drugs (59.3%) observed in Russian population sample in 2015–2018 [46]. The standardized DM awareness proportion in our study (61.2%) was also higher compared to Czech republic 54% [3], similar to Switzerland (65%) [47], Luxembourg (68%) [32] but lower than in Italy (77%) [48], Poland (77%) [3], and Portugal 87% [49].

Even though our awareness estimates of hypercholesterolemia and DM were low and our awareness of hypertension quite high, they are within the variability range observed in the countries with lower CVD mortality rates. However, our cross-country comparisons must be interpreted cautiously because of the comparability limitations. For example, different definitions of hypercholesterolemia (total cholesterol \geq 4.9/5.0 and LDL \geq 2.9) were used in the Czech Republic and Italy [3, 32, 35]. The participants in studies from Germany [42], Greece [43], and the Netherlands [41] were generally older, ranging from 55 to 95 years of age, compared to participants in our study who were 35-69 years of age. Contrary to our study, Italian estimates [35] disregarded the medication use of participants. Despite methodological differences, we believe in the reliability of the comparisons. Thus, low CVD awareness rates cannot solely explain the higher CVD mortality in Russia [6].

4.2. Factors Associated with Awareness. In agreement with previous research [17, 50–53], our study showed that older people were more likely to be aware of hypertension, hypercholesterolemia, and DM compared to younger people. The discrepancy might be attributable to the lack of healthcare utilization due to the overall low disease prevalence among the young and the correspondingly lower probability of exposure to the risk information [54]. Our findings of the higher awareness of hypertension and hypercholesterolemia in females compared to males are also similar to those of earlier studies [15, 19, 20, 55–59]. This might be explained by female's higher engagement with healthcare services and accordingly their increased likelihood of being informed about any health risks [15].

International Journal of Hypertension



FIGURE 2: Adjusted and unadjusted odds ratios and confidence intervals for awareness of hypertension, hypercholesterolemia and diabetes mellitus by age, sex, education, income, smoking status, drinking status, BMI, and previous cvd experience. (a) Hypertension. (b) Hypercholesterolemia. (c) Diabetes mellitus. (d) Hypertension. (e) Hypercholesterolemia. (f) Diabetes mellitus.



FIGURE 3: Proportions of participants taking medication for hypertension, hypercholesterolemia, and diabetes mellitus depending on the awareness of having these risk factors.

The associations observed between socioeconomic factors and the awareness of risk factors appear inconsistent with prior research. Contrary to earlier associations between high income and awareness [3, 16, 60], our study found higher awareness of DM only among low-income participants. Despite the existing evidence of the higher awareness levels among those with university education [16, 61–65] and also the opposite findings in a French study [15], we found no association between awareness and education. These findings may indicate specific features of the studied population, including the equal accessibility of the healthcare services, regardless of the socioeconomic status.

Other researchers have described the associations of the awareness of CVD risk factors with alcohol drinking [3, 19, 52, 60, 63, 66] and smoking [19, 56, 59, 67]. Our findings were similar with respect to smoking, but hazardous drinking was associated with the awareness of hypertension and hypercholesterolemia only before the adjustments for

	Crude estimates	Standardized estimates ^a
	Percent (95% CI)	
Total sample		
Hypertension	58.0 (56.4; 59.6)	53.5 (52.0; 55.1)
Hypercholesterolemia	83.4 (82.2; 84.5)	81.6 (80.3; 82.9)
Diabetes mellitus	8.7 (7.8; 9.6)	7.1 (6.4; 7.9)
Novosibirsk		
Hypertension	58.9 (56.4; 61.3)	52.2 (49.6; 54.8)
Hypercholesteremia	85.2 (83.3; 86.9)	82.8 (80.6; 84.9)
Diabetes mellitus	9.8 (8.4; 11.4)	7.9 (6.6; 9.3)
Arkhangelsk		
Hypertension	57.4 (55.4; 59.5)	54.4 (52.5; 56.4)
Hypercholesterolemia	82.2 (80.6; 83.7)	80.9 (79.1; 82.5)
Diabetes mellitus	7.9 (6.9; 9.1)	6.7 (5.8; 7.7)

TABLE 3: Prevalence of hypertension, hypercholesterolemia, and diabetes mellitus in the total study population (N = 3803) and by study site (Novosibirsk, N = 1512; Arkhangelsk, N = 2291).

^aAge- and sex-standardized to European standard population 2013 (ESP2013).

smoking and other covariates. This may be explained by a known association between smoking and hazardous drinking—the commonly cohabiting components of an unhealthy lifestyle. The adjustment for smoking may have excessively attenuated the association between the awareness and hazardous drinking. Based on this assumption, the overall negative effect of the unhealthy lifestyle on the awareness could be inferable.

With respect to obesity and previous cardiovascular events, our findings agree with the existing evidence that living with these conditions increases awareness of hypertension, hypercholesterolemia, and DM [15, 18, 20, 64, 68–70], which may be explained by higher health concerns and frequent health care contacts.

Finally, our study has demonstrated that those aware of their risk factors were more likely to be receiving proper medication compared to those unaware. Notably, our findings reveal a disparity between the awareness rates (44.7%) and the treatment rates for hypercholesterolemia, which were relatively low (11.6%). Conversely, participants with hypertension and diabetes mellitus (DM) displayed comparatively higher rates of treatment. Despite our identification of low treatment rates for hypercholesterolemia and relatively favorable treatment rates for hypertension, prior studies report extremely low antihypertensive and lipid-lowering medication adherence in Russia [71]. On the other hand, self-medication for high blood pressure without having hypertension diagnosed by a doctor is an acknowledged problem in Russia because of the non-prescription sales of antihypertensives [71]. As for antidiabetics, their intake by those not aware of having DM might be explained by the common prescription of glucose-lowering medication to those at the so-called prediabetes stage [72]. The role of awareness in a Russian setting is therefore crucial not only in recognising the risk of hypertension, hypercholesterolemia, and DM but also in treatment, adherence, control, and overall prevention of unnecessary self-medication.

4.3. Strengths and Limitations. A strength of this study is that we investigated awareness of three major CVD risk factors using data from a large population-based sample of Russian adults, where self-reports could be assessed against the objective presence of the same risk factors, as defined by examinations performed by health professionals. However, the relatively low prevalence of DM resulted in a rather low precision of the awareness estimates and the power to identify the associated characteristics.

An important limitation is that the KYH sample is comprised of only urban residents of two Russian cities, which may not be representative of the Russian population overall. Urban residents have better access to healthcare, which might lead to an overestimation of the presented awareness proportions [3]. Therefore, the generalizability of the findings is limited to the Russian urban population.

The overall response proportion in the KYH was relatively low and varied between the cities 68% in Arkhangelsk and 41% in Novosibirsk [28]. We assessed nonresponse bias by comparing the risk factors and the awareness proportions between the sites which resulted in minor differences (Table 3) indirectly indicating that selection bias was not substantial. However, assessments do not exclude the possibility of selection bias for the whole. KYH was presented as a study focused on cardiovascular health, and people more concerned with their health may have been more likely to participate in it.

The ascertainment of the risk factors included using the self-reported data on medication use, which could limit the objectivity. However, we believe the bias could not be large since participants were asked to show their regularly taken medications or to list their commercial names, doses, and frequencies. Antihypertensives, lipid-lowering drugs, and antidiabetics require long-time daily intake, so failure to remember their names was unlikely with adherence. In addition, several studies show a good agreement between self-reported CVD medication and pharmacy records [73, 74].

5. Conclusion

Awareness of hypertension is relatively high, but for high cholesterol and diabetes mellitus awareness is relatively low. Lower awareness may lead to lower levels of preventative treatment for key CVD risk factors which contribute to higher CVD morbidity and mortality in Russia. We observed that awareness levels varied by age, sex, income, smoking status, obesity, and comorbidities. These findings may guide targeted awareness-raising interventions.

Data Availability

The data used to support the findings of this study may be released upon application on the website https://metadata. knowyourheart.science after contacting David Leon at david.leon@lshtm.ac.uk.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Acknowledgments

The authors would like to acknowledge The Know Your Heart (KYH) study leadership for granting the free of charge data access. The authors also acknowledge the KYH study participants for their time and effort. The Know Your Heart (KYH) study was a component of the International Project on Cardiovascular Disease in Russia (IPCDR) funded by Wellcome Trust Strategic Award (100217), UiT The Arctic University of Norway (UiT), Norwegian Institute of Public Health, and Norwegian Ministry of Health and Social Affairs. MS and MSh were supported by the Russian Academy of Science, FWNR-2024-0002.

Supplementary Materials

Supplementary file 1: adjusted and unadjusted odds ratios and confidence intervals for awareness of hypertension, hypercholesterolemia, and diabetes mellitus by age, sex, education, income, smoking status, drinking status, BMI, and previous cvd experience. (*Supplementary Materials*)

References

- World Health Organisation, "Cardiovascular diseases," 2021, https://www.who.int/health-topics/cardiovascular-diseases#tab= tab_1.
- [2] G. A. Roth, G. A. Mensah, C. O. Johnson et al., "Global burden of cardiovascular diseases and risk factors, 1990-2019: update from the GBD 2019 study," *Journal of the American College of Cardiology*, vol. 76, no. 25, pp. 2982–3021, 2020.
- [3] W. Lu, H. Pikhart, A. Tamosiunas et al., "Prevalence, awareness, treatment and control of hypertension, diabetes and hypercholesterolemia, and associated risk factors in the Czech Republic, Russia, Poland and Lithuania: a crosssectional study," *BMC Public Health*, vol. 22, no. 1, p. 883, 2022.

- [4] Eurostat, "WHO mortality database," 2021, https://ec.europa.eu/ eurostat/statistics-explained/index.php?title=Cardiovascular_dis eases_statistics#Deaths_from_cardiovascular_diseases.
- [5] Rossat, "Russia in numbers," 2020, https://dmitriypushin.ru/ wp-content/uploads/2021/02/Spravochnik-Rossiya-v-tsifrah-2020_compressed.pdf.
- [6] World Health Organisation, WHO Mortality Database, WHO, Geneva, Switzerland, 2022.
- [7] S. Trias-Llimos, L. Pennells, A. Tverdal et al., "Quantifying the contribution of established risk factors to cardiovascular mortality differences between Russia and Norway," *Scientific Reports*, vol. 10, no. 1, Article ID 20796, 2020.
- [8] O. Lakunchykova, M. Averina, T. Wilsgaard et al., "Why does Russia have such high cardiovascular mortality rates? Comparisons of blood-based biomarkers with Norway implicate non-ischaemic cardiac damage," *Journal of Epidemiology & Community Health*, vol. 74, no. 9, pp. 698–704, 2020.
- [9] D. Boateng, F. Wekesah, J. L. Browne et al., "Knowledge and awareness of and perception towards cardiovascular disease risk in sub-Saharan Africa: a systematic review," *PLoS One*, vol. 12, no. 12, Article ID e0189264, 2017.
- [10] Centers for Disease Control and Prevention, "Heart disease and stroke," 2023, https://www.betterhealth.vic.gov.au/ health/conditionsandtreatments/heart-disease-and-stroke.
- [11] National Health Services, "What is high cholesterol?" 2019, https://www.nhs.uk/conditions/high-cholesterol/.
- [12] M. Clinic, "High blood pressure," 2020, https://www. mayoclinic.org/diseases-conditions/high-blood-pressure/ symptoms-causes/syc-20373410.
- [13] National Health Services, "Diabetes," 2019, https://www.nhs. uk/conditions/diabetes/.
- [14] S. G. Chrysant, "A new paradigm in the treatment of the cardiovascular disease continuum: focus on prevention," *Hippokratia*, vol. 15, no. 1, pp. 7–11, 2011.
- [15] G. Fenech, A. Vallee, M. Cherfan et al., "Poor awareness of hypertension in France: the CONSTANCES population-based study," *American Journal of Hypertension*, vol. 33, no. 6, pp. 543–551, 2020.
- [16] U. G. Sendur and M. Adas, "Determinants of awareness on diabetes and its complications," *Experimental and Clinical Endocrinology & Diabetes*, vol. 129, no. 04, pp. 269–275, 2021.
- [17] O. A. Sanuade, R. B. Awuah, and M. Kushitor, "Hypertension awareness, treatment and control in Ghana: a cross-sectional study," *Ethnicity and Health*, vol. 25, no. 5, pp. 702–716, 2020.
- [18] A. Oraii, A. Shafiee, A. Jalali, F. Alaeddini, S. Saadat, and F. Masoudkabir, "Prevalence, awareness, treatment, and control of type 2 diabetes mellitus among the adult residents of tehran: tehran Cohort Study," *BMC Endocrine Disorders*, vol. 22, no. 1, p. 248, 2022.
- [19] C. Wang, Y. Yu, X. Zhang et al., "Awareness, treatment, control of diabetes mellitus and the risk factors: survey results from northeast China," *PLoS One*, vol. 9, no. 7, Article ID e103594, 2014.
- [20] S. T. Yen, A. K. G. Tan, and F. I. Mustapha, "Awareness of diabetes, hypertension, and hypercholesterolemia in Malaysia," *Journal of Diabetes*, vol. 9, no. 9, pp. 874–883, 2017.
- [21] S. A. Boytsov, S. A. Shalnova, A. D. Deev et al., "Arterial hypertension among individuals of 25–64 years old: prevalence, awareness, treatment and control. BY the data from eccd," *Cardiovascular- Therapy and Prevention*, vol. 13, no. 4, pp. 4–14 2014, 2014.
- [22] P. M. Grinshtein YuI, V. V. Shabalin, R. R. Ruf et al., "The prevalence of arterial hypertension in Krasnoyarsky territory:

the data from the epidemiological study ESSE-RF," Arterial'naya Gipertenziya (Arterial Hypertension), vol. 52, 2016.

- [23] Y. A. Balanova, S. A. Shalnova, A. E. Imaeva et al., "Prevalence, awareness, treatment and control of hypertension in Russian federation (data of observational ESSERF-2 study)," *Rational Pharmacotherapy in Cardiology*, vol. 15, no. 4, pp. 450–466, 2019.
- [24] M. Cybulsky, S. Cook, A. V. Kontsevaya, M. Vasiljev, and D. A. Leon, "Pharmacological treatment of hypertension and hyperlipidemia in Izhevsk, Russia," *BMC Cardiovascular Disorders*, vol. 16, no. 1, p. 122, 2016.
- [25] S. K. Malyutina, E. V. Mazdorova, M. Y. Shapkina et al., "[The frequency and profile of drug treatment in subjects with dyslipidemias and cardimetabolic diseases in an urban Russian population older then 55 years]," *Kardiologiia*, vol. 61, no. 12, pp. 49–58, 2021.
- [26] M. Shapkina, A. Ryabikov, E. Mazdorova et al., "The determinants of the 13-year risk of incident atrial fibrillation in a Russian population cohort of middle and elderly age," *Journal of Personalized Medicine*, vol. 12, no. 1, p. 122, 2022.
- [27] S. K. Malyutina, E. V. Mazdorova, M. Y. Shapkina et al., "[The profile of drug treatment in subjects aged over 50 years with hypertension in an urban Russian population]," *Kardiologiia*, vol. 60, no. 3, pp. 21–29, 2020.
- [28] S. Cook, S. Malyutina, A. V. Kudryavtsev et al., "Know Your Heart: rationale, design and conduct of a cross-sectional study of cardiovascular structure, function and risk factors in 4500 men and women aged 35-69 years from two Russian cities, 2015-18," *Wellcome Open Res*, vol. 3, p. 67, 2018.
- [29] World Health Organisation, "ATC/DDD index," 2023, https://www.whocc.no/atc_ddd_index/.
- [30] F. Babor Thomas, J. Higgins-Biddle, J. Saunders, and M. Mastela, AUDIT, World Health Organisation, Geneva, Switzerland, 2001.
- [31] Eurostat, "Revision of the European standard population- report of eurostat's task force- 2013," 2013, https://ec.europa.eu/eurostat/ web/products-manuals-and-guidelines/-/ks-ra-13-028.
- [32] A. Alkerwi, S. Pagny, M. L. Lair, C. Delagardelle, and J. Beissel, "Level of unawareness and management of diabetes, hypertension, and dyslipidemia among adults in Luxembourg: findings from ORISCAV-LUX study," *PLoS One*, vol. 8, no. 3, Article ID e57920, 2013.
- [33] T. Sehestedt, H. Ibsen, and T. Jorgensen, "Awareness, treatment and control of hypertension in Denmark. The Inter99 study," *Blood Pressure*, vol. 16, no. 5, pp. 312–319, 2007.
- [34] M. E. De Macedo, M. J. Lima, A. O. Silva, P. Alcantara, V. Ramalhinho, and J. Carmona, "Prevalence, awareness, treatment and control of hypertension in Portugal. The PAP study," *Revista Portuguesa de Cardiologia*, vol. 26, no. 1, pp. 21–39, 2007.
- [35] S. Omboni, G. Carabelli, E. Ghirardi, and S. Carugo, "Awareness, treatment, and control of major cardiovascular risk factors in a small-scale Italian community: results of a screening campaign," *Vascular Health and Risk Management*, vol. 9, pp. 177–185, 2013.
- [36] R. Cifkova, J. Bruthans, L. Strilchuk et al., "Longitudinal trends in blood pressure, prevalence, awareness, treatment, and control of hypertension in the Czech population. Are there any sex differences?" *Frontiers in Cardiovascular Medicine*, vol. 9, Article ID 1033606, 2022.
- [37] J. R. Banegas, P. Guallar-Castillon, F. Rodriguez-Artalejo, A. Graciani, E. Lopez-Garcia, and L. M. Ruilope, "Association between awareness, treatment, and control of hypertension,

and quality of life among older adults in Spain," *American Journal of Hypertension*, vol. 19, no. 7, pp. 686–693, 2006.

- [38] A. Santosa, Y. Zhang, L. Weinehall et al., "Gender differences and determinants of prevalence, awareness, treatment and control of hypertension among adults in China and Sweden," *BMC Public Health*, vol. 20, no. 1, p. 1763, 2020.
- [39] T. Zdrojewski, B. Wyrzykowski, R. Szczech et al., "Epidemiology and prevention of arterial hypertension in Poland," *Blood Pressure*, vol. 14, no. 2, pp. 10–16, 2005.
- [40] M. Joffres, E. Falaschetti, C. Gillespie et al., "Hypertension prevalence, awareness, treatment and control in national surveys from England, the USA and Canada, and correlation with stroke and ischaemic heart disease mortality: a crosssectional study," *BMJ Open*, vol. 3, no. 8, Article ID e003423, 2013.
- [41] C. T. van Rossum, H. van de Mheen, J. C. Witteman, A. Hofman, J. P. Mackenbach, and D. E. Grobbee, "Prevalence, treatment, and control of hypertension by sociodemographic factors among the Dutch elderly," *Hypertension*, vol. 35, no. 3, pp. 814–821, 2000.
- [42] S. Muli, C. Meisinger, M. Heier, B. Thorand, A. Peters, and U. Amann, "Prevalence, awareness, treatment, and control of hypertension in older people: results from the populationbased KORA-age 1 study," *BMC Public Health*, vol. 20, no. 1, p. 1049, 2020.
- [43] A. Triantafyllou, S. Douma, K. Petidis et al., "Prevalence, awareness, treatment and control of hypertension in an elderly population in Greece," *Rural and Remote Health*, vol. 10, no. 2, p. 1225, 2010.
- [44] T. Zdrojewski, B. Solnica, B. Cybulska et al., "Prevalence of lipid abnormalities in Poland. The NATPOL 2011 survey," *Kardiologia Polska*, vol. 74, no. 3, pp. 213–223, 2016.
- [45] M. M. Bikbov, R. R. Fayzrakhmanov, G. M. Kazakbaeva et al., "Prevalence, awareness and control of diabetes in Russia: the Ural Eye and Medical Study on adults aged 40+ years," *PLoS One*, vol. 14, no. 4, Article ID e0215636, 2019.
- [46] S. Malyutina, E. Mazurenko, E. Mazdorova et al., "The profile of glucose lowering therapy in persons with type 2 diabetes mellitus in an aging Russian population," *Journal of Personalized Medicine*, vol. 12, no. 10, p. 1689, 2022.
- [47] A. Kaiser, P. Vollenweider, G. Waeber, and P. Marques-Vidal, "Prevalence, awareness and treatment of type 2 diabetes mellitus in Switzerland: the CoLaus study," *Diabetic Medicine*, vol. 29, no. 2, pp. 190–197, 2012.
- [48] A. Gnasso, M. C. Calindro, C. Carallo et al., "Awareness, treatment and control of hyperlipidaemia, hypertension and diabetes mellitus in a selected population of southern Italy," *European Journal of Epidemiology*, vol. 13, no. 4, pp. 421–428, 1997.
- [49] M. Barreto, I. Kislaya, V. Gaio et al., "Prevalence, awareness, treatment and control of diabetes in Portugal: results from the first National Health examination Survey (INSEF 2015)," *Diabetes Research and Clinical Practice*, vol. 140, pp. 271–278, 2018.
- [50] C. Dong, P. Ge, X. Ren, H. Fan, and X. Yan, "Prevalence, awareness, treatment and control of hypertension among adults in rural north-western China: a cross-sectional population survey," *Journal of International Medical Research*, vol. 41, no. 4, pp. 1291–1300, 2013.
- [51] F. O'Brien, P. McCallion, R. Carroll, M. O'Dwyer, E. Burke, and M. McCarron, "The prevalence, awareness, treatment, and control of hypertension in older adults with an intellectual disability in Ireland: a cross sectional study,"

European Journal of Cardiovascular Nursing, vol. 20, no. 4, pp. 315–323, 2021.

- [52] M. Pereira, A. Azevedo, and H. Barros, "Determinants of awareness, treatment and control of hypertension in a Portuguese population," *Revista Portuguesa de Cardiologia*, vol. 29, no. 12, pp. 1779–1792, 2010.
- [53] S. Shirani, R. Kelishadi, N. Sarrafzadegan et al., "Awareness, treatment and control of hypertension, dyslipidaemia and diabetes mellitus in an Iranian population: the IHHP study," *Eastern Mediterranean Health Journal*, vol. 15, no. 6, pp. 1455–1463, 2009.
- [54] B. Everett and A. Zajacova, "Gender differences in hypertension and hypertension awareness among young adults," *Biodemography and Social Biology*, vol. 61, no. 1, pp. 1–17, 2015.
- [55] C. K. Chow, K. K. Teo, S. Rangarajan, S. Islam, R. Gupta, and A. Avezum, "Prevalence, awareness, treatment, and control of hypertension in rural and urban communities in high-middle-and low-income countries," *JAMA*, vol. 310, no. 9, pp. 959–968, 2013.
- [56] L. Shahab, J. Mindell, N. R. Poulter, and R. West, "Hypertension and its identification among current, past and never smokers in an english population sample," *European Journal* of Cardiovascular Prevention & Rehabilitation, vol. 17, no. 1, pp. 63–70, 2010.
- [57] J. L. Wellman, B. Holmes, and S. Y. Hill, "Accuracy of selfreported hypertension: effect of age, gender, and history of alcohol dependence," *Journal of Clinical Hypertension*, vol. 22, no. 5, pp. 842–849, 2020.
- [58] G. S. Stergiou, A. Menti, N. Kalpourtzi et al., "Prevalence, awareness, treatment and control of hypertension in Greece: EMENO national epidemiological study," *Journal of Hypertension*, vol. 39, no. 5, pp. 1034–1039, 2021.
- [59] R. E. K. Man, A. H. W. Gan, E. K. Fenwick et al., "Prevalence, determinants and association of unawareness of diabetes, hypertension and hypercholesterolemia with poor disease control in a multi-ethnic Asian population without cardiovascular disease," *Population Health Metrics*, vol. 17, no. 1, p. 17, 2019.
- [60] J. Lu, Y. Lu, X. Wang et al., "Prevalence, awareness, treatment, and control of hypertension in China: data from 1.7 million adults in a population-based screening study (China PEACE Million Persons Project)," *The Lancet*, vol. 390, no. 10112, pp. 2549–2558, 2017.
- [61] Q. Cao, P. Pei, J. Zhang et al., "Hypertension unawareness among Chinese patients with first-ever stroke," *BMC Public Health*, vol. 16, no. 1, p. 170, 2016.
- [62] M. A. Foma, Y. Saidu, S. A. Omoleke, and J. Jafali, "Awareness of diabetes mellitus among diabetic patients in the Gambia: a strong case for health education and promotion," *BMC Public Health*, vol. 13, no. 1, p. 1124, 2013.
- [63] X. Liu, W. Gu, Z. Li, H. Lei, G. Li, and W. Huang, "Hypertension prevalence, awareness, treatment, control, and associated factors in Southwest China: an update," *Journal of Hypertension*, vol. 35, no. 3, pp. 637–644, 2017.
- [64] Y. Okura, L. H. Urban, D. W. Mahoney, S. J. Jacobsen, and R. J. Rodeheffer, "Agreement between self-report questionnaires and medical record data was substantial for diabetes, hypertension, myocardial infarction and stroke but not for heart failure," *Journal of Clinical Epidemiology*, vol. 57, no. 10, pp. 1096–1103, 2004.
- [65] A. Supiyev, T. Nurgozhin, Z. Zhumadilov, A. Peasey, J. A. Hubacek, and M. Bobak, "Prevalence, awareness, treatment and control of dyslipidemia in older persons in

urban and rural population in the Astana region, Kazakhstan," *BMC Public Health*, vol. 17, no. 1, p. 651, 2017.

- [66] H. He, Y. Q. Yu, Y. Li et al., "Dyslipidemia awareness, treatment, control and influence factors among adults in the Jilin province in China: a cross-sectional study," *Lipids in Health and Disease*, vol. 13, no. 1, p. 122, 2014.
- [67] P. Mathur, S. Leburu, and V. Kulothungan, "Prevalence, awareness, treatment and control of diabetes in India from the countrywide national NCD monitoring survey," *Frontiers in Public Health*, vol. 10, Article ID 748157, 2022.
- [68] I. S. Okosun and G. A. Dever, "Abdominal obesity and ethnic differences in diabetes awareness, treatment, and glycemic control," *Obesity Research*, vol. 10, no. 12, pp. 1241–1250, 2002.
- [69] R. Safari-Faramani, F. Rajati, K. Tavakol et al., "Prevalence, awareness, treatment, control, and the associated factors of diabetes in an Iranian Kurdish population," *Journal of Diabetes Research*, vol. 2019, Article ID 5869206, 9 pages, 2019.
- [70] X. Liu, S. Yu, Z. Mao et al., "Dyslipidemia prevalence, awareness, treatment, control, and risk factors in Chinese rural population: the Henan rural cohort study," *Lipids in Health and Disease*, vol. 17, no. 1, p. 119, 2018.
- [71] S. Cook, L. A. Hopstock, A. E. Eggen et al., "Pharmacological management of modifiable cardiovascular risk factors (blood pressure and lipids) following diagnosis of myocardial infarction, stroke and diabetes: comparison between population-based studies in Russia and Norway," BMC Cardiovascular Disorders, vol. 20, no. 1, p. 234, 2020.
- [72] P. Mukhopadhyay and S. Chowdhury, "Drug therapy in prediabetes," *Journal of the Indian Medical Association*, vol. 103, no. 11, pp. 603–608, 2005.
- [73] O. H. Klungel, A. de Boer, A. H. Paes, R. M. Herings, J. C. Seidell, and A. Bakker, "Agreement between self-reported antihypertensive drug use and pharmacy records in a population-based study in The Netherlands," *Pharmacy World and Science: PWS*, vol. 21, no. 5, pp. 217–220, 1999.
- [74] J. D. Hafferty, A. I. Campbell, L. B. Navrady et al., "Selfreported medication use validated through record linkage to national prescribing data," *Journal of Clinical Epidemiology*, vol. 94, pp. 132–142, 2018.