

Research Article

Prevalence and Antimicrobial Resistance of Bacteria Isolated from Marine and Freshwater Fish in Tanzania

Esther Marijani 

Open University of Tanzania, P.O. Box 23409, Dar es Salaam, Tanzania

Correspondence should be addressed to Esther Marijani; esthermarijani@gmail.com

Received 16 September 2021; Revised 6 December 2021; Accepted 14 February 2022; Published 4 March 2022

Academic Editor: Clemencia Chaves Lopez

Copyright © 2022 Esther Marijani. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This study aimed to determine the prevalence and antimicrobial resistance of bacteria isolated from retail fish and shrimp in Tanzania. A total of 92 fish and 20 shrimp samples were analyzed. Fish samples consisted of 24 Nile tilapia, 24 Nile perch, and 24 red snapper. The isolates were identified by their morphological characteristics, conventional biochemical tests, and analytical profile index test kits. The antibiotic susceptibility of selected bacteria was determined by the disc diffusion method. Out of the 92 samples analyzed, 96.7% were contaminated with 7 different bacterial species. *E. coli* was the most prevalent bacteria (39%), followed by *Klebsiella* spp. (28%) and *Salmonella* spp. (16%). Other species isolated from this study were *Staphylococcus* spp. (8%), *Citrobacter* (4%), *Shigella* spp. (3%), and *Pseudomonas* spp. (1%). All samples were analyzed for *Campylobacter* spp.; however, none of the samples tested were positive for *Campylobacter* spp. Fish from the open-air market were contaminated by six bacterial species: *E. coli* (40%), *Klebsiella* spp. (26%), *Salmonella* spp. (24%), *Shigella* spp. (6.7%), *Citrobacter* spp. (6.5%), and *Pseudomonas* spp. (2%), while *E. coli* (37%), *Klebsiella* spp. (33%), *Staphylococcus* spp. (23%), and *Shigella* spp. (2%) were isolated in supermarket samples. According to the International Commission on Microbiological Specifications for Foods criteria, 54 (58.7%) and 38 (41.3%) samples were good and marginally acceptable, respectively. *E. coli* isolates were resistant to penicillin (PEN), erythromycin (ERY), gentamicin (GEN), azithromycin (AZM), and tetracycline (TET), while *Salmonella* spp. isolates exhibited resistance to gentamicin (CN), tetracycline (TET), penicillin (PEN), and erythromycin (ERY). These results suggest that the presence of these bacteria might cause a health risk/hazard to human beings and may cause disease to susceptible individuals, especially immune-compromised consumers.

1. Introduction

Fish are a vital source of food for people and are the most important source of high-quality protein. Fish provide approximately 16% of the animal protein consumed by the world's population [1]. Approximately 60% of developing countries derive 30% of their annual protein from fish [2]. Fish provide 19% of protein and are one of the cheapest sources of protein in Africa. In Tanzania, per capita, fish consumption is 8.0 kilograms, and fish make up 19.7% of the country's animal protein intake [3]. However, fish consumption is much higher in coastal regions and Zanzibar (almost 20 kg per capita), as well as along the coasts of Lake Tanganyika and Lake Victoria. Apart from being a rich source of protein, fish also play a unique role in providing a range of micronutrients and essential fatty acids, especially

long-chain polyunsaturated fatty acids, which cannot be easily substituted by other food commodities [4, 5]. Despite these numerous advantages of fish, they are susceptible to a wide variety of bacterial pathogens, most of which can cause disease and are considered by some to be saprophytic in nature [1]. Fish and fishery products are known to be sources of transmitting foodborne infections and intoxication if not properly handled/processed [6].

It has been reported that the microbiological diversity of fresh fish muscle depends on the fishing grounds and environmental factors around it [7]. According to Clucas and Ward [8], the type of microorganisms that are associated with a particular fish depends on their habitat.

Bacterial pathogens associated with fish are classified as indigenous and nonindigenous. Examples of nonindigenous bacterial pathogens include *Escherichia coli*, *Clostridium*

botulinum, *Shigella dysenteriae*, *Staphylococcus aureus*, *Listeria monocytogenes*, and *Salmonella*. These nonindigenous organisms contaminate the fish or the habitat in one way or the other. Examples of indigenous bacterial pathogens are *Vibrio* species and *Aeromonas* species, which are found naturally living in fish habitats [9, 10].

The bacteria from fish only become pathogens when there are stressors, such as poor water quality and overstocking, which allow opportunistic bacterial infections to prevail [1]. Additionally, studies have reported the presence of indicator microorganisms of fecal contamination, opportunistic, and pathogenic bacteria to humans in fish samples [11, 12].

Human infections caused by pathogens transmitted from fish are widespread and depend on the season, the patients' contact with fish and related environments, dietary habits, and the immune system status of the individual [13]. Pathogens can be transmitted through food or the handling of fish. Foodborne illnesses such as dysentery and diarrhea resulting from the consumption of contaminated fish can result in economic losses. Microbial associations with fish compromise the safety and quality of human consumption; this is particularly critical when microorganisms are opportunistic and/or pathogenic in nature [11, 13].

Poor fish handling, processing, and packaging methods and practices in retail markets are still common in Tanzania [14]. There are limited studies on the prevalence of bacteria isolated from retail fish and shrimps in Tanzania. The risk of getting foodborne diseases or food poisoning by fish consumers and handlers may be high in Tanzania. Therefore, this study investigates the prevalence of bacteria isolated from retail fish and shrimps in Tanzania.

Several antibiotics used in agriculture and aquaculture are critical for human medicine by the World Health Organization (WHO), including the antibiotic classes of tetracyclines, quinolones, and penicillin [15].

Resistance to all antibiotic classes has been observed in a wide range of bacteria, including those pathogenic to humans [16, 17].

Antimicrobial resistance is widespread in Tanzania, and there is widespread misuse of antimicrobials in the livestock and aquaculture industries. However, there is a scarcity of information regarding antibiotic resistance in Tanzania [18]. The scarcity of data may have contributed to the use of drugs without a prescription or prescribing drugs without carrying out a laboratory test to identify the pathogens and test their susceptibility to drugs for effective treatment.

Hence, another goal of this study was to determine the prevalence and antimicrobial resistance of foodborne pathogens isolated from retail fish and shrimps in Tanzania.

2. Methodology

2.1. Sample Collection. A total of 92 fresh fish samples of both marine and freshwater were purchased from two different open-air fish markets ($n=60$) and two supermarkets ($n=32$) in the Ilala district, Dar es Salaam, Tanzania. Out of the 92 samples collected, twenty-four samples were of Nile tilapia, Nile perch, red snapper, and twenty

shrimp. After the collection, samples were aseptically and immediately transported in a sterile polypropylene bag placed in a cooler box. The cooler box contained crushed ice, and the temperatures were between 4°C and 8°C during transportation. The samples were transported to the Open University Laboratory, and for further analysis, the samples were analyzed at the College of Veterinary and Biomedical Sciences, Sokoine University of Agriculture.

2.2. Culture, Isolation, and Identification. For isolation and identification of bacteria, culturing was performed by chopping a piece of fish flesh aseptically and spreading it over the surface of blood agar and MacConkey agar before incubation between 24 and 48 hours at 37°C. For *Campylobacter* spp. identification, a piece of fish samples was first enriched in Bolton broth overnight and later inoculated onto modified charcoal cefoperazone deoxycholate agar (mCCDA) (Oxoid Ltd., Basingstoke, Hampshire, England) containing the *Campylobacter* mCCDA selective supplement SR155 E (Oxoid Ltd., Basingstoke, Hampshire, England). Incubation was performed as previously described by Kurekci et al. [19] at 37°C for 48 h under microaerophilic conditions generated by CampyGen™ gas sachets (Oxoid, Basingstoke, England, UK).

Gram staining was done to differentiate organisms based on the structure of their cell walls [20]. Furthermore, the classical identification of bacterial colonies and biotyping were performed according to the methods described by Abbott et al. [20] and El Deen et al. [21] with slight modifications. Briefly, the isolates were conventionally studied for their macro- and micromorphological characteristics and then by biochemical assays that consisted of 21 phenotypic characteristics tests. The assays included lactose, raffinose, trehalose, dulcitol, maltose, mannose, D-mannitol, melibiose, sucrose, citrate, urea, indole, catalase, motility, ampicillin resistance, m-inositol, oxidase, nitrate, cellobiose, and xylose. Triple sugar iron agar and IMViC were also used for the characterization of members of the family Enterobacteriaceae.

2.3. Microbial Load by Total Viable Count. To assess the level of microbial contamination, 10 g of fish pieces was rinsed in 10 mL of sterile normal saline. Then, 1 mL of the rinse was serially diluted 10-fold using 10 universal bottles containing sterile normal saline. In each dilution, 1 mL was poured on the plate count of medium Petri dishes in duplicate. The plates were incubated at 37°C for 24 h. Then, colonies were counted, and the average colony counts were used to establish the colony-forming units (CFU/mL or CFU/g).

2.4. Antimicrobial Assay. Antimicrobial susceptibility testing was performed using the disc diffusion method. The isolates were tested against quinolones, namely, ciprofloxacin (CIP); macrolides, erythromycin (E), and azithromycin (AZM); and aminoglycosides, gentamicin (CN) (0.06–64 µg/mL), and tetracycline (TE). Other antimicrobials were clindamycin (DA) and penicillin (P). All

antimicrobials were supplied by Sigma-Aldrich (St. Louis, MO, USA). For antimicrobial susceptibility assays, a pool of bacterial colonies was used to prepare suspensions corresponding to 0.5 McFarland standards (1.5×10^8 CFU/mL) using normal saline, and then bacteria were spread on top of Müller-Hinton agar using a sterile swab. Discs were placed on top of the medium, and the plates were then incubated at 37°C for 24 h. Zones of inhibition were measured by means of a simple ruler, and the diameter was recorded in millimeters (mm).

Isolates were defined as susceptible, intermediate, or resistant in accordance with the CLSI [22] *Enterobacteriaceae* breakpoints.

2.5. Data Analysis. Data from counts (in log CFU/g) were analyzed (means, standard deviations, and standard errors) using the Statistical Package SPSS v21 (IBM Corporation, Armonk, NY, USA). Descriptive statistics were used to describe the results of the prevalence analysis. The prevalence was estimated as the number of samples detected positive for *Salmonella* spp., *Campylobacter* spp., *E. coli*, *Shigella* spp., and *Staphylococcus* spp. isolation from the total sample analyzed.

3. Results

3.1. Total Viable Count (TVC) and Microbiological Quality Data. The average TVC for the 92 samples was 4.57 log CFU/g, ranging from 3.82 to 4.70 and 5.22 log CFU/g in tilapia, Nile perch, and red snapper, respectively. A significant difference ($P < 0.05$) was observed between Nile tilapia and red snapper samples, as shown in Table 1. When comparing total samples (Table 1), red snapper had the highest mean TVC at 5.22 log CFU/g, and Nile tilapia had the lowest at 3.82 log CFU/g ($P < 0.05$). There is no significant difference ($P > 0.05$) in TVC log CFU/g between samples from the supermarket and retail market. However, samples from the supermarket have a lower TVC at 3.77 log CFU/g compared to samples from the open-air market (4.99 log CFU/g) (data not shown).

Based on the International Commission on Microbiological Specifications for Foods criteria, 54 (58.7%) and 38 (41.3%) samples were good and marginally acceptable, respectively (Table 1). Nile tilapia had a higher good quality sample, followed by shrimp, Nile perch, and red snapper. None of the samples had unacceptable microbiological quality.

3.2. Prevalence of *Campylobacter*, *E. coli*, *Listeria*, *Salmonella*, and *Shigella*. The prevalence and contamination rates of open-air market fish with pathogenic bacteria are presented in Table 2). Seven bacterial species were identified from all analyzed samples. The most prevalent bacterial species recovered in the samples was *E. coli* (39%), followed by *Klebsiella* spp. (28%) and *Salmonella* spp. (16%). Other species isolated from this study were *Staphylococcus* spp. (8%), *Citrobacter* (4%), *Shigella* spp. (3%), and *Pseudomonas* spp. (1%). Nile tilapia were highly contaminated by *E. coli*

(48%), followed by shrimp (45%). Nile perch were highly contaminated by *Salmonella* spp., followed by Nile tilapia. Only shrimp and Nile perch were contaminated by *Shigella* spp., while *Staphylococcus* spp. were isolated in Nile tilapia and Nile perch only.

All samples were screened for *Campylobacter*, but no *Campylobacter* was recovered by culture.

Samples from the open-air market were contaminated by six bacterial species, whereas four species were detected in supermarket samples (Figure 1). Fish from the open-air market were contaminated by *E. coli* (40%), *Klebsiella* spp. (26%), and *Salmonella* spp. (24%) while supermarket samples were contaminated by *E. coli* (37%), *Klebsiella* spp. (33%), and *Staphylococcus* spp. (23%). *Salmonella* spp. was not detected in the supermarket samples, while *Staphylococcus* spp. was not detected in the open-air market samples.

3.3. Antibiogram Profile of Isolated Bacteria. In general, the isolates showed a low level of resistance to most of the antimicrobial agents. Of the 48 *E. coli* isolates, 14 (29%) were resistant to penicillin, 12 (25%) were resistant to erythromycin, 8 (17%) were resistant to gentamicin, and 7 (15%) and 5 (10%) were resistant to azithromycin and ciprofloxacin, respectively. For *Salmonella* spp. isolates, 8 (31%), 7 (27%), 6 (23%), and 3 (12%) showed resistance to gentamicin, tetracycline, penicillin, and erythromycin, respectively.

“For *Staphylococcus* spp., among the 16 isolates, 31% and 5% were resistant to tetracycline and gentamicin, respectively. However, resistance to ciprofloxacin, clindamycin, and penicillin was observed in 19%, 13%, and 13% of isolates, respectively.

Klebsiella spp. exhibited resistance to six antimicrobial agents comprising gentamicin, tetracycline, penicillin, erythromycin, azithromycin, and ciprofloxacin, whereas more than 20% were resistant to gentamicin (Table 3).

As observed in Figure 2, bacterial isolates from marine fish were more resistant to antibiotics than those isolated from freshwater fish. *Shigella* spp. isolated from marine fish were resistant to tetracycline 28 (17%), gentamicin, and ciprofloxacin, while no isolates from freshwater were resistant to antibiotics.

Salmonella spp. isolates recovered from marine fish showed resistance to six antimicrobial agents comprising gentamicin, tetracycline, penicillin, erythromycin, azithromycin, and ciprofloxacin. *Salmonella* spp. isolates recovered from freshwater fish exhibited resistance to only three antimicrobial agents comprising gentamicin, tetracycline, and erythromycin.

4. Discussion

Most outbreaks of food poisoning related to fish are derived from the consumption of raw or insufficiently heat-treated fish and cross-contamination during handling. According to Huss et al. [24] and Aberoumand [25], approximately 12% of foodborne outbreaks related to the consumption of fish are caused by bacteria.

TABLE 1: Mean total viable counts of retail fish in log₁₀ CFU per gram ± standard error (SE).

Fish species	N	*Mean ± SE	% of samples that fell into the quality category**		
			Good	Marginally acceptable	Unacceptable
Nile perch	24	4.70 ± 0.10 ^a	16.30	8.70	0
Shrimps	20	4.53 ± 0.29 ^a	15.22	6.52	0
Nile tilapia	24	3.82 ± 0.41 ^b	19.57	6.52	0
Red snapper	24	5.22 ± 0.13 ^a	7.61	19.57	0
Total	92		58.70	41.30	0

*Means followed by similar letters do not differ significantly ($P < 0.05$). **Microbiological quality categories of good ($< 5 \times 10^5$ CFU/g), marginally acceptable (5×10^5 to 1×10^7 CFU/g), and unacceptable ($> 1 \times 10^7$ CFU/g) [23].

TABLE 2: Occurrence of different bacteria on retail fish from Dar es Salaam.

Bacterial species	Prevalence (%)				
	Red snapper (n = 24)	Shrimps (n = 20)	Nile perch (n = 24)	Nile tilapia (n = 24)	Overall (n = 92)
<i>Citrobacter</i> spp.	1 (4)*	2 (10)	1 (4)	0	4 (4)
<i>E. coli</i>	10 (40)	9 (45)	6 (26)	12 (48)	36 (39)
<i>Klebsiella</i> spp.	9 (36)	7 (35)	4 (17)	6 (24)	26 (28)
<i>Pseudomonas</i> spp.	1 (4)	0	0	0	1 (1)
<i>Salmonella</i> spp.	3 (12)	2 (10)	6 (26)	4 (24)	15 (16)
<i>Shigella</i> spp.	1 (4)	0	2 (9)	0	3 (3)
<i>Staphylococcus</i> spp.	0	0	4 (17)	3 (12)	7 (8)

*Values in parentheses indicate the % incidence.




FIGURE 1: Prevalence of different bacteria on fish from open-air markets and supermarkets.

TABLE 3: Resistance profiles of bacterial spp. isolated from open-air markets and supermarkets.

*Antibiotics	Disc conc. (ug)	% of resistant isolates						
		<i>E. coli</i>	<i>Klebsiella</i> spp.	<i>Shigella</i> spp.	<i>Salmonella</i> spp.	<i>Pseudomonas</i> spp.	<i>Citrobacter</i> spp.	<i>Staphylococcus</i> spp.
TET	30	4	16	50	27	0	22	31
GEN	10	17	28	25	31	0	17	25
CIP	5	10	10	25	8	0	11	19
CLI	2	0	0	0	0	0	0	13
PEN	10	29	18	0	23	0	11	13
AZM	15	15	16	0	0	0	22	0
ERY	15	25	12	0	12	0	17	0

*Antibiotics: TET = tetracycline; GEN = gentamicin; CIP = ciprofloxacin; CLI = clindamycin; PEN = penicillin; AZM = azithromycin; ERY = erythromycin.




FIGURE 2: Prevalence of bacterial spp. resistance to antibiotics isolated from open-air markets and supermarkets. *TET = tetracycline; GEN = gentamicin; CIP = ciprofloxacin; CLI = clindamycin; PEN = penicillin; AZM = azithromycin; ERY = erythromycin.

In this study, a total of 7 different species of bacteria were isolated. The most prevalent bacteria were *E. coli* (39%), followed by *Klebsiella* spp. (28%) and *Salmonella* spp. (16%). Other bacteria isolated in this study are *Staphylococcus* spp. (8%) and *Shigella* spp. (3%). *Citrobacter* (4%) and *Pseudomonas* spp. (1%) were also isolated in this study.

Salmonella spp. was detected in fish samples (16%) collected from different retail markets in this study. Higher results were reported by Kumar et al. [26], Jegadeeshkumar et al. [27], and Budiati et al. [28] at rates of 30%, 43.8%, and 90% in fish samples, respectively. In contrast, other studies from Trinidad and Turkey detected *Salmonella* spp. at 0% and 5% in fish samples, respectively [29, 30].

The presence of some bacterial species isolated from this study can cause food poisoning, diarrhea, typhoid fever, and shigellosis. Sichewo et al. [1] suggested that, when present in food, pathogens such as *S. aureus*, *Salmonella*, *Shigella*, and *Pseudomonas* are most likely to cause foodborne diseases. The presence of *Salmonella*, *Shigella*, and *E. coli* in fish indicates faecal and environmental contamination [1]. Coliforms such as *E. coli* are not normal bacteria in fish and are commonly present where there has been faecal contamination from warm-blooded animals [31]. *E. coli* is known to be a reliable indicator of faecal contamination in small numbers and large numbers, and it is an indicator of mishandling [32]. The presence of *E. coli* in this study could be attributed to poor mishandling of fish by traders. The presence of these bacteria isolated from the open-air market samples might also possibly be due to unhygienic handling during transportation and storage. The use

of contaminated water for cleaning and processing fish in the fish market is presumably the cause of secondary contamination. The lack of proper drainage facilities and heavy fly infestation in these fish markets also promotes tertiary contamination to a great extent.

The isolation rate of *Staphylococcus* spp. in the retail fish samples in the present study (8%) is relatively low compared with previous reports, which had rates ranging from 19.9% to 61.7% [33–36]. However, in this study, *Staphylococcus* spp. were isolated in samples from supermarkets only, which might be associated with inappropriate handling of fish.

The findings of this study show that the mean TVC for 92 samples was 4.57 log CFU/g, ranging from 3.82 to 4.70 and 5.22 log CFU/g in Nile tilapia, Nile perch, and red snapper, respectively. This is similar to the study of Wang et al. [37], who found the TVCs for salmon and tilapia to be 5.58 and 4.37 log CFU/g, respectively. According to the International Commission on Microbiological Specifications for Foods criteria, the microbiological quality of the fish examined in this study is acceptable and does not pose a potential risk to public health.

The results from this study also showed that the enteric bacteria isolated from fish were resistant to some antibiotics. It was observed that a small percentage of *Salmonella* spp. were resistant to gentamicin, tetracycline, penicillin, and erythromycin. Similar to our results (except erythromycin), Ponce et al. [38] found that the majority of isolates were resistant to ampicillin, tetracycline, chloramphenicol, gentamicin, sulfisoxazole, streptomycin, and kanamycin.

A total of 29, 25, 17, 15, and 13% of the *E. coli* isolates exhibited resistance to penicillin, erythromycin, gentamicin, azithromycin, and tetracycline, respectively. This is in contrast to the results obtained by Kibret and Abera [39], where erythromycin and tetracycline were highly resistant to *E. coli* isolates at 89.4 and 72.6%, respectively. The high prevalence of resistance to tetracycline, ampicillin, and cotrimoxazole in *E. coli* in the region has also been reported by Sifuna et al. [40], in which *E. coli* demonstrated resistance mostly to ampicillin and tetracycline. According to Sifuna et al. [40], the resistance pattern reported in their study can also be linked to the use of these drugs in veterinary practice and lead to resistance in humans.

In this study, bacteria isolated from marine fish were more resistant to antibiotics than those from freshwater fish. *Salmonella* spp. isolates recovered from marine samples were resistant to six antimicrobial agents, while isolates recovered from freshwater fish were resistant to three antimicrobial agents. *Shigella* isolates recovered from marine fish were resistant to tetracycline, gentamicin, and ciprofloxacin. However, *Shigella* isolates recovered from freshwater samples were not resistant to any antimicrobial agents. In contrast, a study on the antimicrobial resistance in isolates from marine species showed a high occurrence of antibiotic resistance, with 68% of isolates demonstrating resistance to at least one antibiotic, which is higher than the occurrence of antibiotic resistance observed in the current study [41]. Another study examining antimicrobial resistance in marine mammals reported *E. coli* was resistant to doxycycline (5%), amoxicillin (4%), and gentamicin (1%). Lockwood et al. [42] carried out a study on bacteria isolated from stranded harbor seals over a 12-year period. The author reported that bacterial isolates exhibit resistance most frequently to ampicillin (74% resistant) and cephalothin (64% resistant). Antibiotic-resistant microbes and genes within the aquatic environment have been documented in various marine species from cephalopods to marine mammals and elasmobranchs, such as sharks [43–47].

Rose et al. [41] and Ahmed et al. [48] concluded that the presence of antibiotic-resistant bacteria from animals, and marine animals, in particular, indicates not only the widespread presence of the microbes but often a significant percentage of the bacteria demonstrating resistance to multiple antibiotics.

Oates et al. [49] reported that when wastewater discharge and runoff from agricultural activity, which carries antibiotics and resistant bacteria into terrestrial waterways, finds its way to marine coastlines, it may cause disease in marine organisms, contributing to antibiotic resistance. This, along with other anthropogenic contributions, may contribute to the elevation of natural background levels of antibiotic resistance genes in aquatic environments, encouraging their transfer into pathogens or serving as a means for antibiotic resistance propagation [44]. When resistant bacteria are introduced to animals or their environment, the animals may become sick, or resistance traits may be transferred to other bacterial species, or they may become a reservoir that transfers the bacteria and resistance back to humans and the environment [45]. The antibiotics used by the majority of the

Tanzanian population are unregulated and used indiscriminately, and statistics are rarely collected, hence increasing the risk to the environment and human and animal health. Therefore, due to the widespread acquisition of resistance, it is important that susceptibility tests be routinely performed to guide antibiotic treatment and policy.

5. Conclusion

In conclusion, this study demonstrated the presence of highly pathogenic agents such as *Salmonella* and *Shigella* species and opportunistic pathogens, and some carried antimicrobial resistance.

Their presence is a potential health risk/hazard to human beings and may cause disease to susceptible individuals, especially immune-compromised consumers.

This study has therefore proven the need for the adoption of proper hygienic measures from farm to fork and hygienic education for fish handlers/traders and consumers.

In addition, bacterial isolates recovered from marine fish samples were more resistant to antibiotics than isolates recovered from freshwater fish samples. The environment might be a reservoir of resistance genes and dispersal vectors due to the influence of anthropogenic activities in marine environments. The resistance patterns observed in this study imply that some components of resistance are likely related to environmental origins and may spread without the selective pressure of antibiotic use.

There is a need for research on antibiotic susceptibility surveillance in aquatic environments where fresh fish and marine fish are obtained for human consumption.

Data Availability

Data will be available on request.

Conflicts of Interest

The author declares that there are no conflicts of interest.

Acknowledgments

This study was supported by The World Academy of Science (TWAS), Grant number: 19-085G/BIO/AF/AC_I-FR3240310155 dated 30 October 2019. The funding provided the equipment and reagents used for this study. The author wishes to gratefully thank Mr. Mahona and Mr. Lazack of the chemistry and biology laboratory at the Open University of Tanzania for their assistance in the field and laboratory work. The author is also grateful to Dr. Alexander Mzula, Mr. Erick Oswald, and Mr. Mikidadi Abdallah, who assisted in the laboratory work at the College of Veterinary Medicine and Biomedical Sciences at the Sokoine University of Agriculture. The author also thank fish traders who participated in this study by offering the fish samples.

References

- [1] P. R. Sichewo, R. K. Gono, J. V. Muzvondiwa, and N. Sizanobuhle, "Isolation and identification of pathogenic

- bacteria in edible fish: a case study of fletcher dam in," *International Journal of Science and Research*, vol. 3, no. 11, pp. 897–904, 2013.
- [2] B. F. Abisoye, S. K. S. Ojo, R. Adeyemi, and O. Olajuyigbe, "Bacteriological assessment of some commonly sold fishes in Lagos metropolis market, Nigeria," *Prime Journal of Microbiology Research*, vol. 2, no. 1, pp. 23–26, 2011.
 - [3] Food and Agricultural Organisation, *The State of World Fisheries and Aquaculture*, Food and Agricultural Organisation, Rome, Italy, 2016.
 - [4] C. Béné, R. Arthur, H. Norbury et al., "Contribution of fisheries and aquaculture to food security and poverty reduction: assessing the current evidence," *World Development*, vol. 79, pp. 177–196, 2016.
 - [5] C. Y. Chan, N. Tran, S. Pethiyagoda, C. C. Crissman, T. B. Sulser, and M. J. Phillips, "Prospects and challenges of fish for food security in Africa," *Global Food Security*, vol. 20, pp. 17–25, 2019.
 - [6] United Republic of Tanzania, *The United Republic of Tanzania Ministry of Livestock and Fisheries Development Basic Data for Livestock and Fisheries Sectors*, United Republic of Tanzania, Dar es salaam, Tanzania, 2013.
 - [7] M. M. Cahill, "Bacterial flora of fishes: a review," *Microbial Ecology*, vol. 19, no. 1, pp. 21–41, 1990.
 - [8] I. J. Clucas and A. R. Ward, *Post-Harvest Fisheries Development: A Guide to Handling, Preservation, Processing and Quality*, Natural Resources Institute (NRI), Chatham, UK, 1996.
 - [9] B. Austin, "Bacterial pathogens of marine fish," *Oceans and Health: Pathogens in the Marine Environment*, pp. 391–413, Springer US, New York, NY, USA, 2005.
 - [10] O. L. M. Haenen, J. J. Evans, and F. Berthe, "Bacterial infections from aquatic species: potential for and prevention of contact zoonoses," *Revue Scientifique et Technique de L'OIE*, vol. 32, no. 2, pp. 497–507, 2013.
 - [11] M. Mhango, S. Mpuchane, and B. Mpuchane, "Incidence of indicator organisms, opportunistic and pathogenic bacteria in fish," *African Journal of Food, Agriculture, Nutrition and Development*, vol. 10, no. 10, pp. 4202–42018, 2010.
 - [12] E. K. Lipp and J. B. Rose, "The role of seafood in foodborne diseases in the United States of America," *Revue Scientifique et Technique de L'OIE*, vol. 16, no. 2, pp. 620–640, 1997.
 - [13] L. Novotny, L. Dvorska, A. Lorencova, V. Beran, and I. Pavlik, "Fish: a potential source of bacterial pathogens for human beings," *Veterinari Medicina*, vol. 49, no. 9, pp. 343–358, 2004.
 - [14] J. Jumbe, P. Kibas, D. Kakongoro, and R. Tumwebaze, *Current State of Handling, Processing and Quality of Omena (Rastrionebola argentea) in Mfangano and Rusinga Islands, Kenya*, pp. 57–68, Kenyatta University, Nairobi, Kenya, 2010.
 - [15] H. Y. Done and R. U. Halden, "Reconnaissance of 47 antibiotics and associated microbial risks in seafood sold in the United States," *Journal of Hazardous Materials*, vol. 282, pp. 10–17, 2015.
 - [16] R. Laxminarayan, A. Duse, C. Wattal et al., "Antibiotic resistance-the need for global solutions," *The Lancet Infectious Diseases*, vol. 13, no. 12, pp. 1057–1098, 2013.
 - [17] R. A. Stein, "Antibiotic resistance: a global, interdisciplinary concern," *The American Biology Teacher*, vol. 73, no. 6, pp. 314–321, 2011.
 - [18] R. Z. Sangeda, J. Kibona, C. Munishi et al., "Assessment of implementation of antimicrobial resistance surveillance and antimicrobial stewardship programs in Tanzanian health facilities a year after launch of the national action plan," *Frontiers in Public Health*, vol. 8, p. 454, 2020.
 - [19] C. Kurekci, S. L. Bishop-Hurley, P. E. Vercoe, Z. Durmic, R. A. M. Al Jassim, and C. S. McSweeney, "Screening of Australian plants for antimicrobial activity against *Campylobacter jejuni*," *Phytotherapy Research*, vol. 26, no. 2, pp. 186–190, 2012.
 - [20] S. L. Abbott, W. K. W. Cheung, and J. M. Janda, "The genus *Aeromonas*: biochemical characteristics, atypical reactions, and phenotypic identification schemes," *Journal of Clinical Microbiology*, vol. 41, no. 6, pp. 2348–2357, 2003.
 - [21] A. E. N. El Deen, M. Dorgham, A. H. M. Hassan, and A. S. Hakim, "Studies on *Aeromonas hydrophila* in cultured *Oreochromis niloticus* at Kafr el Sheikh governorate, Egypt with reference to histopathological alterations in some," *World Journal of Fish and Marine Sciences*, vol. 3, no. 6, pp. 233–240, 2014.
 - [22] Clinical and Laboratory Standards Institute (CLSI), *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically*, Clinical and Laboratory Standards Institute (CLSI), Wayne, IL, USA, 2018.
 - [23] International Commission on Microbiological Specifications for Foods (ICMSF), *Micro-organisms in Food-2, Sampling for Microbiological Analysis: Principles and Specific Applications*, Blackwell Scientific Publications, Oxford, UK, 1987.
 - [24] H. Huss, L. V. Jørgensen, and B. F. Vogel, "Control options for *Listeria monocytogenes* in seafoods," *International Journal of Food Microbiology*, vol. 62, no. 3, pp. 267–274, 2000.
 - [25] A. Aberoumand, "Estimation of microbiological variations in minced lean fish products," *World Journal of Fish and Marine Sciences*, vol. 2, no. 3, pp. 204–207, 2010.
 - [26] H. S. Kumar, R. Sunil, M. N. Venugopal, I. Karunasagar, and I. Karunasagar, "Detection of salmonella spp. in tropical seafood by polymerase chain reaction," *International Journal of Food Microbiology*, vol. 88, no. 1, pp. 91–95, 2003.
 - [27] D. Jegadeeshkumar, V. Saritha, K. Moorthy, and B. T. Sureshkumar, "Prevalence, antibiotic resistance and RAPD analysis of food isolates of *Salmonella* species," *International Journal of Bio-Technology and Research*, vol. 1, pp. 50–55, 2010.
 - [28] T. Budiati, G. Rusul, W. N. Wan-Abdullah, Y. M. Arip, R. Ahmad, and K. L. Thong, "Prevalence, antibiotic resistance and plasmid profiling of *Salmonella* in catfish (*Clarias gariepinus*) and tilapia (*Tilapia mossambica*) obtained from wet markets and ponds in Malaysia," *Aquaculture*, vol. 372–375, pp. 127–132, 2013.
 - [29] A. A. Adesiyun, "Prevalence of *Listeria* spp., *Campylobacter* spp., *Salmonella* spp., *Yersinia* spp. and toxigenic *Escherichia coli* meat and seafoods in Trinidad," *Food Microbiology*, vol. 10, no. 5, pp. 395–403, 1993.
 - [30] N. Ertas Onmaz, S. Abay, F. Karadal, H. Hizlisoy, N. Telli, and S. Al, "Occurrence and antimicrobial resistance of *Staphylococcus aureus* and *Salmonella* spp. in retail fish samples in Turkey," *Marine Pollution Bulletin*, vol. 90, no. 1–2, pp. 242–246, 2015.
 - [31] K. K. Chao, C. C. Chao, and W. L. Chao, "Suitability of the traditional microbial indicators and their enumerating methods in the assessment of fecal pollution of subtropical freshwater environments," *Journal of Microbiology, Immunology, and Infection*, vol. 36, no. 4, pp. 288–293, 2003.
 - [32] B. C. Echezona and E. C. Uzodinma, "Isolation and identification of pathogenic bacteria associated with frozen mackerel fish (*Scomber scombrus*) in a humid tropical

- environment," *African Journal of Agricultural Research*, vol. 6, no. 8, pp. 1947–1951, 2011.
- [33] K. Kusunoki, H. Ushioda, M. Jin et al., "Results of bacterial contamination of commercial raw fish and shellfish in Tama, Tokyo (1986–1996)," *Japanese Journal of Food Microbiology*, vol. 15, no. 3/4, pp. 161–165, 1998.
- [34] K. Omoe, M. Ishikawa, Y. Shimoda, D.-L. Hu, S. Ueda, and K. Shinagawa, "Detection of seg, seh, and sei genes in *Staphylococcus aureus* isolates and determination of the enterotoxin productivities of *S. aureus* isolates harboring seg, seh, or sei genes," *Journal of Clinical Microbiology*, vol. 40, no. 3, pp. 857–862, 2002.
- [35] E. Saito, N. Yoshida, J. Kawano, A. Shimizu, and S. Igimi, "Isolation of *Staphylococcus aureus* from raw fish in relation to culture methods," *Journal of Veterinary Medical Science*, vol. 73, no. 3, pp. 287–292, 2011.
- [36] P. Bujamma and P. Padmavathi, "Prevalence of *Staphylococcus aureus* in fish samples of local domestic fish market," *International Journal of Current Microbiology and Applied Sciences*, vol. 4, no. 5, pp. 427–433, 2015.
- [37] F. Wang, L. Jiang, Q. Yang et al., "Prevalence and antimicrobial susceptibility of major foodborne pathogens in imported seafood," *Journal of Food Protection*, vol. 74, no. 9, pp. 1451–1461, 2011.
- [38] E. Ponce, A. A. Khan, C.-M. Cheng, C. Summage-West, and C. E. Cerniglia, "Prevalence and characterization of *Salmonella enterica* serovar weltevreden from imported seafood," *Food Microbiology*, vol. 25, no. 1, pp. 29–35, 2008.
- [39] M. Kibret and B. Abera, "Antimicrobial susceptibility patterns of *E. coli* from clinical sources in northeast Ethiopia," *African Health Sciences*, vol. 11, no. Suppl 1, pp. S40–S45, 2011.
- [40] A. W. Sifuna, E. N. M. Njagi, P. Okemo, A. Munyalo, G. O. Orinda, and S. Kariuki, "Microbiological quality and safety of *Restrineobola argentea* retailed in Kisumu town markets, Kenya," *East African Medical Journal*, vol. 85, no. 10, pp. 509–513, 2008.
- [41] J. M. Rose, R. J. Gast, A. Bogomolni et al., "Occurrence and patterns of antibiotic resistance in vertebrates off the northeastern United States coast," *FEMS Microbiology Ecology*, vol. 67, no. 3, pp. 421–431, 2009.
- [42] S. K. Lockwood, J. L. Chovan, and J. K. Gaydos, "Aerobic bacterial isolations from harbor seals (*Phoca vitulina*) stranded in Washington: 1992–2003," *Journal of Zoo and Wildlife Medicine*, vol. 37, no. 3, pp. 281–291, 2006.
- [43] S. P. Johnson, S. Nolan, and F. M. Gulland, "Antimicrobial susceptibility of bacteria isolated from pinnipeds stranded in central and northern California," *Journal of Zoo and Wildlife Medicine*, vol. 29, pp. 288–294, 1998.
- [44] T. W. Greig, J. A. Bemiss, B. R. Lyon, G. D. Bossart, and P. A. Fair, "Prevalence and diversity of antibiotic resistant *Escherichia coli* in bottlenose dolphins (*Tursiops truncatus*) from the Indian river lagoon, Florida, and Charleston Harbor area, south Carolina," *Aquatic Mammals*, vol. 33, no. 2, pp. 185–194, 2007.
- [45] A. M. Schaefer, J. D. Goldstein, J. S. Reif, P. A. Fair, and G. D. Bossart, "Antibiotic-resistant organisms cultured from atlantic bottlenose dolphins (*Tursiops truncatus*) inhabiting estuarine waters of charleston, SC and Indian river lagoon, FL," *EcoHealth*, vol. 6, no. 1, pp. 33–41, 2009.
- [46] J. K. Blackburn, M. A. Mitchell, M.-C. H. Blackburn, A. Curtis, and B. A. Thompson, "Evidence of antibiotic resistance in free-swimming, top-level marine predatory fishes," *Journal of Zoo and Wildlife Medicine*, vol. 41, no. 1, pp. 7–16, 2010.
- [47] J. R. Stewart, R. J. Gast, R. S. Fujioka et al., "The coastal environment and human health: microbial indicators, pathogens, sentinels and reservoirs," *Environmental Health: A Global Access Science Source*, vol. 7, no. SUPPL. 2, pp. 1–14, 2008.
- [48] A. M. Ahmed, Y. Motoi, M. Sato et al., "Zoo animals as reservoirs of gram-negative bacteria harboring integrons and antimicrobial resistance genes," *Applied and Environmental Microbiology*, vol. 73, no. 20, pp. 6686–6690, 2007.
- [49] S. C. Oates, M. A. Miller, B. A. Byrne et al., "Epidemiology and potential land-sea transfer of enteric bacteria from terrestrial to marine species in the monterey bay region of California," *Journal of Wildlife Diseases*, vol. 48, no. 3, pp. 654–668, 2012.