A FIXED POINT THEOREM FOR A NONLINEAR TYPE CONTRACTION

S. A. HUSAIN and V. M. SEHGAL

Department of Mathematics
University of Wyoming
Laramie, Wyoming 82071

(Received January 31, 1978)

ABSTRACT. A well-known result of Boyd and Wong [1] on nonlinear contractions is extended. Several other known results are obtained as special cases.

INTRODUCTION.

In this paper, we extend a well-known result of Boyd and Wong [1] and obtain as consequences several other known results (see [2], [3], [4], [5]).

Throughout this paper, let (X, d) be a complete metric space, $\mathbb{R}^+$ the nonnegative reals and $\phi = \phi(t_1, t_2, t_3, t_4, t_5): (\mathbb{R}^+)^5 \rightarrow \mathbb{R}^+$ a function which is (a) continuous from right in each coordinate variable (b) nondecreasing in t_2, t_3, t_4, t_5 , and satisfies the inequality (c) $\phi(t, s, s, s, s) < \text{Max}\{t, s\}$ if $\text{Max}\{t, s\} \neq 0$ where $\{a, b\} \subseteq \{0, 1, 2\}$ with $a + b = 2$. Note that (c) implies that $\phi(t, t, t, t, t) < t$ for any $t > 0$.

2. MAIN RESULTS.

The following is the main result of this paper.

THEOREM 1. Let $f, g: X \rightarrow X$ be two commutative mappings such that

$$(i) \quad fX \subseteq gX,$$

$$(ii) \quad g \text{ is continuous,}$$

$$(iii) \quad d(fx, fy) \leq \phi(d(gx, gy), d(fx, gx), d(fy, gy), d(fx, gy), d(fy, gx)),$$

for each $x, y \in X$. Then, there exists a unique $u \in X$ with $fu = gu = u$.

We first prove the following lemma which simplifies the proof of the above theorem.

LEMMA. Under the conditions of Theorem 1, if there exists a $v \in X$ such that $fv = gv$, then there exists a unique $u \in X$ with $fu = gu = u$.

PROOF. We show that for any $w \in X$

$$f(w) = g(w) \text{ implies } f(v) = f(w) \quad (2.1)$$

Suppose $t = d(fv, fw) > 0$. Then it follows by (iii) that

$$t \leq \phi(t, 0, 0, t, t) \leq \phi(t, t, t, t, t) < t,$$

a contradiction. Thus $fv = fw$. Now, since $fw = gw$, therefore, $f(fw) = g(fw)$ and consequently by (2.1)

$$f(w) = f(fw) = g(fw).$$

Thus, if we set $u = f(w)$, then $fu = gu = u$. The uniqueness of u now follows from (2.1).

PROOF OF THEOREM 1. Let x_0 be an arbitrary point in X . Construct a sequence $\{y_n\}$ in X as follows. Let $y_0 = fx_0$. By (i) there exists a $x_1 \in X$ such that $y_0 = gx_1$. Set $y_1 = fx_1$. Thus, if $y_0, y_1, \dots, y_n$ are obtained with $y_n = fx_n$, there exists by (i) a $x_{n+1} \in X$ such that $y_n = gx_{n+1}$. Let $y_{n+1} = fx_{n+1}$. Thus, for each $n \in I$ (nonnegative Integers),

$$y_n = fx_n = gx_{n+1}. \quad (2.2)$$

We shall show that $\{y_n\}$ is a Cauchy sequence in X . For this, let for each $n \in I$, $d_n = d(y_n, y_{n+1})$. Then by (i) and (b),

$$d_{n+1} = d(fx_{n+1}, fx_{n+2}) \leq \phi(d_n, d_n, d_{n+1}, 0, d_n + d_{n+1}). \quad (2.3)$$

Now, if for some $n \in I$, $d_{n+1} > d_n$, then by (b) and (c)

$$d_{n+1} \leq \phi(d_n, d_{n+1}, d_{n+1}, 0, 2d_{n+1}) < d_{n+1},$$

a contradiction. Thus for each $n \in I$, $d_{n+1} \leq d_n$, that is $\{d_n\}$ is a nonincreasing sequence of nonnegative reals and consequently there exists a $d \in \mathbb{R}^+$ such that $\{d_n\} \rightarrow d$. Clearly $d = 0$, for otherwise by (2.3) and (c),

$$d \leq \phi(d, d, d, 0, 2d) < d,$$

a contradiction. Thus,

$$d_n \rightarrow 0. \quad (2.4)$$

Suppose, now that $\{y_n\}$ is not a Cauchy sequence. Then there exists a $E > 0$ such that for each $k \in I$, there exist integers $n(k)$, $m(k)$ with $k \leq n(k) < m(k)$ satisfying

$$E_k = d(y_{n(k)}, y_{m(k)}) > E.$$

Let $m(k)$ be the least integer greater than $n(k)$ such (2.4) holds. This implies that for each $k \in I$, $d(y_{n(k)}, y_{m(k)-1}) \leq E$. Consequently, for each $k \in I$,

$$E < E_k \leq d(y_{n(k)}, y_{m(k)-1}) + d(y_{m(k)-1}, y_{m(k)}) \leq E + d_k. \quad (2.5)$$

Hence, it follows by (2.4) that as $k \rightarrow \infty$, $E_k \rightarrow E$.

However, for each $k \in I$,

$$\begin{aligned} E_k &\leq d_{n(k)} + d(fx_{n(k)+1}, fx_{m(k)+1}) + d_{m(k)}, \\ &\leq 2d_k + \phi(E_k, d_k, d_k, E_k + d_k, E_k + d_k), \end{aligned}$$

Therefore, as $k \rightarrow \infty$,

$$E \leq \phi(E, 0, 0, E, E) < E,$$

contradicting the existence of $E > 0$. Thus, $\{y_n\}$ is a Cauchy sequence in X .

Consequently, there is a $v \in X$ such that $\{y_n\} \rightarrow v$, that is

$$fx_n = gx_{n+1} \rightarrow v. \quad (2.6)$$

We show that for this v ,

$$\alpha = d(fv, gv) = 0.$$

Suppose $\alpha > 0$. Now by (ii) and (2.6) we have,

$$fgx_n = gfx_n \rightarrow gv \text{ and } g^2x_n \rightarrow gv.$$

Also, it follows by (b) and (iii) that,

$$d(fgx_n, fv) \leq \phi(d(g^2x_n, gv), d(fgx_n, g^2x_n), \alpha, d(fgx_n, gv), \alpha + d(gv, g^2x_n)).$$

Therefore, as $n \rightarrow \infty$, the above inequality yields that

$$\alpha = d(gv, fv) \leq \phi(0, 0, \alpha, 0, \alpha) < \alpha,$$

a contradiction. Thus $fv = gv$ and hence by the above lemma, there is a unique $u \in X$ satisfying $fu = gu = u$.

In the special case when g is taken to be the identity map of x in Theorem 1, we have

COROLLARY 1. Let $f: X \rightarrow X$ satisfy either of the following conditions: for all $x, y \in X$,

$$(A). \quad d(fx, fy) \leq \phi(d(x, y), d(x, fx), d(y, fy), d(x, fy), d(y, fx)).$$

$$(B). \quad d(fx, fy) \leq \alpha(d(x, fx) + d(y, fy)) + \beta(d(x, fy) + d(y, fx)) + \Psi(d(x, y))$$

where $\alpha \geq 0$, $\beta \geq 0$ and $\Psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a right continuous function satisfying $\Psi(t) < (1-2\alpha-2\beta)t$ if $t > 0$. Then f has a unique fixed point in X .

PROOF. The conclusion is an obvious consequence of Theorem 1 if (A) holds.

In case of condition (B), let $\phi: (\mathbb{R}^+)^5 \rightarrow \mathbb{R}^+$ be defined by

$$\phi(t_1, t_2, t_3, t_4, t_5) = \Psi(t_1) + \alpha(t_2 + t_3) + \beta(t_4 + t_5).$$

then ϕ satisfies conditions (a), (b) and (c). Thus the conclusion again follows by Theorem 1.

It may be remarked that if $\alpha = \beta = 0$ in (B) then Corollary 1 yields a well-known result of Boyd and Wong [1]. If $\Psi(t) = \alpha t$, then Corollary 1 yields certain results of Hardy and Rogers [2], Kannan [3], Reich [4], Sehgal [5]. All these results are special cases of Theorem 1.

REFERENCES

1. Boyd, D. W. and S. W. Wong. On Nonlinear Contractions, Proc. Amer. Math Soc. 20 (1969) 458-464.
2. Hardy, G. E. and T. D. Rogers. A Generalization of a Fixed Point Theorem of Reich, Canad. Math. Bull. 16 (2) (1973) 201-206.
3. Kannan, R. Some Remarks on Fixed Points, Bull. Cal. Math. Soc. 60 (1968) 71-76.
4. Reich, S. Kannan's Fixed Point Theorems, Bull. U.M.I. 4 (1971) 1-11.
5. Sehgal, V. M. Some Fixed and Common Fixed Point Theorems in Metric Spaces, Canad. Math. Bull. 16 (1974) 257-259.

KEY WORDS AND PHRASES. *Nonlinear type contraction, Fixed point theorems.*

AMS(MOS) SUBJECT CLASSIFICATIONS (1970). 47H10.