A LOWER BOUND ON THE NUMBER OF FINITE SIMPLE GROUPS

MICHAEL E. MAYS

Department of Mathematics
West Virginia University
Morgantown, West Virginia 26506
U.S.A.

(Received March 5, 1980)

ABSTRACT. Let $S(n) = |\{m < n: \text{there is a (non-cyclic) simple group of order } m\}|$.

Investigation of known families of simple groups provides the lower bound

$S(n) \gg n^{1/4}/\log n$.

KEY WORDS AND PHRASES. Simple group, asymptotic formula.

1980 MATHEMATICAL SUBJECT CLASSIFICATION. Primary 20 D 06.

The non-specialist reader should refer first to Hurley and Rudvalis (4).

Write $S(n) = |\{m < n: \text{there is a simple group of order } m\}|$ and $S'(n) = |\{G: G \text{ is a simple group and } |G| < n\}|$. Dornhoff (1), Dornhoff and Spitznagel (2), and Erdős (3) got successively better upper bounds for $S(n)$ by refining an argument which uses the Sylow theorems to generate a necessary criterion for a simple group of order m to exist. From the observation that $S(n) \leq |\{m < n: \text{for any prime } p|m \text{ there is a } d|m \text{ such that } d > 1 \text{ and } d \equiv 1 \pmod{p}\}|$ Dornhoff found that

$S(n) = o(n)$ and Erdős derived a complicated bound better than that of Dornhoff but not as good as $o(n^{1-\varepsilon})$. It should be noted that in general $S(n) < S'(n)$ because it occasionally (in fact infinitely often) happens that non-isomorphic simple groups of the same order exist.

We offer the following lower bound for $S(n)$, hence for $S'(n)$

THEOREM. $S(n) \gg n^{1/4}/\log n$.

PROOF. We estimate the number of integers $m < n$ which can be the order of a simple group in one of the known families and note that in all but finitely many cases the orders of the groups in that family are distinct.

From a list of known families of simple groups (4, p. 708) we see that one family dominates in the sense that for $F_1(n) = |\{m < n: m \text{ is the order of a simple group in family 1}\}|$, $F_1(n) = O(F_i(n))$ for any i . $F_1(n)$ is the number of simple projective special linear groups of order less than n .

Thus to estimate $S(n)$ from below, we count tripletons (k, p, a) such that

- 1) k is an integer greater than 1,
- 2) a is an integer ≥ 1 , and if $p = 2$ or $p = 3$ and $k = 2$ then $a > 1$, and
- 3) p is a prime, and writing $q = p^a$ we have

$$f(k, p, a) = q^{k(k-1)/2} \prod_{i=2}^k (q^i - 1)/(q^i - p^i) = |\text{PSL}_k(q)| < n.$$

Artin (5) showed that in exactly two cases distinct tripletons give rise to isomorphic groups, and in one case there are non-isomorphic groups of the same order in that family. Since $f(k, p, a) < q^{k(k-1)/2} q^{(k(k+1)/2)-1} < q^{k^2}$, $S(n) \gg |\{m < n: \text{there exists } (k, p, a) \text{ satisfying 1), 2), and 3) such that}$

$m = p^{ak^2}\}$. Such tripletons may be counted by a triple sum, and we have

$$S(n) \gg \sum_{a=1}^{\infty} \sum_{k=2}^{\infty} \sum_{p < n^{1/ak^2}} 1. \quad \text{Constraining } a \text{ and } k \text{ so that } n^{1/ak^2} \geq 2,$$

$$S(n) \gg \sum_{a=1}^{\log n/4 \log 2} (\log n/a \log 2)^{1/2} \sum_{k=2}^{\pi(n^{1/ak})^2},$$

and the Prime Number Theorem using $a = 1$ and $k = 2$ yields $S(n) \gg n^{1/4}/\log n$.

This theorem is of interest because it has been conjectured (3) that $S'(n) = o(n^{1-\varepsilon})$, or even $S'(n) = o(n^{1/3})$. We have that $1/4 - \varepsilon$ is a lower bound on the exponent of n , and if when all simple groups are classified no new family denser than the projective special linear groups appears, analyzing a perhaps more complicated triple sum carefully should yield the best exponent b in the estimate $S'(n) = o(n^b)$.

ACKNOWLEDGMENT: I wish to thank Professor Raymond Ayoub of the Pennsylvania State University for his advice and help.

REFERENCES

1. Dornhoff, L. Simple groups are scarce, Proc. Am. Math. Soc. 19(1968) 692-696.
2. Dornhoff, L. and E. L. Spitznagel, Jr. Density of finite simple group orders, Math Zeitschrift 106(1968) 175-177.
3. Erdős, P. Remarks on some problems in number theory, Math. Balcanica 4.32 (1974) 197-202.
4. Hurley, J. and A. Rudvalis. Finite simple groups, Am. Math. Monthly 84(1977) 693-714.
5. Artin, E. The orders of the linear groups, Comm. Pure and Appl. Math. 8(1952) 355-365.