GENERALIZED KÖTHE-TOEPLITZ DUALS

I.J. MADDOX

Department of Pure Mathematics
Queen's University of Belfast
Belfast BT7 1NN
Northern Ireland

(Received November 9, 1979)

ABSTRACT. The α and β -duals spaces of generalized ℓ_p spaces are characterized, where $0 < p \leq \infty$. The question of when the α and β dual spaces coincide is also considered.

KEY WORDS AND PHRASES. Generalized Köthe-Toeplitz dual spaces, Sequences of linear operators, Generalized ℓ_p spaces.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES. 40C05, 40J05.

1. INTRODUCTION.

X and Y denote complex Banach spaces with zero elements θ , and $||\cdot||$ denotes the norm in either X or Y . The continuous dual of X is written X^* . By $s(X)$ we mean the space of all X -valued sequences $x = (x_k)$, where $x_k \in X$ for $k \in N = \{1, 2, 3, \dots\}$.

If $0 < p < \infty$, we mean by $\ell_p(X)$ the space of all X -valued sequences $x = (x_k)$ such that $\sum \|x_k\|^p < \infty$. Sums are over $k \in \mathbb{N}$, unless otherwise indicated.

By $\ell_\infty(X)$ we denote the space of all $x = (x_k)$ such that $\sup \|x_k\| < \infty$.

In case $X = \mathbb{C}$, the space of complex numbers, we write ℓ_p instead of $\ell_p(\mathbb{C})$.

Let $A = (A_k)$ denote a sequence of linear, but not necessarily bounded, operators on X into Y . If E is any nonempty subset of $s(X)$ then the α -dual of E is defined to be

$$E^\alpha = \{A : \sum \|A_k x_k\| < \infty, \text{ for all } x \in E\}.$$

The β -dual of E is defined to be

$$E^\beta = \{A : \sum A_k x_k \text{ converges, for all } x \in E\}.$$

Since Y is complete, we have $E^\alpha \subset E^\beta$. The α and β duals of E may be regarded as generalized Köthe-Toeplitz duals, since in case $X = Y = \mathbb{C}$, when the A_k may be identified with complex numbers a_k , the duals reduce to the classical spaces first considered by Köthe and Toeplitz [1].

Using the notation $(1/p) + (1/q) = 1$, where $1 \leq p \leq \infty$, with the convention that $q = \infty$ when $p = 1$, and $q = 1$ when $p = \infty$, it is well-known that

$$\ell_p^\alpha = \ell_p^\beta = \ell_q. \quad (1.1)$$

We shall see that, in general, $\ell_p^\alpha(X) \subset \ell_p^\beta(X)$, where the inclusion may be strict. However, when $0 < p \leq 1$ the α and β duals coincide. Also, when $1 < p \leq \infty$, the α and β duals coincide provided that Y is finite dimensional.

2. CHARACTERIZATION OF THE DUALS.

THEOREM 1. Let $0 < p \leq 1$. Then $A \in \ell_p^\beta(X)$ if and only if there exists $m \in \mathbb{N}$ such that A_k is bounded, for all $k \geq m$, and

$$H = \sup_{k \geq m} \|A_k\| < \infty. \quad (2.1)$$

PROOF. Sufficiency. Let (2.1) hold and $\sum \|x_k\|^p < \infty$. By a familiar inequality, see for example Maddox [2], page 22,

$$\begin{aligned} \left(\sum_{k=m}^{\infty} \|A_k x_k\| \right)^p &\leq \sum_{k=m}^{\infty} \|A_k x_k\|^p \\ &\leq \sum_{k=m}^{\infty} \|A_k\|^p \|x_k\|^p \\ &\leq H^p \sum_{k=m}^{\infty} \|x_k\|^p. \end{aligned}$$

Hence $\sum A_k x_k$ is absolutely convergent, and so convergent.

Necessity. Let $A \in \ell_p^\beta(X)$ and suppose, if possible, that no such m exists.

Then there are natural numbers $k(1) < k(2) < \dots$ and $z_i \in X$, $\|z_i\| \leq 1$, such that for $i \in \mathbb{N}$,

$$\|A_{k(i)} z_i\| > i^{2/p}. \quad (2.2)$$

Define $x_k = z_i / i^{2/p}$ for $k = k(i)$ and $x_k = \theta$ otherwise. Then $x \in \ell_p(X)$ since $\sum \|x_k\|^p \leq \pi^2/6$, but $\|A_k x_k\| > 1$ for infinitely many k , contrary to the fact that $\sum A_k x_k$ converges.

Now suppose, if possible, that $\sup_{k \geq m} \|A_k\| = \infty$. Then there are natural numbers $k(1) < k(2) < \dots$ with $k(1) \geq m$ such that for $i \in \mathbb{N}$,

$$\|A_{k(i)}\| > 2i^{2/p}. \quad (2.3)$$

Choose $z_i \in X$ with $\|z_i\| \leq 1$ such that $2\|A_{k(i)} z_i\| \geq \|A_{k(i)}\|$, so by (2.3)

we see that (2.2) holds with the new $k(i)$ and z_i . We may define $x \in \ell_p(X)$ as above and obtain a contradiction. Hence (2.1) must hold, and the proof is complete.

If we examine the proof of Theorem 1 we see that in the sufficiency we had $\sum \|A_k x_k\| < \infty$, so that $A \in \ell_p^\alpha(X)$. Also, in the necessity, the constructions involved $x \in \ell_p(X)$ such that $\sum \|A_k x_k\|$ was divergent. Hence we have:

THEOREM 2. If $0 < p \leq 1$ then

$$\ell_p^\alpha(X) = \ell_p^\beta(X).$$

Next we consider the case $1 < p < \infty$.

THEOREM 3. Let $1 < p < \infty$. Then $A \in \ell_p^\alpha(X)$ if and only if there exists $m \in \mathbb{N}$ such that A_k is bounded for all $k \geq m$, and

$$M = \sum_{k=m}^{\infty} \|A_k\|^q < \infty. \quad (2.4)$$

PROOF. Sufficiency. Let (2.4) hold and $x \in \ell_p(X)$. By Hölder's inequality,

$$\sum_{k=m}^{\infty} \|A_k x_k\| \leq M^{1/q} (\sum_{k=m}^{\infty} \|x_k\|^p)^{1/p} < \infty.$$

Necessity. Since $\ell_p^\alpha(X) \subset \ell_1^\alpha(X)$ when $p > 1$, the existence of the m in the theorem follows from Theorems 1 and 2.

Now for $k \geq m$ we may choose $z_k \in X$ with $\|z_k\| \leq 1$ such that $2\|A_k z_k\| \geq \|A_k\|$.

For all $\lambda \in \ell_p$ we have $(\lambda_k z_k) \in \ell_p(X)$, so

$$\sum_{k=m}^{\infty} |\lambda_k| \|A_k z_k\| < \infty$$

for all $\lambda \in \ell_p$. By (1.1) it follows that

$$H = \sum_{k=m}^{\infty} \|A_k z_k\|^q < \infty,$$

whence $M \leq 2^q H$, so (2.4) holds, and the proof is complete.

THEOREM 4. Let $1 < p < \infty$. Then $A \in \ell_p^\beta(X)$ if and only if there exists $m \in \mathbb{N}$ such that A_k is bounded for all $k \geq m$, and

$$\sup \sum_{k=m}^{\infty} \|A_k^* f\|^q < \infty, \quad (2.5)$$

where the supremum is over all $f \in Y^*$ with $\|f\| \leq 1$.

PROOF. With the restriction that all the A_k are bounded, and with different notation, this result was proved by Thorp [3]. Only the existence of m in the necessity needs attention, and this follows from Theorems 1 and 2, and the fact that $\ell_p^\beta(X) \subset \ell_1^\beta(X)$.

Finally, we examine the case $p = \infty$. The proofs are left to the reader. We remark that with the restriction that all the A_k are bounded, the result concerning $\ell_\infty^\beta(X)$ was given by Maddox [4].

THEOREM 5. $A \in \ell_\infty^\alpha(X)$ if and only if there exists $m \in \mathbb{N}$ such that A_k is bounded for all $k \geq m$, and

$$\sum_{k=m}^{\infty} \|A_k\| < \infty. \quad (2.6)$$

THEOREM 6. $A \in \ell_\infty^\beta(X)$ if and only if there exists $m \in \mathbb{N}$ such that A_k is bounded for all $k \geq m$, and

$$\sup \left\| \sum_{k=m}^{m+n} A_k x_k \right\| < \infty, \quad (2.7)$$

$$\sup \left\| \sum_{k=m}^{m+n} A_k x_k \right\| \rightarrow 0 \quad (m \rightarrow \infty), \quad (2.8)$$

where the suprema are over all $n \geq 0$ and all $x_k \in X$ with $\|x_k\| \leq 1$.

3. COINCIDENCE OF DUALS.

It was shown in Theorem 2 that, when $0 < p \leq 1$, $\ell_p^\alpha(X) = \ell_p^\beta(X)$ for any Banach spaces X and Y .

We next shown that, when $1 < p < \infty$, the inclusion $\ell_p^\alpha(X) \subset \ell_p^\beta(X)$ may be strict.

THEOREM 7. If $1 < p < \infty$ then there are Banach spaces X and Y such that $\ell_p^\alpha(X) \subset \ell_p^\beta(X)$ with strict inclusion.

PROOF. Take $X = Y = \ell_p$ and write

$$e_k = (0, 0, \dots, 1, 0, 0, \dots)$$

where 1 is in the k -place and there are zeros elsewhere. Define bounded linear operators A_k on ℓ_p into itself by

$$A_k x = x_k e_k$$

for each $x = (x_k) \in \ell_p$. Then $\|A_k\| = 1$ for all $k \in \mathbb{N}$, so A is not in $\ell_p^\alpha(X)$ by Theorem 3.

Let us now show that (2.5) holds. Take any $f \in \ell_p^*$ with $\|f\| \leq 1$. Then for $x \in \ell_p$ we have

$$f(x) = \sum_i f_i x_i$$

for some (f_i) such that $\sum |f_i|^q \leq 1$. Hence, by definition of A_k^* ,

$$(A_k^* f)(x) = f(A_k x) = f_k x_k$$

and so $\|A_k^* f\| = |f_k|$. Hence

$$\sum \|A_k^* f\|^q = \sum |f_k|^q \leq 1,$$

so by Theorem 4 we have $A \in \ell_p^\beta(X)$.

Still with the case $1 < p < \infty$ we have:

THEOREM 8. If $1 < p < \infty$ and Y is finite dimensional then for any X we have

$$\ell_p^\alpha(X) = \ell_p^\beta(X).$$

PROOF. We have to show that $A \in \ell_p^\beta(X)$ implies $A \in \ell_p^\alpha(X)$. Now if $A \in \ell_p^\beta(X)$ then by Theorem 4 there exists $m \in \mathbb{N}$ such that A_k is bounded for all $k \geq m$. Suppose Y has finite dimension n and that $(b_1, b_2, \dots, b_n)$ is a Hamel base for Y . Then $y \in Y$ implies

$$y = \sum_{i=1}^n \lambda_i(y) b_i$$

where each $\lambda_i \in Y^*$. Take $z \in X$ and $k \geq M$. Then

$$A_k z = \sum_{i=1}^n \lambda_i(A_k z) b_i \quad (2.9)$$

and $\lambda_i \circ A_k \in X^*$. Since $\sum_{k=m}^{\infty} A_k x_k$ converges for all $x \in \ell_p(X)$ we have

$$\sum_{k=m}^{\infty} (\lambda_i \circ A_k) x_k$$

convergent for all $x \in \ell_p(X)$ and each i .

Choose $z_k \in X$, $\|z_k\| \leq 1$ such that $2|(\lambda_i \circ A_k)z_k| \geq \|\lambda_i \circ A_k\|$.

If $t \in \ell_p$ then $(t_k z_k) \in \ell_p(X)$ so that

$$\sum_{k=m}^{\infty} t_k (\lambda_i \circ A_k) z_k$$

converges for all $t \in \ell_p$, whence for each i ,

$$\sum_{k=m}^{\infty} \|\lambda_i \circ A_k\|^q < \infty. \quad (2.10)$$

By (2.9) and Hölder's inequality,

$$\|A_k\|^q \leq \sum_{i=1}^n \|\lambda_i \circ A_k\|^q \cdot \left(\sum_{i=1}^n \|b_i\|^p \right)^{q/p}. \quad (2.11)$$

Denoting the final term in (2.11) by H ,

$$\sum_{k=m}^{\infty} \|A_k\|^q \leq H \sum_{i=1}^n \sum_{k=m}^{\infty} \|\lambda_i \circ A_k\|^q. \quad (2.12)$$

It follows from (2.10) and (2.12) that (2.4) holds, so by Theorem 3 we have

$$A \in \ell_p^\alpha(X).$$

For certain values of p , and any X , the next result is the converse of Theorem 8.

THEOREM 9. If $2 < p < \infty$ and $\ell_p^\alpha(X) = \ell_p^\beta(X)$ then Y must be finite dimensional.

PROOF. Suppose, if possible, that Y is infinite dimensional. Since $q < 2$, if $c_k = k^{-2/q}$ then $\sum c_k < \infty$. By the Dvoretzky-Rogers theorem [5], there exists an unconditionally convergent series $\sum y_k$ in Y such that $\|y_k\|^2 = c_k$ for $k \in \mathbb{N}$. Hence

$$\sum \|y_k\|^q \text{ diverges.} \quad (2.13)$$

Take $f \in X^*$ with $\|f\| = 1$ and define rank one operators $A_k = y_k \otimes f$.

Then $\|A_k\| = \|y_k\|$, so by (2.13) and Theorem 3, A is not in $\ell_p^\alpha(X)$.

Now if $x \in \ell_p(X)$ then

$$\sum A_k x_k = \sum f(x_k) y_k.$$

But $(f(x_k)) \in \ell_\infty$ and $\sum y_k$ is unconditionally convergent, so that $\sum f(x_k) y_k$ converges, whence $A \in \ell_p^\beta(X)$, which gives a contradiction.

We remark that it would appear that the argument of Theorem 9 cannot be used in the case $p = 2$, since in a general Hilbert space Y the unconditional convergence of $\sum y_k$ implies that $\sum \|y_k\|^2$.

However, we can deal with the case $p = 2$ of Theorem 9 when Y is a Hilbert space:

THEOREM 10. Let Y be a Hilbert space and suppose $\ell_2^\alpha(X) = \ell_2^\beta(X)$.

Then Y must be finite dimensional.

PROOF. Suppose, if possible, that Y is infinite dimensional. Choose an orthonormal sequence (e_k) in Y and denote the inner product in Y by (y_1, y_2) . Take $g \in X^*$, $\|g\| = 1$ and define rank one operators $A_k = e_k \otimes g$, so that $\|A_k\| = 1$. Now let $f \in Y^*$ with $\|f\| \leq 1$. Then there exists $y \in Y$ such that

$$f(z) = (z, y)$$

for all $z \in Y$, with $\|y\| = \|f\| \leq 1$. Then for $x \in X$,

$$(A_k^* f)(x) = (g(x) e_k, y) = g(x) (e_k, y).$$

Hence $\|A_k^* f\| \leq |(e_k, y)|$, so by Bessel's inequality,

$$\sum \|A_k^* f\|^2 \leq \|y\|^2 \leq 1.$$

Thus (2.5) holds with $q = 2$, and so $A \in \ell_2^\beta(X)$. But $A \notin \ell_2^\alpha(X)$ since $\|A_k\| = 1$ for all k . This contradiction implies our result.

The case $p = \infty$ is due essentially to Thorp [3], who shows that

$\ell_{\infty}^{\alpha}(X) = \ell_{\infty}^{\beta}(X)$ if and only if Y is finite dimensional.

REFERENCES

1. Köthe, G. and Toeplitz, O. Lineare Räume mit unendlich vielen Koordinaten und Ringe unendlicher Matrizen, J. reine angew. Math. 171 (1934), 193-226.
2. Maddox, I.J. Elements of Functional Analysis, Cambridge University Press, 1970.
3. Thorp, B.L.D. Sequential-evaluation convergence, J. London Math. Soc. 44 (1969), 201-209.
4. Maddox, I.J. Matrix maps of bounded sequences in a Banach space, Proc. American Math. Soc. 63 (1977), 82-86.
5. Dvoretzky, A. and Rogers, C.A. Absolute and unconditional convergence in normed linear spaces, Proc. Nat. Acad. Sci. (U.S.A.). 36 (1950) 192-197.