

## **a\*-FAMILIES OF ANALYTIC FUNCTIONS**

**G.P. KAPOOR and A.K. MISHRA**

Department of Mathematics  
Indian Institute of Technology, Kanpur  
Kanpur - 208016, U.P., India

(Received February 15, 1983)

**ABSTRACT.** Using convolutions, a new family of analytic functions is introduced. This family, called  $a^*$ -family, serves in certain situations to unify the study of many previously well known classes of analytic functions like multivalent convex, starlike, close-to-convex or prestarlike functions, functions starlike with respect to symmetric points and other such classes related to the class of univalent or multivalent functions. A necessary and sufficient condition on the Taylor series coefficients so that an analytic function with negative coefficients is in an  $a^*$ -family is obtained and sharp coefficients bound for functions in such a family is deduced. The extreme points of an  $a^*$ -family of functions with negative coefficients are completely determined. Finally, it is shown that Zmorvic conjecture is true if the concerned families consist of functions with negative coefficients.

*KEY WORDS AND PHRASES.* Univalent Functions, Multivalent Functions, Convolution,  $p$ -valent starlike functions,  $p$ -valent close-to-convex functions,  $p$ -valent prestarlike functions, starlike functions with respect to symmetric points, Coefficients bound, Extreme points etc.  
*1980 MATHEMATICS SUBJECT CLASSIFICATION (1980):* 30C45, 30C50, 30C55

### 1. INTRODUCTION.

Let  $A_p$ ,  $p = 1, 2, \dots$ , denote the family of functions  $f$ , analytic in  $E = \{z : |z| < 1\}$  and having Taylor series expansion

$$f(z) = z^p + \sum_{k=1}^{\infty} a_k z^{k+p}. \quad (1.1)$$

In the present paper we introduce the concept of an  $a^*$ -family of functions in  $A_p$ . It turns out that many familiar subfamilies of  $A_p$ , related to univalent and multivalent functions, are  $a^*$ -families. We determine a sufficient condition, on the coefficients, such that a function  $f$  in  $A_p$ , given by (1.1), is an  $a^*$ -family. Further, we show that such a condition is also necessary when  $f$  is in  $A[p]$ , the family of functions  $f$  in  $A_p$  having Taylor series expansion

$$f(z) = z^p - \sum_{k=1}^{\infty} |a_k| z^{k+p}. \quad (1.2)$$

Using these results we determine the extreme points of an  $a^*$ -family in  $A[p]$ . Finally, we give some applications of our results in Section 4.

### 2. DEFINITION AND EXAMPLES.

The Hadamard product or convolution  $f * g$  of functions  $f$  and  $g$ , analytic in  $E$  and

given by  $f(z) = \sum_{k=0}^{\infty} a_k z^k$  and  $g(z) = \sum_{k=0}^{\infty} b_k z^k$ , is defined as the analytic function  $(f * g)(z) = \sum_{k=0}^{\infty} a_k b_k z^k$ .

Definition. A family of functions  $f$  in  $A_p$ ,  $p = 1, 2, \dots$ , is said to be an  $a^*$ -family if there exist functions  $s_o$  and  $g_o$  analytic in  $E$  defined by

$$s_o(z) = \sum_{k=0}^{\infty} c_k z^{k+p}, \quad c_o > 0, \quad c_k \geq 0, \quad k = 1, 2, \dots \tag{2.1}$$

and

$$g_o(z) = \sum_{k=0}^{\infty} d_k z^{k+p}, \quad d_o > 0, \quad d_k \geq 0, \quad k = 1, 2, \dots \tag{2.2}$$

satisfying

$$\frac{c_k}{c_o} - \frac{d_k}{d_o} > 0, \quad k = 1, 2, \dots \tag{2.3}$$

and a number  $\rho$ ,  $0 \leq \rho \leq c_o / d_o$ , such that for every  $f$  in  $F$

$$(g_o * f)(z) \neq 0, \quad 0 < |z| < 1 \tag{2.4}$$

and

$$\operatorname{Re} \left( \frac{(s_o * f)(z)}{(g_o * f)(z)} \right) > \rho \tag{2.5}$$

for  $z$  in  $E$ .

We write- $F$  is an  $a^*$ -family with the tuple  $(s_o, g_o, \rho)$ - when the tuple  $(s_o, g_o, \rho)$  is explicitly needed. Further, we denote an  $a^*$ -family in  $A[p]$  by  $[a^*]$ -family.

The following well known families in  $A_p$  are some examples of  $a^*$ -families.

Example 1 For  $0 \leq \alpha < 1$  and  $p = 1, 2, \dots$ , let  $S^*(p, \alpha)$  denote the  $a^*$ -family in  $A_p$  with the tuple

$$\left( \left\{ \frac{z^m}{1-z} \right\}_{m=1}^{p-1}, \left\{ \frac{z^m}{1-z} \right\}_{m=1}^p, p, \alpha \right) \tag{2.6}$$

and  $K(p, \alpha)$  denote the  $a^*$ -family in  $A_p$  with the tuple

$$\left( \left\{ \frac{z(1+z)}{(1-z)^2} \right\}_{m=1}^{p-1}, \left\{ \frac{z}{(1-z)^2} \right\}_{m=1}^p, p, \alpha \right) \tag{2.7}$$

It is easily seen that  $S^*(p, \alpha)$  is the family of  $p$ -valent starlike functions of order  $\alpha$  i.e. family of functions  $f$  in  $A_p$  satisfying  $\operatorname{Re}(zf'(z)/f(z)) > p\alpha$ . Also,  $K(p, \alpha)$  is the family of  $p$ -valent convex functions of order  $\alpha$  i.e. family of functions  $f$  in  $A_p$  such that  $zf'/p$  is in  $S^*(p, \alpha)$  [4].

Example 2. Let  $K(n+p-1)$  denote the class of functions  $f$  in  $A_p$  satisfying the condition

$$\operatorname{Re} \left( \frac{(z^n f)^{(n+p)}}{(z^{n-1} f)^{(n+p-1)}} \right) > \frac{n+p}{2}$$

where  $p = 1, 2, \dots$ ,  $n = -p+1, -p+2, \dots, p$ ; and  $z$  is in  $E$ . It is known [1] that  $f$  in  $A_p$  is in  $K(n+p-1)$ , if and only if,  $\operatorname{Re} G(z) > \frac{1}{2}$  where

$$G(z) = \frac{\frac{z^p}{(1-z)^{n+p+1}} * f(z)}{\frac{z^p}{(1-z)^{n+p}} * f(z)}$$

A brief calculation shows that, for  $h(z) = (z^p/(1-z)^{n+p}) * f(z)$  and  $f$  in  $K(n+p-1)$

$$\operatorname{Re}(zh'(z)/h(z)) = (p+n) \operatorname{Re} G(z) - n > (p-n)/2 > 0$$

Thus,  $h(z) \neq 0$  in  $0 < |z| < 1$  and it follows that  $K(n+p-1)$  is an a\*-family with the tuple

$$\left( \frac{z^p}{(1-z)^{n+p+1}}, \frac{z}{(1-z)^{n+p}}, \frac{1}{2} \right). \tag{2.8}$$

Example 3. Let  $J(p, \alpha)$  denote the a\*-family in  $A_p$  with tuple

$$\left( \left\{ \frac{z}{(1-z)^2} - \sum_{m=1}^{p-1} m z^m \right\}, z^p, p\alpha \right) \tag{2.9}$$

where  $0 \leq \alpha < 1$ . It is easily seen that  $J(p, \alpha)$  is a known subfamily of the family of  $p$ -valent close-to-convex functions [5]. The family  $J(1, \alpha)$  is the family of univalent functions  $f$  in  $A_1$  satisfying  $\operatorname{Re}(f'(z)) > \alpha$  for  $z$  in  $E$ .

Example 4. The family  $I(p, \alpha)$ ,  $0 \leq \alpha < 1$ ,  $p = 1, 2, \dots$ , of functions  $f$  in  $A_p$  satisfying  $\operatorname{Re} \{f(z)/z^p\} > p\alpha$ , for  $z$  in  $E$ , is an a\*-family with the tuple

$$\left( \frac{z^p}{(1-z)}, z^p, p\alpha \right). \tag{2.10}$$

Example 5. The family  $N(\alpha)$  of univalent functions starlike of order  $\alpha$ ,  $0 \leq \alpha < 1$ , with respect to symmetric points i.e. the family of functions  $f$  in  $A_1$  satisfying

$$\operatorname{Re} (zf'(z)/(f(z) - f(-z))) > \alpha, \text{ is an a*-family with the tuple}$$

$$\left( \frac{z}{(1-z)^2}, \frac{z}{(1-z)^2}, \alpha \right). \tag{2.11}$$

Example 6. A function  $f$  in  $A_p$ ,  $p = 1, 2, \dots$ , is said to be  $p$ -valent prestarlike of order  $\alpha$ ,  $0 \leq \alpha < 1$ , if the function  $h(z) = (z^p/(1-z)^{2p(1-\alpha)}) * f(z)$  is  $p$ -valent starlike of order  $\alpha$ . A brief calculation shows that

$$\begin{aligned} F(z) &= \frac{p(1-2\alpha)}{2(1-\alpha)} + \frac{1}{2(1-\alpha)} \frac{zh'(z)}{h(z)} \\ &= \frac{\left[ \frac{p(1-2\alpha)}{2(1-\alpha)} \left\{ \frac{z}{(1-z)} - \sum_{m=2}^{p-1} z^m \right\} + \frac{1}{2(1-\alpha)} \left\{ \frac{z}{(1-z)^2} - \sum_{m=2}^{\infty} m z^m \right\} \right] * h(z)}{h(z)} \\ &= \left[ \sum_{m=0}^{\infty} \frac{2p(1-\alpha)+m}{2(1-\alpha)} z^{m+p} \right] * h(z)/h(z) \\ &= \frac{p z^p}{(1-z)^{2p(1-\alpha)+1}} * f(z) / \frac{z^p}{(1-z)^{2p(1-\alpha)}} * f(z). \end{aligned}$$

Further, a function  $f$  is  $p$ -valent prestarlike of order  $\alpha$ , if and only if,  $\operatorname{Re} F(z) > p/2$ . Now, it follows that the family  $PS^*(p, \alpha)$ , of  $p$ -valent prestarlike functions of order  $\alpha$ , is an a\*-family with the tuple

$$\left( \frac{z^p}{(1-z)^{2p(1-\alpha)+1}}, \frac{z^p}{(1-z)^{2p(1-\alpha)}}, \frac{1}{2} \right). \tag{2.12}$$

We note that  $PS^*(1, \alpha)$  is the class of prestarlike functions of order  $\alpha$  studied by Ruscheweyh [7].

3. NECESSARY AND SUFFICIENT CONDITIONS AND EXTREME POINTS.

We have the following sufficient condition on  $|a_n|$  for a function  $f$  in  $A_p$ , given by (1.1), to be in an  $a^*$ -family.

**Theorem 1.** Let  $f$ , in  $A_p$ , be given by (1.1). Let  $\{c_k\}_{k=0}^\infty$  and  $\{d_k\}_{k=0}^\infty$ , the sequences of real numbers with  $c_0 > 0, d_0 > 0, (c_k/c_0) - (d_k/d_0) \geq 0$ ; and  $\rho, 0 \leq \rho < (c_0/d_0)$  be such that

$$\sum_{k=1}^\infty (c_k - \rho d_k) |a_k| \leq (c_0 - \rho d_0). \tag{3.1}$$

Then,  $f$  is in the  $a^*$ -family with the tuple  $(s_0, g_0, \rho)$ , where  $s_0$  and  $g_0$  are defined by (2.1) and (2.2) respectively.

**Proof.** Let  $h(z) = (g_0 * f)(z)$ . Then, for  $z$  in  $E$ ,

$$\operatorname{Re} (h(z)/z^p) \geq d_0 - \sum_{k=1}^\infty d_k |a_k| |z|^k. \tag{3.2}$$

Now, since  $(c_k/c_0) - (d_k/d_0) \geq 0$  and (3.1) holds, we have

$$1 - \sum_{k=1}^\infty \frac{d_k}{d_0} |a_k| \geq 1 - \sum_{k=1}^\infty \frac{(c_k - \rho d_k)}{(c_0 - \rho d_0)} |a_k| \geq 0. \tag{3.3}$$

Thus, (3.2) and (3.3) give that  $h(z) \neq 0$  in  $0 < |z| < 1$ .

Next, for  $s_0$  and  $g_0$  defined by (2.1) and (2.2) and  $|z| = r$ ,

$$\begin{aligned} \operatorname{Re} \left( \frac{(s_0 * f)(z)}{(g_0 * f)(z)} - \rho \right) &= \operatorname{Re} \left( \frac{(c_0 - \rho d_0) + \sum_{k=1}^\infty (c_k - \rho d_k) a_k z^k}{d_0 + \sum_{k=1}^\infty d_k a_k z^k} \right) \\ &= \frac{(c_0 - \rho d_0) - \sum_{k=1}^\infty (c_k - \rho d_k) |a_k| r^k}{d_0 - \sum_{k=1}^\infty d_k |a_k| r^k}. \end{aligned}$$

Thus, by (3.1) and (3.3), we get

$$\operatorname{Re} \left( \frac{(s_0 * f)(z)}{(g_0 * f)(z)} \right) > \rho$$

Hence,  $f$  is in the  $a^*$ -family with the tuple  $(s_0, g_0, \rho)$  and this completes the proof of the theorem.

The following theorem gives a necessary and sufficient condition on  $|a_n|$  for a function  $f$  in  $A[p]$ , given by (1.2), to be in an  $[a^*]$ -family.

**Theorem 2.** Let the sequences  $\{c_k\}_{k=0}^\infty, \{d_k\}_{k=0}^\infty$  and the nonnegative number  $\rho$  be defined as in Theorem 1. Then a function  $f$  in  $A[p]$ , given by (1.2), is in the  $[a^*]$ -family with the tuple  $(s_0, g_0, \rho)$ , if and only if,

$$\sum_{k=1}^\infty (c_k - \rho d_k) |a_k| \leq (c_0 - \rho d_0) \tag{3.4}$$

where  $s_0$  and  $g_0$  are defined by (2.1) and (2.2) respectively.

**Proof.** Let  $f$  be in the  $[a^*]$ -family with the tuple  $(s_0, g_0, \rho)$ . Then, for  $|z| = r < 1$ ,

$$\operatorname{Re} \left( \frac{(s_0 * f)(z)}{(g_0 * f)(z)} - \rho \right) = \operatorname{Re} \left( \frac{(c_0 - \rho d_0) - \sum_{k=1}^\infty (c_k - \rho d_k) |a_k| z^k}{d_0 - \sum_{k=1}^\infty d_k |a_k| z^k} \right) > 0. \tag{3.5}$$

Now, let  $-1 < z < 1$ . By the condition (2.4),

$h(z) = d_0 z^p - \sum_{k=1}^{\infty} d_k |a_k| z^{k+p} \neq 0$  in  $0 < |z| < 1$ , so that

$d_0 - \sum_{k=1}^{\infty} d_k |a_k| z^{k+p} > 0$  for  $-1 < z < 1$ . It follows now by (3.5) that

$$\sum_{k=1}^{\infty} (c_k - \rho d_k) |a_k| z^k \leq (c_0 - \rho d_0). \tag{3.6}$$

Now, taking limit  $z \rightarrow 1$  along the real axis in (3.6), we get (3.4). Thus, the proof of the theorem is complete in view of Theorem 1.

Corollary 1. Let  $f$ , given by (1.2), be in an  $ia^*J$ -family with the tuple  $(s_0, g_0, \rho)$ , where  $s_0, g_0$  and  $\rho$  are defined as in the definition in Section 2.

Then, for  $k = 1, 2, \dots$

$$|a_k| \leq \frac{c_0 - \rho d_0}{c_k - \rho d_k}. \tag{3.7}$$

The inequality (3.7) is sharp, the extremal function being

$$f_k(z) = z^p - ((c_0 - \rho d_0)/(c_k - \rho d_k)) z^{k+p} \tag{3.8}$$

for each  $k = 1, 2, \dots$

Proof. The corollary is a direct consequence of the necessary and sufficient condition (3.4).

Corollary 2. Let  $F$  and  $G$  be two  $a^*$ -families with the tuples  $(s_0, g_0, 0)$  and  $(s_0, g_1, 0)$  respectively. Then,

$$F \cap A[p] = G \cap A[p]. \tag{3.9}$$

Remark 1. Choosing  $\{c_k\} \equiv \{k+p\}$ ,  $\{d_k\} \equiv 1$  and  $\rho = p\alpha$ ;  $k = 0, 1, 2, \dots$ ;  $p = 1, 2, \dots$ ; and  $0 \leq \alpha < 1$ , it follows from Theorem 1 that the condition

$$\sum_{k=1}^{\infty} (k+p - p\alpha) |a_k| \leq p(1-\alpha) \tag{3.10}$$

is sufficient for a function  $f$ , given by (1.1), to be in  $S^*(p, \alpha)$ . Further, by Theorem 2, it follows that the condition (3.10) is both necessary and sufficient for a function  $f$ , given by (1.2), to be in  $S^*[p, \alpha] \equiv S^*(p, \alpha) \cap A[p]$ .

If we choose  $\{c_k\} \equiv \{(k+p)^2\}$ ,  $\{d_k\} = \{k+p\}$  and  $\rho = p\alpha$ ,  $k = 0, 1, 2, \dots$ ,  $p = 1, 2, \dots$ ,  $0 \leq \alpha < 1$ , then Theorem 1 gives that the condition

$$\sum_{k=1}^{\infty} \frac{(k+p)}{p} (k+p-p\alpha) |a_k| \leq p(1-\alpha) \tag{3.11}$$

is sufficient for a function  $f$ , given by (1.1), to be in  $K(p, \alpha)$ . Further, by Theorem 2 we have that the condition (3.11) is both necessary and sufficient for a function  $f$ , given by (1.2), to be in  $K[p, \alpha] \equiv K(p, \alpha) \cap A[p]$ .

We note that the sufficient conditions (3.10) and (3.11) have been obtained by Ozaki [6], Goodman [2] and Schild [8] respectively in the particular cases  $\alpha = 0, p \geq 1$ ;  $\alpha = 0, p = 1$  and  $\alpha = \frac{1}{2}, p = 1$ . Silverman [9] has obtained similar necessary and sufficient conditions in the particular case  $p = 1$  and  $0 \leq \alpha < 1$ .

Remark 2. Choosing  $\{c_k\}_{k=0}^{\infty} \equiv \{k+p\}_{k=0}^{\infty}$ ,  $d_0 = 1$ ,  $d_k = 0, k=1, 2, \dots$  and  $\rho = p\alpha$ ,  $p = 1, 2, \dots$ ,  $0 \leq \alpha < 1$ , it follows from Theorem 1 that the condition

$$\sum_{k=1}^{\infty} (k+p) |a_k| \leq p(1-\alpha) \tag{3.12}$$

is sufficient for a function  $f$ , given by (1.1), to be in  $J(p, \alpha)$ . Further, by Theorem 2, it follows that the condition (3.12) is both necessary and sufficient for a function  $f$ , given by (1.2), to be in  $J[p, \alpha] \equiv J[p, \alpha] \cap A[p]$ .

**Remark 3.** With the choice  $\{c_k\}_{k=0}^\infty \equiv \{1\}$ ,  $d_0 = 1$  and  $d_k = 0$ ,  $k = 1, 2, \dots$  and  $\rho = p\alpha$ ,  $p = 1, 2, \dots$ ,  $0 \leq \alpha < 1$ , we have that, if

$$\sum_{k=1}^\infty |a_k| \leq p(1-\alpha) \tag{3.13}$$

then  $f$ , given by (1.1) is in  $I(p, \alpha)$ . Further, the condition (3.13) is both necessary and sufficient for a function  $f$ , given by (1.2), to be in  $I[p, \alpha] \equiv I[p, \alpha] \cap A[p]$ .

**Remark 4.** If we choose  $\{c_k\}_{k=0}^\infty \equiv \{k\}_{k=0}^\infty$ ,  $d_k = 1$  when  $k$  is odd and  $d_k = 0$  when  $k$  is even; and  $\rho = \alpha$ ,  $0 \leq \alpha < 1$ , then it follows from Theorem 1 that the condition

$$\sum_{k=1}^\infty (2k+1-\alpha) |a_{2k+1}| + \sum_{k=1}^\infty 2k |a_k| \leq (1-\alpha) \tag{3.14}$$

is sufficient for a function  $f$ , given by (1.1), to be in  $N(\alpha)$ . Further, it follows from Theorem 2 that the condition (3.14) is both necessary and sufficient for a function  $f$ , given by (1.2), to be in  $N[\alpha] \equiv N(\alpha) \cap A[p]$ .

**Remark 5.** Let  $z^p/(1-z)^{2p(1-\alpha)} = \sum_{k=0}^\infty C_o(\alpha, k+p) z^{k+p}$  and  $z^p/(1-z)^{2p(1-\alpha)+1} = \sum_{k=0}^\infty O_1(\alpha, k+p) z^{k+p}$ .

Then, by choosing  $\{c_k\}_{k=0}^\infty \equiv \{C_1(\alpha, k+p)\}_{k=0}^\infty$ ,  $\{d_k\}_{k=0}^\infty \equiv \{C_o(\alpha, k+p)\}_{k=0}^\infty$  and  $\rho = \frac{1}{2}$  it follows from Theorem 3.1 that the condition

$$\sum_{k=\frac{1}{2}}^\infty (k+p-p\alpha) C_o(\alpha, \rho+\alpha) |a_k| \leq p(1-\alpha) \tag{3.15}$$

is sufficient for  $f$ , given by (1.1), to be in  $PS^*(p, \alpha)$ .

Further, from Theorem 2 it follows that the condition (3.15) is both necessary and sufficient for a function  $f$ , given by (1.2) to be in  $PS^*[p, \alpha] \equiv PS^*(p, \alpha) \cap A[p]$ .

We note that (3.15) includes a recent result of Silverman and Silvia [10].

In view of Theorem 2, it follows that an  $[a^*]$ -family is a closed convex subset of the space of analytic functions in  $E$  with the compact open topology. Thus, the closed convex hull of an  $[a^*]$ -family  $F$  is equal to itself. In the next theorem we determine the extreme points of an  $[a^*]$ -family.

**Theorem 3.** Let the sequences  $\{c_k\}_{k=0}^\infty$ ,  $\{d_k\}_{k=0}^\infty$ , the nonnegative number  $\rho$  and the functions  $s_o$  and  $g_o$  be defined as in the definition in Section 2. Further, let  $f_o(z) = z^p$ ,

$f_k(z) = z^p - (c_o - \rho d_o)/(c_k - \rho d_k) z^{k+p}$ ,  $k = 1, 2, \dots$ . Then, the extreme points of the  $[a^*]$ -family with the tuple  $(s_o, g_o, \rho)$  are precisely the set of functions  $\{f_k\}_{k=0}^\infty$ .

**Proof.** We show that a function  $f$  is in the  $[a^*]$ -family with the tuple  $(s_o, g_o, \rho)$ ,

if and only if, it can be written in the form  $\sum_{k=0}^\infty t_k f_k(z)$ , where  $t_k \geq 0$  and

$\sum_{k=0}^\infty t_k = 1$ . The conclusion in the theorem about the extreme points is equivalent to this result.

First, let  $f(z) = \sum_{k=0}^\infty t_k f_k(z)$ , where  $t_k \geq 0$  and  $\sum_{k=0}^\infty t_k = 1$ . Then

$$f(z) = z^p - \sum_{k=1}^\infty t_k \frac{(c_o - \rho d_o)}{(c_k - \rho d_k)} z^{k+p}.$$

Now,

$$\sum_{k=1}^{\infty} \frac{(c_k - \rho d_k)}{(c_0 - \rho d_0)} t_k \frac{(c_0 - \rho d_0)}{(c_k - \rho d_k)} = \sum_{k=1}^{\infty} t_k = 1 - t_0 \leq 1.$$

Thus, by Theorem 2, f is in the [a\*]-family with the tuple (s<sub>0</sub>, g<sub>0</sub>, ρ).

Conversely, let f, defined by the Taylor series (1.2), be in the [a\*]-family with the tuple (s<sub>0</sub>, g<sub>0</sub>, ρ). Then, by Corollary 1, for k = 1, 2, ...,

$$|a_k| \leq \frac{c_0 - \rho d_0}{c_k - \rho d_k}.$$

We let, for k = 1, 2, ..., t<sub>k</sub> = ((c<sub>k</sub> - ρd<sub>k</sub>)/(c<sub>0</sub> - ρd<sub>0</sub>)) |a<sub>k</sub>| and t<sub>0</sub> = 1 -  $\sum_{k=1}^{\infty} t_k$ . Note that 0 ≤ t<sub>k</sub> ≤ 1 for k = 1, 2, 3, ... . Now with this choice of t<sub>k</sub>, we can write

f(z) =  $\sum_{k=0}^{\infty} t_k f_k(z)$  where  $\sum_{k=0}^{\infty} t_k = 1$ . This completes the proof of the theorem.

**Remark 6.** The extreme points of individual [a\*]-families can be obtained from Theorem 3 by substituting appropriate values of c<sub>k</sub>, d<sub>k</sub> and ρ as in Remarks 1-5.

4. APPLICATIONS.

In our next theorem, using Theorem 2, we determine the sharp values of β ≡ β(α) and γ ≡ γ(α) such that K[p, α] ⊂ S\* [p, β] and Re {f(z)/z<sup>p</sup>} > γ for f in K[p, α] and z in E. It is to be noted that, in general, there does not exist β = β(α) > α such that K(p, α) ⊂ S\* (p, α), p = 2, 3, ... . Further, the value of Re {f(z)/z<sup>p</sup>} can well be negative for |z| < 1 and f in K(p, α), p = 2, 3, ... [3].

**Theorem 4.** Let f, given by (1.2), be in K [p, α]. Then,

(i) f is in S\*[p, β] where

$$\beta \equiv \beta(\alpha) = \frac{p + 1}{2p + 1 - p\alpha} \tag{4.1}$$

and

(ii) Re{f(z)/z<sup>p</sup>} > γ, for z in E, where

$$\gamma \equiv \gamma(\alpha) = \frac{p^2(1-\alpha)+p+1}{(p+1)(p+1-p\alpha)}. \tag{4.2}$$

Both the results in (i) and (ii) are sharp.

**Proof.** Let f be in K [p, α]. Then, by the necessary condition (3.11), we have

$$\sum_{k=1}^{\infty} \frac{(k+p)(k+p-p\alpha)}{p^2(1-\alpha)} |a_k| \leq 1. \tag{4.3}$$

In view of the sufficient condition (3.10), we first determine the maximum value of β such that (4.3) implies

$$\sum_{k=1}^{\infty} \frac{(k+p-p\beta)}{p(1-\beta)} |a_k| \leq 1. \tag{4.4}$$

Again, it is sufficient to determine the maximum value of β such that for k = 1, 2, ...,

$$\frac{(k+p)(k+p-p\alpha)}{p(1-\alpha)} \geq \frac{(k+p-p\beta)}{1-\beta}$$

or, equivalently, β ≤ (k+p)/(2p+k-pα). Since the sequence {φ<sub>k</sub>} = {(k+p)/(2p+k-pα)} is increasing in k, we choose

$$\beta \equiv \beta(\alpha) = (p+1)/(2p+1-p\alpha). \tag{4.5}$$

Further, by using (3.10) and (3.11), it is easily seen that the function

$$h(z) \equiv z^p - \frac{p^2(1-\alpha)}{(p+1)(p+1-p\alpha)} z^{p+1} \quad (4.6)$$

in  $K[p, \alpha]$  is in  $S^*[p, \beta]$  with  $\beta$  defined by (4.1) but is not in  $S^*[p, \beta_1]$  for any  $\beta_1 > \beta$ . This shows that the value of  $\beta \equiv \beta(\alpha)$  is precise.

In view of (3.13), to prove (ii), we first find the maximum value of  $\gamma = \gamma(\alpha)$  such that (4.2) implies

$$\sum_{k=1}^{\infty} \frac{|a_k|}{p(1-\gamma)} \leq 1 \quad (4.7)$$

Adopting the proof for part (i), we get

$$\gamma \equiv \gamma(\alpha) = \frac{p^2(1-\alpha)+p+1}{(p+1)(p+1-p\alpha)}.$$

Further, the function  $h(z)$  given by (4.6) shows that  $\operatorname{Re}\{h(z)/z^p\} > p\gamma$  but  $\operatorname{Re}\{h(z)/z^p\} \leq p\gamma_1$  for  $\gamma_1 > \gamma$  and  $z = r < 1$  where  $\gamma$  is defined by (4.2). This completes the proof of the theorem.

We, next, consider Zmorovic conjecture for functions in  $A[1]$ . Zmorovic [11] conjectured that  $J(1,0) \subset S^*(1,0)$ . Subsequently, this conjecture was proved to be false, by Zmorovic himself [12] among others, by showing the existence of a function in  $J(1,0)$  which is not in  $S^*(1,0)$ . On the other hand,  $K(1,0) \not\subset J(1,0)$ , for the function  $h_1(z) = z/(1-z)$  is in  $K(1,0)$  and  $\operatorname{Re} h'(z) < 0$  in a region  $-\frac{1}{2} < 1,0 < \arg z < \pi/2$ . Thus, it follows that there is no inclusion relation between the three classes  $K(1,0)$ ,  $S^*(1,0)$  and  $J(1,0)$ . However, it follows easily from Corollary 2 that

$$S^*[1,0] = J[1,0] = N[0].$$

Further, for  $p = 1$ , (3.11)  $\implies$  (3.12)  $\implies$  (3.14)  $\implies$  (3.10)  $\implies$  (3.13). Thus, we have

$$K[1, \alpha] \not\subset J[1, \alpha] \not\subset N[\alpha] \not\subset S^*[1, \alpha] \not\subset I[1, \alpha].$$

#### REFERENCES

1. GOEL, R.M. and SOHI, N.S., A new criterion for  $p$ -valent functions, Proc. Amer. Math. Soc. 78 (1980) 353-357.
2. GOODMAN, A.W., Univalent functions and nonanalytic curves, Proc. Amer. Math. Soc. 8 (1957) 591-601.
3. HALLENBECK, D.J. and LIVINGSTON, A.E., Applications of extreme point theory to classes for multivalent functions, Trans. Amer. Math. Soc. 221 (1976) 339-359.
4. KAPOOR, G.P. and MISHRA, A.K., Convex hulls and extreme points for some classes of multivalent functions, J. Math. Anal. Appl. (To appear).
5. LIVINGSTON, A.E.,  $p$ -Valent close-to-convex functions, Trans. Amer. Math. Soc. 115 (1965) 161-179.
6. OZAKI, S., Some remarks on univalence and multivalence of functions, Science reports of the Tokyo Bunrika Daigaku Sect A, Vol. 2, No. 32 (1934) 41-55.
7. RUSCHEWEYH, St., Linear operators between classes of prestarlike functions, Comment. Math. Helv. 52 (1977) 497-509.
8. SCHILD, A., On a class of univalent star shaped mappings, Proc. Amer. Math. Soc. 9 (1958) 751-757.
9. SILVERMAN, H., Univalent functions with negative coefficients, Proc. Amer. Math. Soc. 51 (1975) 109-116.
10. SILVERMAN, H. and SILVIA, E.M., Prestarlike functions with negative coefficients, Int. J. Math. and Math. Sci. 2 (1979) 427-439.
11. ZMOROVIC, V.A., An open problem in the theory of univalent functions, Nauk Zaniski Kiev. Lerjeabvii Uni. 11 (1952) 83-94 (Russian).
12. ZMOROVIC, V.A., On certain special classes of analytic functions univalent in a circle, Uspehi Mat. Nauk 9 (1954).





# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

