

THE POULSEN SIMPLEX IS NOT A TENSOR PRODUCT

THOMAS E. ARMSTRONG

Department of Economics
University of Minnesota
Minneapolis, Minnesota 55455

(Received September 29, 1982)

ABSTRACT. It is shown that the Poulsen simplex is not a projective tensor product of non-trivial Choquet simplexes.

KEY WORDS AND PHRASES. *Poulsen simplex, projective tensor product, projective limits.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 46A55.

1. INTRODUCTION

The Poulsen simplex P is an example of a metrisable Choquet simplex whose extreme points $\xi(P)$ are dense in P . Such a simplex was constructed by Poulsen in [11]. In [6] Lazar and Lindenstrauss showed how to represent metrisable Choquet simplexes S as projective limits of an affine projective system $\{\{\Delta_n : n \in \mathbb{N}\}, \{\pi_n : n \in \mathbb{N}\}\}$ where each Δ_n is an $(n-1)$ -simplex with $\xi(\Delta_n) = \{e_1, \dots, e_n\} \subset \xi(\Delta_{n+1})$ and with $\pi_n : \Delta_{n+1} \rightarrow \Delta_n$ described by the requirement that $\pi_n(e_j) = e_j$ if $1 \leq j \leq n$ and $\pi_n(e_{n+1}) = a_{n1}e_1 + \dots + a_{nn}e_n$ where $a_{nj} > 0$ and $\sum a_{nj} = 1$. The triangular matrix $A = (a_{ij})$ is called a representing matrix for S . There are many representing matrices for S as there are many realizations of S as such a projective limit. It was established by Lindenstrauss, Olsen, and Sternfeld [7] that $S = \overline{\xi(S)}$ iff the sequence of rows of A form a dense subset of the positive face of the unit ball of ℓ^1 when each was regarded as a sequence. Lindenstrauss, Olsen and Sternfeld showed in [7] that, up to affine homeomorphism, P is the only metrisable Choquet simplex with $P = \overline{\xi(P)}$.

It is well known that the Poulsen simplex P is prime in that $A(K)$ is an anti-lattice (Asimow and Ellis, [2]). We show here that P is prime in the semigroup of

convex metrisable compact sets with multiplication being projective tensor product. The proof involves a fairly straight forward application of the properties of projective limits and of representing matrices for metrisable simplexes.

2. MAIN RESULTS

We refer the reader to E.B. Davies and G.F. Vincent-Smith [3] for the details concerning projective tensor products both finite and infinite. For any family $\{S_i : i \in I\}$ of Choquet simplexes there is defined, up to affine homeomorphism, a Choquet simplex $\bigotimes_{i \in I} S_i$, the projective tensor product, which has the property that there is a continuous multi-affine embedding \otimes of $\prod_{i \in I} S_i$ into $\bigotimes_{i \in I} S_i$ given by $(x_i : i \in I) \rightarrow \bigotimes_{i \in I} x_i$ so that if $m : \prod_{i \in I} S_i \rightarrow E$ is continuous multi-affine for some locally convex space E there exists a linear $n : \bigotimes_{i \in I} S_i \rightarrow E$ with $n \circ \otimes = m$. It is shown that $\xi(\bigotimes_{i \in I} S_i)$ is $\{\bigotimes_{i \in I} x_i : x_i \in \xi(S_i) : i \in I\}$ and that $\otimes : \prod_{i \in I} \xi(S_i) \rightarrow \xi(\bigotimes_{i \in I} S_i)$ is a homeomorphism. It is easily checked that if each S_i is a projective limit of a sequence $\{S_{in} : n \in \mathbb{N}\}$ of simplexes under projections $\{P_{in} : n \in \mathbb{N}\}$ then $\bigotimes_{i \in I} S_i$ is a projective limit of $\{\bigotimes_{i \in I} S_{in} : n \in \mathbb{N}\}$ under $\{P_n : n \in \mathbb{N}\}$ where $P_n : \bigotimes_{i \in I} S_{in+1} \rightarrow \bigotimes_{i \in I} S_{in}$ is the map induced by the multi-affine transformation $(x_i : i \in I) \rightarrow \bigotimes_{i \in I} P_{in}(x_i)$ from $\prod_{i \in I} S_{in+1}$ to $\prod_{i \in I} S_{in}$.

PROPOSITION. The Poulsen simplex is not a tensor product.

PROOF. Suppose that $P = X \otimes Y$ with X and Y each at least one dimensional metrisable Choquet simplexes. Let $A = (a_{ij})$ and $B = (b_{ij})$ be representing matrices for X and Y respectively. Let $p_n : \Delta_{n+1} \rightarrow \Delta_n$ and $q_n : \Delta_{n+1} \rightarrow \Delta_n$ for $n \in \mathbb{N}$ be the sequences of projections associated with A and B respectively so that X is the projective limit of $\{\Delta_n : n \in \mathbb{N}\}$ under $\{p_n : n \in \mathbb{N}\}$ and Y is the projective limit of $\{\Delta_n : n \in \mathbb{N}\}$ under $\{q_n : n \in \mathbb{N}\}$. Then P is the projective limit of $\{\Delta_n \otimes \Delta_n : n \in \mathbb{N}\}$ under the system $\{r_n : n \in \mathbb{N}\}$ of projections where $r_n(e_i \otimes e_j) = e_i \otimes e_j$ if $1 \leq i, j \leq n$, $r_n(e_{n+1} \otimes e_j) = \sum_{i=1}^n a_{ni} e_i \otimes e_j$, $r_n(e_i \otimes e_{n+1}) = \sum_{j=1}^n b_{nj} e_i \otimes e_j$ and $r_n(e_{n+1} \otimes e_{n+1}) = \sum_{i,j=1}^n a_{ni} b_{nj} e_i \otimes e_j$.

For any $n \in \mathbb{N}$, let $D_{n2} = \Delta_n \otimes \Delta_n$. For $1 \leq k \leq n$ define D_{n2+k} to be $\text{conv}(D_{n2+k-1}, e_k \otimes e_{n+1}) \subset \Delta_{n+1} \otimes \Delta_{n+1}$. Define D_{n2+n+k} to be $\text{conv}(D_{n2+n+k-1}, e_{n+1} \otimes e_k)$. Define, for $1 \leq k \leq n$, the affine surjection $R_{n2+k-1} : D_{n2+k-1} \rightarrow D_{n2+k-1}$ to be the identity on D_{n2+k-1} and to be $\sum_{j=1}^n b_{nj} e_k \otimes e_j$ on $e_k \otimes e_{n+1}$. Similarly define, for $1 \leq k \leq n$, the affine surjection $R_{n2+n+k-1} : D_{n2+n+k-1} \rightarrow D_{n2+n+k-1}$ by setting $R_{n2+n+k-1}$

equal to the identity on $D_{n^2+n+k-1}$ and by setting $R_{n^2+n+k-1}(e_{n+1} \otimes e_k)$ equal to $\sum_{i=1}^n a_{ni} (e_i \otimes e_k)$. Finally, set R_{n^2+2n} equal to the affine surjection from $D_{(n+1)^2}$ to D_{n^2+2n} which is equal to the identity on D_{n^2+2n} and has $R_{n^2+2n}(e_{n+1} \otimes e_{n+1}) = \sum_{i,j=1}^n a_{ni} b_{nj} e_i \otimes e_j$. We then have P equal to the projective limit of $\{D_m : m \in \mathbb{N}\}$ under $\{R_m : m \in \mathbb{N}\}$.

The projections $\{R_m : m \in \mathbb{N}\}$ have a representing matrix $C = (c_{ij})$ which is triangular and has its rows and columns most conveniently indexed by ordered pairs (i,j) . In this set up the entries in row $(n+1,k)$ for $k=1, \dots, n$ are a_{ni} in column (i,k) and 0 otherwise. The entry in column (k,j) for row $(k,n+1)$ is b_{nj} and 0 otherwise. The entry in row $(n+1,n+1)$ in column (i,j) for $1 \leq i, j \leq n$ is $a_{ni} b_{nj}$ with 0 entries elsewhere. Except for the rows (n,n) each row lies in the subspace of $\ell^1(N \times N) \{x : x_{(2,1)} = 0\}$ or the subspace $\{x : x_{(1,2)} = 0\}$. Since the union of these two subspaces is closed and nowhere dense in the positive face of the unit ball of $\ell^1(N \times N)$ the rows indexed by $\{(n,n) : n \in \mathbb{N}\}$ must be dense in the positive face of the unit ball of $\ell^1(N \times N)$ in order that all of the rows be dense (if we are to have the representing matrix of a Poulsen simplex.)

Let $M_1^+(n,n)$ denote all $n \times n$ matrices (c_{ij}) with $c_{ij} \leq 0$ and $\sum_{i,j=1}^n c_{ij} = 1$. $M_1^+(n,n)$ is naturally embedded in $\ell^1(N \times N)$ by setting $c_{ij} = 0$ if $i > n$ or $j > n$. In order that the (n,n) rows of the representing matrix of $\{R_m : m \in \mathbb{N}\}$ be dense it is necessary and sufficient that for all n every $(c_{ij}) \in M_1^+(n,n)$ be approximable by matrices of the form (d_{ij}) where $d_{ij} = a_{mi} b_{mj}$ $1 \leq i, j \leq n$. This in turn implies that each $(c_{ij}) \in M_1^+(n,n)$ be approximable by matrices $(f_{ij}) \in M_1^+(n,n)$ where $f_{ij} = a_i b_j$ with a_i the i -th row sum and b_j the j -th column sum. The set of such matrices (f_{ij}) is $(2n-2)$ -dimensional whereas $M_1^+(n,n)$ is (n^2-1) -dimensional. Since $n^2-1 > 2n-2$ if $n > 1$ this is impossible. This establishes that $X \otimes Y = P$ is impossible.

COROLLARY. P is not $\bigotimes_{n=1}^{\infty} X_n$ for any sequence of non-zero dimensional simplexes $\{X_n : n \in \mathbb{N}\}$.

PROOF. $\bigotimes_{n=1}^{\infty} X_n = X_1 \otimes \left[\bigotimes_{n=2}^{\infty} X_n \right]$.

REFERENCES

1. Alfsen, E., Convex Compact Sets and Boundary Integrals, Springer, New York, 1971.
2. Asimow, D., Ellis, A.J., Convexity Theory and Its Applications in Functional Analysis, Academic Press, London, 1980.
3. Davies, E.B., Vincent-Smith, G.F., Tensor Products, Infinite Products and Projective Limits of Choquet Simplexes, Math. Scand., 22 (1968), 145-164.
4. Hulanicki, A., Phelps, R.R., Some Applications of Tensor Products of Partially Ordered Linear Spaces, J. Functional Anal., 2 (1968), 177-201.
5. Lazar, A.J., Affine Products of Simplexes, Math. Scand., 22 (1968), 165-175.
6. Lazar, A.J. Lindenstrauss, J., Banach Spaces Whose Duals are L-spaces and Their Representing Matrices, Acta Math., 126 (1971), 165-194.
7. Lindenstrauss, J., Olsen, G., Sternfeld, Y., The Poulsen Simplex, Ann. Inst. Fourier (Grenoble), 28 (1978), 91-114.
8. Lusky, W. Note on the Paper "The Poulsen Simplex" by Lindenstrauss, Olsen and Sternfeld. Ann. Inst. Fourier (Grenoble), 28 (1978), 233-243.
9. Lusky, W., On the Primariness of the Poulsen Simplex Space, Israel J. Math., 37 (1980), 151-163.
10. Namioka, I., Phelps, R.R., Tensor Products of Compact Convex Sets, Pacific J. Math., 31 (1969), 469-480.
11. Poulsen, E.T., A Simplex with Dense Extreme Points, Ann. Inst. Fourier (Grenoble), 11 (1961), 355-370.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

