

ON n^{th} -ORDER DIFFERENTIAL OPERATORS WITH BOHR-NEUGEBAUER TYPE PROPERTY

ARIBINDI SATYANARAYAN RAO

Department of Mathematics, Concordia Univ., Montreal

(Received July 2, 1985 and in revised form July 31, 1986)

ABSTRACT. Suppose B is a bounded linear operator in a Banach space. If the differential operator $\frac{d^n}{dt^n} - B$ has a Bohr-Neugebauer type property for Bochner almost periodic functions, then, for any Stepanov almost periodic continuous function $g(t)$ and any Stepanov-bounded solution of the differential equation $\frac{d^n}{dt^n} u(t) - Bu(t) = g(t)$, $u^{(n-1)}, \dots, u', u$ are all almost periodic.

KEY WORDS AND PHRASES. Bounded linear operator, Bohr-Neugebauer property, Bochner (Stepanov or weakly) almost periodic function, completely continuous normal operator.
1970 AMS SUBJECT CLASSIFICATION SCHEME. PRIMARY 34C25, 34G05; SECONDARY 43A60.

1. INTRODUCTION.

Suppose X is a Banach space and J is the interval $-\infty < t < \infty$. A function $f \in L^p_{\text{loc}}(J; X)$ with $1 \leq p < \infty$ is said to be Stepanov - bounded or S^p -bounded on J if

$$\|f\|_{S^p} = \sup_{t \in J} \left[\int_t^{t+1} \|f(s)\|^p ds \right]^{1/p} < \infty. \quad (1.1)$$

For the definitions of almost periodicity, weak almost periodicity and S^p -almost periodicity, we refer the reader to pp. 3, 39 and 77, Amerio-Prouse [1].

Suppose that B is a bounded linear operator having domain and range in X . We say that the differential operator $\frac{d^n}{dt^n} - B$ has Bohr-Neugebauer property if, for any almost periodic X -valued function $f(t)$ and any bounded (on J) solution of the equation

$$\frac{d^n}{dt^n} u(t) - Bu(t) = f(t) \quad \text{on } J, \quad (1.2)$$

$u^{(n-1)}, \dots, u', u$ are all almost periodic.

Our main result is as follows.

THEOREM 1. For a bounded linear operator B with domain $D(B)$ and range $R(B)$ in a Banach space X , let the differential operator $\frac{d^n}{dt^n} - B$ be such that, for any almost periodic X -valued function $f(t)$ and any S^p -bounded solution $u: J \rightarrow D(B)$ of the equation (1.2), $u^{(n-1)}, \dots, u', u$ are all S^1 -almost periodic. If $p > 1$, then, for any S^p -bounded solution $u: J \rightarrow D(B)$ of the equation

$$\frac{d^n}{dt^n} u(t) - Bu(t) = g(t) \quad \text{on } J, \quad (1.3)$$

$u^{(n-1)}, \dots, u', u$ are all almost periodic.

REMARK 1. Theorem 1 is a generalization of a result of Zaidman [6].

2. PROOF OF THEOREM 1.

By (1.3), we have the representation

$$u^{(n-1)}(t) = u^{(n-1)}(0) + \int_0^t B u(s) ds + \int_0^t g(s) ds \quad \text{on } J. \quad (2.1)$$

If $0 < t_2 - t_1 < 1$ and $p^{-1} + q^{-1} = 1$, then, by the

Hölder's inequality,

$$\begin{aligned} \left\| \int_{t_1}^{t_2} B u(s) ds \right\| &\leq \|B\| \cdot \int_{t_1}^{t_2} \|u(s)\| ds \\ &\leq \|B\| \cdot \left[\int_{t_1}^{t_2} \|u(s)\|^p ds \right]^{p^{-1}} \cdot (t_2 - t_1)^{q^{-1}} \\ &\leq \|B\| \cdot \left[\int_{t_1}^{t_1+1} \|u(s)\|^p ds \right]^{p^{-1}} \cdot (t_2 - t_1)^{q^{-1}} \\ &\leq \|B\| \cdot \|u\|_{S^p} \cdot (t_2 - t_1)^{q^{-1}}. \end{aligned} \quad (2.2)$$

Hence $\int_0^t B u(s) ds$ is uniformly continuous on J . Further, by Theorem 8, p. 79, Amerio-Prouse [1], $\int_0^t g(s) ds$ is uniformly continuous on J . Consequently, $u^{(n-1)}$ is uniformly continuous on J .

Now consider a sequence $\{\rho_k(t)\}_{k=1}^\infty$ of non-negative continuous functions on J such that

$$\rho_k(t) = 0 \quad \text{for } |t| \geq k^{-1}, \quad \int_{-k^{-1}}^{k^{-1}} \rho_k(t) dt = 1. \quad (2.3)$$

The convolution between u and ρ_k is defined by

$$(u * \rho_k)(t) = \int_J u(t-s) \rho_k(s) ds = \int_J u(s) \rho_k(t-s) ds. \quad (2.4)$$

From (1.3), it follows that

$$\frac{d^n}{dt^n} (u * \rho_k)(t) - B(u * \rho_k)(t) = (g * \rho_k)(t) \quad \text{on } J. \quad (2.5)$$

Again by Hölder's inequality,

$$\begin{aligned} \|(u * \rho_k)(t)\| &= \left\| \int_{-1}^1 u(t-s) \rho_k(s) ds \right\| \\ &\leq \left[\int_{-1}^1 \|u(t-s)\|^p ds \right]^{p^{-1}} \cdot \left[\int_{-1}^1 [\rho_k(s)]^q ds \right]^{q^{-1}} \\ &= c_{\rho_k} \left[\int_{t-1}^{t+1} \|u(\sigma)\|^p d\sigma \right]^{p^{-1}} \\ &\leq 2 c_{\rho_k} \|u\|_{S^p} \quad \text{for all } t \in J \text{ and } k = 1, 2, \dots \end{aligned} \quad (2.6)$$

Similarly, the S^1 -almost periodicity of $g(t)$ implies the almost periodicity of $(g * \rho_k)(t)$ for all $k = 1, 2, \dots$.

Consequently, it follows from our assumption on the operator $\frac{d^n}{dt^n} - B$ that $(u * \rho_k)^{(n-1)}(t), \dots, (u * \rho_k)'(t), (u * \rho_k)(t)$ are all S^1 -almost periodic from J to X for all $k \geq 1$.

Further, since $u^{(n-1)}(t)$ is uniformly continuous on J , given $\varepsilon > 0$, there

exists $\delta > 0$ such that

$$\|u^{(n-1)}(t_1) - u^{(n-1)}(t_2)\| \leq \varepsilon \text{ for } |t_1 - t_2| \leq \delta. \quad (2.7)$$

Consequently, we have, for $|t_1 - t_2| \leq \delta$,

$$\begin{aligned} & \| (u^{(n-1)} * \rho_k)(t_1) - (u^{(n-1)} * \rho_k)(t_2) \| \\ & \leq \int_{-k}^{k-1} \| u^{(n-1)}(t_1 - s) - u^{(n-1)}(t_2 - s) \| \rho_k(s) ds \\ & \leq \varepsilon \int_{-k}^{k-1} \rho_k(s) ds = \varepsilon, \text{ by (2.3).} \end{aligned} \quad (2.8)$$

Hence $(u * \rho_k)^{(n-1)}(t) = (u^{(n-1)} * \rho_k)(t)$ is uniformly continuous on J . So, by Theorem 7, p. 78, Amerio-Prouse [1], $(u^{(n-1)} * \rho_k)(t)$ is almost periodic.

Furthermore, by the uniform continuity of $u^{(n-1)}(t)$ on J , the sequence of convolutions $(u^{(n-1)} * \rho_k)(t)$ converges to $u^{(n-1)}(t)$ as $k \rightarrow \infty$, uniformly on J . Hence $u^{(n-1)}(t)$ is almost periodic from J to X , and so is bounded on J . Therefore $u^{(n-2)}(t)$ is uniformly continuous on J . Consequently, $(u^{(n-2)} * \rho_k)(t)$ is almost periodic and $(u^{(n-2)} * \rho_k)(t) \rightarrow u^{(n-2)}(t)$ as $k \rightarrow \infty$, uniformly on J . Hence $u^{(n-2)}(t)$ is almost periodic.

Thus we conclude successively that $u^{(n-1)}, \dots, u', u$ are all almost periodic from J to X , which completes the proof of the theorem.

REMARK 2. The conclusion of Theorem 1 remains valid for any S^1 -bounded and uniformly continuous solution of the equation (1.3).

PROOF. By the Lemma of Rao [5], such a solution is bounded on J . Consequently, by the representation (2.1), $u^{(n-1)}$ is uniformly continuous on J .

REMARK 3. If $B = 0$, then Theorem 1 holds for $p \geq 1$.

3. NOTES.

(i) Suppose X is a separable Hilbert space, and consider the differential equation

$$\frac{d^n}{dt^n} u(t) - Bu(t) = f(t) \text{ on } J, \quad (3.1)$$

where $f : J \rightarrow X$ is an almost periodic function, and $B : X \rightarrow X$ is a completely continuous normal operator. Then, if u is a bounded solution of (3.1), $u^{(n)}$ is almost periodic (as shown in the proof of Theorem 1 of Cooke [3]). Therefore, by the Corollary to Lemma 5 of Cooke [3], $u^{(n-1)}, \dots, u', u$ are all almost periodic. That is, the operator $\frac{d^n}{dt^n} - B$ has Bohr-Neugebauer property.

Now assume that u is an S^p -bounded solution ($1 < p < \infty$) of the equation (3.1). If we replace g by f in the proof of our Theorem 1, then, by the Bohr-Neugebauer property of the operator $\frac{d^n}{dt^n} - B$, it follows that $u^{(n-1)}, \dots, u', u$ are all almost periodic. Hence the operator $\frac{d^n}{dt^n} - B$ satisfies the assumption of Theorem 1 for $p > 1$.

(ii) Finally, suppose X is a reflexive space and $B = 0$. Given an almost periodic X -valued function $f(t)$, assume $u(t)$ is a bounded solution of the differential equation

$$\frac{d^n}{dt^n} u(t) = f(t) \text{ on } J. \quad (3.2)$$

Then it follows from Lemma 2 of Cooke [3] that $u^{(n-1)}, \dots, u'$ are all bounded on J . Hence we conclude successively that $u^{(n-1)}, \dots, u', u$ are all almost periodic (see Amerio-Prouse [1], p. 55 and Authors' Remark on p. 82).

Therefore the operator $\frac{d^n}{dt^n}$ has Bohr-Neugebauer property.

Now, given an S^1 -almost periodic continuous X -valued function $g(t)$, suppose $u(t)$ is an S^p -bounded solution ($1 \leq p < \infty$) of the differential equation

$$\frac{d^n}{dt^n} u(t) = g(t) \quad \text{on } J. \quad (3.3)$$

From (3.3), it follows that

$$\frac{d^n}{dt^n} (u * \rho_k)(t) = (g * \rho_k)(t) \quad \text{on } J, \quad (3.4)$$

where $\{\rho_k(t)\}_{k=1}^\infty$ is the sequence defined in the proof of our Theorem 1. Then $(u * \rho_k)(t)$ is bounded on J and $(g * \rho_k)(t)$ is almost periodic from J to X . So, by the Bohr-Neugebauer property of the operator $\frac{d^n}{dt^n}$, $(u * \rho_k)^{(n-1)}(t), \dots, (u * \rho_k)'(t), (u * \rho_k)(t)$ are all almost periodic.

By (3.3), it follows from Theorem 8, p. 79, Amerio-Prouse [1] that $u^{(n-1)}(t)$ is uniformly continuous on J . Consequently, we conclude successively that $u^{(n-1)}(t), \dots, u'(t), u(t)$ are all almost periodic. Hence the operator $\frac{d^n}{dt^n}$ satisfies the assumption of Theorem 1 for $p \geq 1$.

4. CONSEQUENCES OF THEOREM 1.

Let $L(X, X)$ be the Banach space of all bounded linear operators on X into itself, with the uniform operator topology. As consequences of our Theorem 1, we demonstrate the following results.

THEOREM 2. In a reflexive space X , suppose $f : J \rightarrow X$ is an S^p -almost periodic continuous function ($1 \leq p < \infty$), and $B : J \rightarrow L(X, X)$ is almost periodic with respect to the norm of $L(X, X)$. If $u : J \rightarrow X$ is any S^p -almost periodic solution of the differential equation

$$\frac{d^n}{dt^n} u(t) = B(t)u(t) + f(t) \quad \text{on } J, \quad (4.1)$$

then $u^{(n-1)}, \dots, u', u$ are all almost periodic from J to X .

PROOF. Since $B(t)$ is almost periodic from J to $L(X, X)$, and $u(t)$ is S^p -almost periodic from J to X , we can show that $B(t)u(t)$ is S^p -almost periodic from J to X (see Rao [4]). Hence $B(t)u(t) + f(t)$ is S^p -almost periodic from J to X . If we write

$$v(t) = B(t)u(t) + f(t) \quad \text{on } J, \quad (4.2)$$

then (4.1) becomes

$$\frac{d^n}{dt^n} u(t) = v(t) \quad \text{on } J. \quad (4.3)$$

By our Note (ii), the operator $\frac{d^n}{dt^n}$ satisfies the assumption of our Theorem 1 for $p \geq 1$. Since u is S^p -almost periodic, it is S^p -bounded on J . So, by Theorem 1, $u^{(n-1)}, \dots, u', u$ are all almost periodic.

THEOREM 3. In a reflexive space X , suppose $f : J \rightarrow X$ is an S^p -almost periodic continuous function ($1 \leq p < \infty$), and $B : X \rightarrow X$ is a completely continuous linear operator. If $u : J \rightarrow X$ is a weakly almost periodic (strong) solution of the differential equation

$$\frac{d^n}{dt^n} u(t) = Bu(t) + f(t) \quad \text{on } J, \quad (4.4)$$

then $u^{(n-1)}, \dots, u', u$ are all almost periodic.

PROOF. Since B is a bounded linear operator, Bu is also weakly almost periodic. Further, B being a completely continuous operator, the range of Bu is relatively compact. Hence, by Theorem 10, p. 45, Amerio-Prouse [1], Bu is almost periodic. Consequently, $Bu + f$ is S^p -almost periodic. Now, if we write

$$w(t) = Bu(t) + f(t) \quad \text{on } J, \quad (4.5)$$

then (4.4) becomes

$$\frac{d^n}{dt^n} u(t) = w(t) \quad \text{on } J. \quad (4.6)$$

Since u is weakly almost periodic, it is bounded on J . Therefore, by Theorem 1, $u^{(n-1)}, \dots, u', u$ are all almost periodic.

REMARK 4. Suppose X is a Hilbert space and $B \in L(X, X)$ with $B \geq 0$. Consider the differential equation

$$\frac{d^2}{dt^2} u(t) - Bu(t) = f(t) \quad \text{on } J, \quad (4.7)$$

where $f : J \rightarrow X$ is an almost periodic function. Then any bounded solution $u : J \rightarrow X$ of the equation (4.7) is almost periodic (see Zaidman [7]). By (4.7), $u'(t)$ is uniformly continuous on J . Hence, by Theorem 6, p. 6, Amerio-Prouse [1], $u'(t)$ is almost periodic. Therefore the operator $\frac{d^2}{dt^2} - B$ has Bohr-Neugebauer property, and so satisfies the assumption of Theorem 1 for $p > 1$.

REFERENCES

1. AMERIO, L. and PROUSE, G. Almost Periodic Functions and Functional Equations, Van Nostrand Reinhold Company (1971).
2. BOCHNER, S. and NEUMANN, J.V. On Compact Solutions of Operational-differential Equations I, Ann. of Math., 36 (1935), 255-291.
3. COOKE, R. Almost periodicity of Bounded and Compact Solutions of Differential Equations, Duke Math. J., 36 (1969), 273-276.
4. RAO, A.S. On the Stepanov-almost periodic solution of an abstract differential equation, Indiana Univ. Math. J., 23 (1973), 205-208.
5. RAO, A.S. On the Stepanov-bounded primitive of a Stepanov-almost periodic function, Istit. Lombardo Accad. Sci. Lett. Rend. A., 109 (1975), 65-68.
6. ZAIDMAN, S. A remark on Differential Operators with Bohr-Neugebauer property, Istit. Lombardo Accad. Sci. Lett. Rend. A., 105 (1971), 708-712.
7. ZAIDMAN, S. Soluzioni quasi-periodiche per alcune equazioni differenziali in spazi Hilbertiani, Ric. Mat., 13 (1964), 118-134.

