HOW MANY NUMBERS SATISFY THE $3X + 1$ CONJECTURE?

I. KRASIKOV

School of Mathematical Sciences
Tel-Aviv University
Israel

(Received March 25, 1988 and in revised form December 12, 1988)

ABSTRACT. Let $\theta(x)$ be the number of numbers not exceeding x satisfy the $3X + 1$ conjecture. We obtain a system of difference inequalities on functions closely related to θ . Solving this system in the simplest case, we

establish $\theta(x) > cx^{\frac{3}{7}}$. This improves a result of Crandall [1].

KEY WORDS AND PHRASES. $3X + 1$ conjecture, residue class, difference inequality.

1980 AMS SUBJECT CLASSIFICATION CODES. 11A, 11B.

1. INTRODUCTION.

The famous conjecture of Collatz-Kakutani, also known as the Syracuse or the " $3X + 1$ " problem, claims that the sequence

$$\alpha_{n+1} = T(\alpha_n) = \begin{cases} \frac{3\alpha_n + 1}{2}, & \alpha_n \equiv 1 \pmod{2} \\ \frac{\alpha_n}{2}, & \alpha_n \equiv 0 \pmod{2} \end{cases} \quad (1.1)$$

converges to the cycle (1,2) for any $\alpha_0 \in \mathbb{Z}^+$.

The following well-known heuristic argument serves as an evidence for its validity. Consider T as though it were a random walk. It is natural to suppose that odd and even numbers appear independently, with probability $1/2$ at each jump.

Then $T^{(n)}(\alpha_0)$ should converge since the mathematical expectation

of $\frac{T(\alpha)}{\alpha}$ is about $(\frac{3}{2} \cdot \frac{1}{2})^{1/2} < 1$.

Although this conjecture seems to be intractable at present, some supporting results have been obtained. An interesting review on this problem can be found in [2]. In particular, Crandall [1] proved that the conjecture is true for many values of α_0 . Namely, set $\vartheta(x) = |\{u : T^{(k)}(u) = 1 \text{ for some } k \geq 0 \text{ and } u \leq x\}|$. Thus, $\vartheta(x)$ is just the number of numbers not exceeding x satisfying the conjecture. Then Crandall's result is $\vartheta(x) > cx^r$, for appropriate constants $c, r > 0$. However, his proof gives a very poor value for r , about 0.05.

Here we derive a system of difference inequalities on functions closely related to ϑ (Lemma 4). Solving this system in the simplest case, we

establish $\vartheta(x) > cx^{\frac{3}{7}}$. Actually our proof gives a little more, namely:

given any $v \equiv 1 \text{ or } 2 \pmod{3}$ that is not in a cycle, for all $x > 1$

$$|\{n \leq vx : T^{(k)}(n) = v \text{ for some } k \geq 1\}| > c_0 x^{\frac{3}{7}},$$

where c_0 is a positive constant independent of v .

In some sense the proof may be regarded as an attempt to formalize the above mentioned heuristic argument.

2. RESULTS.

Consider the infinite directed graph G on the vertex set $V = \mathbb{Z}^+$ and the edge set $E = \{(T(v), v)\}$, whose edges are oriented from $T(v)$ to v . Denote by $G(v, x)$ an induced subgraph of G whose vertex set consists of all integers n such that some $T^k(n) = v$ and $T^i(n) \leq x$ for $0 \leq i \leq k$. That is, it consists of all integers n whose trajectory hits v and remains below x the entire time. In particular $G(v, x)$ is the empty set if $x < v$. We also put $G(v) = G(v, \infty)$. Observe that $G(v)$ has at most one cycle since the in degree of each vertex, but may be v , is one. Moreover, if v does not lie in a cycle of G then $G(v)$ is a tree.

Here we prefer to deal with U , the mapping inverse to T , namely:

$$U(\alpha) = \begin{cases} 2\alpha, & \alpha \equiv 0, 1 \pmod{3} \\ 2\alpha \cup \frac{2\alpha - 1}{3}, & \alpha \equiv 2 \pmod{3} \end{cases} \quad (2.1)$$

Since only numbers $\alpha \equiv 2 \pmod{3}$ have two inverses under T , we wish to analyze iterates under $U = T^{-1}$ restricted to integers $\equiv 2 \pmod{3}$. To do this we must consider values of $\alpha \pmod{9}$.

Let S_n be a complete system of residue classes modulo 3^n . We split S_n as follows:

$$S_n = \bigcup_{i=0}^2 R_n^i, \text{ where } \alpha \in R_n^i \Leftrightarrow \alpha \equiv i \pmod{3}.$$

Furthermore, put

$$R_n^2 = Q_n^2 \cup Q_n^5 \cup Q_n^8, \text{ where } \alpha \in Q_n^i \Leftrightarrow \alpha \equiv i \pmod{9}.$$

Obviously, $U: R_n^0 \rightarrow R_n^0$ and $U: R_n^1 \rightarrow R_n^2$. The action of U on R_n^2 can be split into the four following operators:

$$U_1: R_n^2 \rightarrow R_n^2, U_1(\alpha) = 4\alpha$$

$$U_2: Q_n^5 \rightarrow R_{n-1}^0, U_2(\alpha) = \frac{2\alpha - 1}{3}$$

$$U_3: Q_n^2 \rightarrow R_{n-1}^2, U_3(\alpha) = \frac{4\alpha - 2}{3}$$

$$U_4: Q_n^8 \rightarrow R_{n-1}^2, U_4(\alpha) = \frac{2\alpha - 1}{3}$$

The following lemma is an easy exercise in elementary number theory:

LEMMA 1.

(i) U_1 is a bijection $R_n^2 \leftrightarrow R_n^2$. Moreover, if $\alpha \in R_n^2$ then $\ell = 3^{n-1}$ is the smallest positive integer such that $U_1^{(\ell)}(\alpha) = \alpha$.

(ii) U_3 is a bijection $Q_n^2 \leftrightarrow R_{n-1}^2$.

(iii) U_4 is a bijection $Q_n^8 \leftrightarrow R_{n-1}^2$.

The action of U on R_n^0 and R_n^1 is much simpler. Namely, $U: R_n^0 \rightarrow R_n^0$ and

$U: R_n^1 \rightarrow R_n^2$ are bijections. Moreover, since $\alpha \in R_n^0$ implies $U(\alpha) = 2\alpha \in R_n^0$ we get

LEMMA 2. If $v \in R_n^0$ then $G(v)$ is a chain.

Now we define the functions we deal with in this paper.

Let $v \equiv m \pmod{3^n}$. We set $f(v, x) = f_n^m(v, x) = |G(v, x)|$. (The reason for using the redundant notation $f_n^m(v, x)$ instead of $f(v, x)$ is to simplify the statement of the difference inequalities that follow.)

Observe that for $v < x$

$$f_n^m(v, x) = 1 + [\log_2 \frac{x}{v}], m \in R_n^0, \quad (2.2)$$

$$f_n^m(v, x) = 1 + f_n^{2m}(2v, x), m \in R_n^1. \quad (2.3)$$

Furthermore, let $W = \{w\}$ be the set of those vertices of G which do not belong to a cycle. For instance, $U^K(4) \in W$ for all $K > 0$. Then $G(w)$ is a tree and we set

$$\phi_n^m(y) = \inf_{v \in W} f_n^m(v, 2^y v) = \inf\{f(v, 2^y v) : v \in W \text{ and } v \equiv m \pmod{3^n}\}.$$

Note that for any $m \equiv 2 \pmod{3}$ and n , the set $\{v : v \in G(u), v \equiv m \pmod{3^n}\} \neq \emptyset$ because $2^k v$ is in this set and 2 is a primitive root $\pmod{3^n}$ for all n .

LEMMA 3. $\phi_n^m(y)$ is nondecreasing function of y .

PROOF. Obviously, $f_n^m(v, x)$ is a nondecreasing function of x .

Hence, $\phi_n^m(y) = \inf f_n^m(v, 2^y v)$ is nondecreasing function of y .

The following lemma gives important recurrent inequalities on $\phi_n^m(y)$.

LEMMA 4. For $y > 0$,

$$\left\{ \begin{array}{l} \phi_n^m(y) > \phi_n^{4m}(y-2) + \phi_{n-1}^{\frac{4m-2}{3}}(y+\alpha-2), m \in Q_n^2 \\ \phi_n^m(y) > \phi_n^{4m}(y-2) + \phi_{n-1}^{\frac{2v-1}{3}}(y+\alpha-1), m \in Q_n^8 \\ \phi_n^m(y) > \phi_n^{4m}(y-2) + [y+\alpha], m \in Q_n^5 \end{array} \right. \quad (2.4)$$

where $\alpha = \log_2 3 \approx 1.585$ and

$$\phi_{n-1}^m(y) = \min(\phi_n^m(y), \phi_n^{m+3^{n-1}}(y), \phi_n^{m+2 \cdot 3^{n-1}}(y)). \quad (2.5)$$

PROOF. (2.5) follows immediately from the definition of $\phi_n^m(y)$. Let us demonstrate (2.4). If $v \equiv m \pmod{3^n}$, $m \in Q_n^5$ then, by (2.1), if $v < x$,

$$|G(v, x)| > |G(4v, x)| + |G(\frac{2v-1}{3}, x)|.$$

If $\frac{2v-1}{3} \equiv 0 \pmod{3}$ then $G(\frac{2v-1}{3}, x)$ is a chain by lemma 2. Thus, by (2.2), if $v < x$,

$$f_n^m(v, x) = f_n^{4m}(4v, x) + 1 + [\log_2 \frac{3x}{2v-1}].$$

Hence, $\phi_n^m(y) > \phi_n^{4m}(y-2) + [y+\alpha]$.

If $m \in Q_n^8$ then $G(v, x)$ is a forest. Hence,

$$|G(v, x)| = |G(4v, x)| + |G(\frac{2v-1}{3}, x)|$$

By $\frac{2v-1}{3} < \frac{2v}{3}$ and by lemma 3 we get, if $y > 0$ and $x = 2^y v$, then

$$\begin{aligned}\phi_n^m(y) &= \inf f_n^m(v, x) = \inf (f_n^{4m}(4v, x) + f_{n-1}^{\frac{2v-1}{3}}(\frac{2v-1}{3}, x)) > \\ &> \inf f_n^{4m}(4v, x) + \inf f_{n-1}^{\frac{2v-1}{3}}(\frac{2v-1}{3}, x) > \phi_n^{4m}(y-2) + \phi_{n-1}^{\frac{2v-1}{3}}(y+\alpha-1).\end{aligned}$$

The case $m \in Q_n^2$ may be considered similarly to the case $m \in Q_n^8$. We omit the details

THEOREM 1. $\theta(x) > c_2 x^{\frac{3}{7}}$.

PROOF. For $n = 2$ the system (2.4) becomes for $y > 0$,

$$\phi_2^2(y) > \phi_2^8(y-2) + \phi_1^2(y+\alpha-2),$$

$$\phi_2^8(y) > \phi_2^5(y-2) + \phi_1^2(y+\alpha-1),$$

$$\phi_2^5(y) > \phi_2^2(y-2),$$

where $\phi_1^2(y) = \min(\phi_2^2(y), \phi_2^8(y), \phi_2^5(y))$. Observe that $\phi_2^8(y) > \phi_1^2(y)$ for $y > 2$ by

$$\phi_2^8(y) > \phi_2^5(y-2) + \phi_1^2(y+\alpha-1) > \phi_1^2(y),$$

since $\phi_1^2(y+\alpha-1) > \phi_1^2(y)$ and $\phi_2^5(y-2) > 0$ if $y > 2$. Hence,

$$\phi_1^2(y) = \min(\phi_2^2(y), \phi_2^5(y)) > \min(\phi_2^2(y), \phi_2^2(y-2)) = \phi_2^2(y-2).$$

This yields if $y > 6$,

$$\begin{aligned}\phi_2^2(y) &> \phi_2^5(y-4) + \phi_1^2(y+\alpha-1) + \phi_1^2(y+\alpha-2) \\ &> \phi_2^2(y-6) + \phi_1^2(y+\alpha-1) + \phi_1^2(y+\alpha-2) \\ &> \phi_2^2(y-6) + \phi_2^2(y+\alpha-5) + \phi_2^2(y+\alpha-4).\end{aligned}$$

The initial conditions $\phi_2^2(0) = 1$ imply $\phi_2^2(y) > 1$ for $y > 6$, whence one proves by induction on n , that for $n \leq y \leq n+1$, one has $\phi_2^2(y) > c_1 \lambda^y$, where $\lambda \approx 1.3534$ is the

largest root of $1 = \lambda^{-6} + \lambda^{\alpha-5} + \lambda^{\alpha-4}$.

Finally, we obtain $\theta(x) > c_2 x^{\frac{\log_2 \lambda}{7}} > c_2 x^{\frac{3}{7}}$, where $\log_2 \lambda \approx 0.436$.

REMARK. Although system (2.4) seems to be very complicated and we were unable to

solve it for $n \geq 3$, averaging it over all residue classes modulo 3^{n-1} looks much more attractive. Namely, define

$$F_n(y) = 3^{-n+1} \sum_{m \in R_n} 2^{\phi_n^m(y)}.$$

Using lemmas 1 and 4 we get

$$\begin{aligned} 3^{n-1} F_n(y) &= \sum_{m \in R_n} 2^{\phi_n^m(y)} > \sum_{m \in R_n} 2^{\phi_n^m(y-2)} + \sum_{m \in R_{n-1}} 2^{\phi_{n-1}^m(y+\alpha-2)} + \sum_{m \in R_{n-1}} 2^{\phi_{n-1}^m(y+\alpha-1)} = \\ &= 3^{n-1} F_n(y-2) + 3^{n-2} F_{n-1}(y+\alpha-2) + 3^{n-2} F_{n-1}(y+\alpha-1). \end{aligned}$$

Thus,

$$F_n(y) > F_n(y-2) + \frac{1}{3} F_{n-1}(y+\alpha-2) + \frac{1}{3} F_{n-1}(y+\alpha-1).$$

Observe that the associated limit equation $1 = \lambda^{-2} + \frac{1}{3} (\lambda^{\alpha-2} + \lambda^{\alpha-1})$ has $\lambda = 2$ as the smallest positive root. Therefore, one might expect that the solution of the

difference inequalities gives $\theta(x) > c_n x^{r_n}$, where $r_n \rightarrow 1$ when n tends to infinity.

REFERENCES

1. CRANDALL, R.E., On the " $3X + 1$ " problem Math. Comp. **32**, (1972), 1281-1292.
2. LAGARIAS, J.C., The $3X + 1$ problem and its generalizations, The American Math. Monthly **92**, (1985), 3-23.