

## ON CERTAIN BAZILEVIĆ FUNCTIONS OF ORDER $\beta$

SHIGEYOSHI OWA

Department of Mathematics  
Kinki University  
Higashi-Osaka, Osaka 577  
JAPAN

(Received May 8, 1991)

**ABSTRACT.** A certain class  $B(n, \alpha, \beta)$  of Bazilević functions of order  $\beta$  in the unit disk is introduced. The object of the present paper is to derive some properties of functions belonging to the class  $B(n, \alpha, \beta)$ . Our result for the class  $B(n, \alpha, \beta)$  is the improvement of the theorem by N. E. Cho ([1]).

**KEY WORDS AND PHRASES.** Analytic function, class  $B(n, \alpha, \beta)$ , Bazilević function.

**1991 AMS SUBJECT CLASSIFICATION CODE.** Primary 30C45.

### 1. INTRODUCTION.

Let  $A(n)$  denote the class of functions of the form

$$f(z) = z + \sum_{k=n+1}^{\infty} a_k z^k \quad (n \in N = \{1, 2, 3, \dots\}) \quad (1.1)$$

which are analytic in the unit disk  $U = \{z: |z| < 1\}$ . A function  $f(z) \in A(n)$  is said to be a member of the class  $B(n, \alpha, \beta)$  if it satisfies

$$\operatorname{Re} \left\{ \frac{f'(z) f(z)^{\alpha-1}}{z^{\alpha-1}} \right\} > \beta \quad (1.2)$$

for some  $\alpha (\alpha > 0)$ ,  $\beta (0 \leq \beta < 1)$ , and for all  $z \in U$ . We note that  $B(n, \alpha, \beta)$  is the subclass of Bazilević functions in the unit disk  $U$  (cf. [1]). Also we say that  $f(z)$  in the class  $B(n, \alpha, \beta)$  is a Bazilević function of order  $\beta$ .

Recently, Cho [1] has studied the class  $B(n, \alpha, 0)$  when  $\beta = 0$ , and has proved

**THEOREM A.** If  $f(z) \in B(n, 2, 0)$  when  $\alpha = 2$  and  $\beta = 0$ , then

$$\operatorname{Re} \left\{ \frac{f(z)}{z} \right\} > \frac{n}{n+2} \quad (z \in U). \quad (1.3)$$

In the present paper, we improve the above theorem by Cho [1].

### 2. PROPERTIES OF THE CLASS $B(n, \alpha, \beta)$ .

In order to establish our main result, we have to recall here the following lemma due to Miller and Mocanu [2].

**LEMMA.** Let  $\phi(u, v)$  be a complex valued function,

$$\phi: D \rightarrow C, D \subset C^2 \text{ (} C \text{ is the complex plane),}$$

and let  $u = u_1 + iu_2$ ,  $v = v_1 + iv_2$ . Suppose that the function  $\phi(u, v)$  satisfies

- (i)  $\phi(u, v)$  is continuous in  $D$ ;
- (ii)  $(1, 0) \in D$  and  $\operatorname{Re}\{\phi(1, 0)\} > 0$ ;
- (iii) for all  $(iu_2, v_1) \in D$  such that  $v_1 \leq -n(1 + u_2^2)/2$ ,

$$\operatorname{Re}\{\phi(iu_2, v_1)\} \leq 0.$$

Let  $p(z) = 1 + p_n z^n + p_{n+1} z^{n+1} + \dots$  be regular in the unit disk  $U$  such that  $(p(z), zp'(z)) \in D$  for all  $z \in U$ . If

$$\operatorname{Re}\{\phi(p(z), zp'(z))\} > 0 \quad (z \in U),$$

then

$$\operatorname{Re}\{p(z)\} > 0 \quad (z \in U).$$

Using the above lemma, we prove

**THEOREM 1.** If  $f(z) \in B(n, \alpha, \beta)$ , then

$$\operatorname{Re}\left\{\frac{f(z)}{z}\right\}^\alpha > \frac{n + 2\alpha\beta}{n + 2\alpha} \quad (z \in U). \quad (2.1)$$

**PROOF.** We define the function  $p(z)$  by

$$\left\{\frac{f(z)}{z}\right\}^\alpha = \gamma + (1 - \gamma)p(z) \quad (2.2)$$

with

$$\gamma = \frac{n + 2\alpha\beta}{n + 2\alpha}. \quad (2.3)$$

Then, we see that  $p(z) = 1 + p_n z^n + p_{n+1} z^{n+1} + \dots$  is regular in  $U$ .

It follows from (2.2) that

$$\frac{f'(z)f(z)^{\alpha-1}}{z^{\alpha-1}} = \gamma + (1 - \gamma)p(z) + \frac{(1 - \gamma)zp'(z)}{\alpha}, \quad (2.4)$$

or

$$\begin{aligned} & \operatorname{Re}\left\{\frac{f'(z)f(z)^{\alpha-1}}{z^{\alpha-1}} - \beta\right\} \\ &= \operatorname{Re}\left\{\gamma - \beta + (1 - \gamma)p(z) + \frac{(1 - \gamma)zp'(z)}{\alpha}\right\} \\ &> 0. \end{aligned} \quad (2.5)$$

Defining the function  $\phi(u, v)$  by

$$\phi(u, v) = \gamma - \beta + (1 - \gamma)u + \frac{(1 - \gamma)v}{\alpha}, \quad (2.6)$$

(note that  $u = p(z)$  and  $v = zp'(z)$ , we have that

- (i)  $\phi(u, v)$  is continuous in  $D = C^2$ ;
- (ii)  $(1, 0) \in D$  and  $\operatorname{Re}\{\phi(1, 0)\} = 1 - \beta > 0$ ;
- (iii) for all  $(iu_2, v_1)$  such that  $v_1 \leq -n(1 + u_2^2)/2$ ,

$$\begin{aligned} \operatorname{Re}\{\phi(iu_2, v_1)\} &= \gamma - \beta + \frac{(1 - \gamma)v_1}{\alpha} \\ &\leq \gamma - \beta - \frac{n(1 - \gamma)(1 + u_2^2)}{2\alpha} \\ &= -\frac{n(1 - \gamma)u_2^2}{2\alpha} \\ &\leq 0. \end{aligned}$$

Therefore, the function  $\phi(u, v)$  satisfies the conditions in Lemma. This implies that  $\operatorname{Re}\{p(z)\} > 0 (z \in U)$ , which is equivalent to

$$\operatorname{Re}\left\{\frac{f(z)}{z}\right\}^\alpha > \gamma = \frac{n + 2\alpha\beta}{n + 2\alpha} \quad (z \in U). \tag{2.7}$$

This completes the assertion of Theorem 1.

Letting  $\beta = 0$  in Theorem 1, we have

COROLLARY 1. If  $f(z) \in B(n, \alpha, 0)$ , then

$$\operatorname{Re}\left\{\frac{f(z)}{z}\right\}^\alpha > \frac{n}{n + 2\alpha} \quad (z \in U). \tag{2.8}$$

REMARK. If we take  $\alpha = 1$  in Corollary 1, then we have the inequality (1.3) by Cho [1].

Making  $\alpha = 1/2$ , Theorem 1 gives

COROLLARY 2. If  $f(z) \in B(n, 1/2, \beta)$ , then

$$\operatorname{Re}\sqrt{\frac{f(z)}{z}} > \frac{n + \beta}{n + 1} \quad (z \in U). \tag{2.9}$$

Finally, we derive

THEOREM 2. If  $f(z) \in B(n, \alpha, \beta)$ , then

$$\operatorname{Re}\left\{\frac{f(z)}{z}\right\}^{\alpha/2} > \frac{n + \sqrt{n^2 + 4\alpha\beta(n + \alpha)}}{2(n + \alpha)} \quad (z \in U). \tag{2.10}$$

PROOF. Defining the function  $p(z)$  by

$$\left\{\frac{f(z)}{z}\right\}^{\alpha/2} = \gamma + (1 - \gamma)p(z) \tag{2.11}$$

with

$$\gamma = \frac{n + \sqrt{n^2 + 4\alpha\beta(n + \alpha)}}{2(n + \alpha)}, \tag{2.12}$$

we easily see that  $p(z) = 1 + p_n z^n + p_{n+1} z^{n+1} + \dots$  is regular in  $U$ . Taking the differentiations of both sides in (2.11), we obtain that

$$\begin{aligned} &\frac{f'(z)f(z)^{\alpha-1}}{z^{\alpha-1}} \\ &= (\gamma + (1 - \gamma)p(z))^2 + \frac{2}{\alpha}(1 - \gamma)(\gamma + (1 - \gamma)p(z))z p'(z), \end{aligned} \tag{2.13}$$

that is, that

$$\begin{aligned} & \operatorname{Re} \left\{ \frac{f'(z)f(z)^{\alpha-1}}{z^{\alpha-1}} - \beta \right\} \\ &= \operatorname{Re} \left\{ (\gamma + (1-\gamma)p(z))^2 + \frac{2}{\alpha}(1-\gamma)(\gamma + (1-\gamma)p(z))zp'(z) - \beta \right\} \\ &> 0. \end{aligned} \tag{2.14}$$

Therefore, letting

$$\phi(u, v) = (\gamma + (1-\gamma)u)^2 + \frac{2}{\alpha}(1-\gamma)(\gamma + (1-\gamma)u)v - \beta, \tag{2.15}$$

(note that  $p(z) = u = u_1 + iu_2$  and  $zp'(z) = v = v_1 + iv_2$ ), we observe that

- (i)  $\phi(u, v)$  is continuous in  $D = C^2$ ;
- (ii)  $(1, 0) \in D$  and  $\operatorname{Re}\{\phi(1, 0)\} = 1 - \beta > 0$ ;
- (iii) for all  $(iu_2, v_1) \in D$  such that  $v_1 \leq -n(1 + u_2^2)/2$ ,

$$\begin{aligned} \operatorname{Re}\{\phi(iu_2, v_1)\} &= \gamma^2 - (1-\gamma)^2u_2^2 + \frac{2}{\alpha}\gamma(1-\gamma)v_1 - \beta \\ &\leq \gamma^2 - \beta - (1-\gamma)^2u_2^2 - \frac{n}{\alpha}\gamma(1-\gamma)(1 + u_2^2) \\ &\leq 0. \end{aligned}$$

Thus, the function  $\phi(u, v)$  satisfies the conditions in Lemma. Applying Lemma, we conclude that

$$\operatorname{Re} \left\{ \frac{f(z)}{z} \right\}^{\alpha/2} > \gamma = \frac{n + \sqrt{n^2 + 4\alpha\beta(n + \alpha)}}{2(n + \alpha)} \quad (z \in U). \tag{2.16}$$

Taking  $\alpha = 1$  in Theorem 2, we have


**COROLLARY 3.** If  $f(z) \in B(n, 1, \beta)$ , then

$$\operatorname{Re} \sqrt{\frac{f(z)}{z}} > \frac{n + \sqrt{n^2 + 4n\beta + 4\beta}}{2(n + 1)} \quad (z \in U). \tag{2.17}$$

**REMARK.** If we take  $\alpha = 2$  and  $\beta = 0$  in Theorem 2, then we have Theorem A by Cho [1].

#### REFERENCES

1. CHO, N.E., On certain subclasses of univalent functions, Bull. Korean Math. Soc. **25** (1988), 215-219.
2. MILLER, S.S. and MOCANU, P.T., Second order differential inequalities in the complex plane, J. Math. Anal. Appl. **65** (1978), 289-305.
3. SINGH, R., On Bazilevič functions, Proc. Amer. Math. Soc. **38** (1973), 261-271.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

