

## AN APPLICATION OF KKM-MAP PRINCIPLE

A. CARBONE

Dipartimento di Matematica  
Università degli studi della Calabria  
87036 Arcavacata di Rende (Cosenza) - ITALY

(Received February 4, 1991 and in revised form June 6, 1991)

**ABSTRACT.** The following theorem is proved and several fixed point theorems and coincidence theorems are derived as corollaries. Let  $C$  be a nonempty convex subset of a normed linear space  $X$ ,  $f: C \rightarrow X$  a continuous function,  $g: C \rightarrow C$  continuous, onto and almost quasi-convex. Assume that  $C$  has a nonempty compact convex subset  $D$  such that the set

$$A = \{y \in C: \|g(x) - f(y)\| \geq \|g(y) - f(y)\| \text{ for all } x \in D\}$$

is compact.

Then there is a point  $y_0 \in C$  such that  $\|g(y_0) - f(y_0)\| = d(f(y_0), C)$ .

**KEY WORDS AND PHRASES.** Almost quasi-convex functions, fixed points, coincidence points.  
**1991 AMS SUBJECT CLASSIFICATION CODE.** Primary 47 H 10, Secondary 54 H 25.

### 1. INTRODUCTION.

There has been given a variety of applications of KKM-map principle by Ky Fan [1] in areas like fixed point theory, approximation theory, minimax theory, potential theory and variational problems. For further applications we refer to [2].

Recently Prolla [4] proved the following result using fixed point theorems for multivalued mappings. In this note we extend his theorem and our proof will follow KKM-map principle.

Let  $C$  be a compact, convex subset of a Banach space  $X$ ,  $f: C \rightarrow X$  a continuous function and  $g: C \rightarrow C$  a continuous, almost affine and onto map. Then there is a  $y_0 \in C$  such that

$$\|g(y_0) - f(y_0)\| = d(f(y_0), C).$$

Recall that a map  $g: C \rightarrow X$  is almost affine if

$$\|g(\lambda x_1 + (1 - \lambda)x_2) - y\| \leq \lambda \|g(x_1) - y\| + (1 - \lambda) \|g(x_2) - y\|$$

for all  $x_1, x_2 \in C$  and  $y \in X$ .

Clearly a linear map is almost affine, but not conversely.

We have taken an almost quasi-convex map  $g$ .

**DEFINITION.** A map  $g: C \rightarrow X$  is said to be almost quasi-convex if, for every  $t \in X$  and  $r > 0$ , the set  $\{u \in C: \|g(u) - t\| < r\}$  is convex.

An almost quasi-convex condition is more general than almost affine condition.

We use the following well-known result (Lin [3]) to derive our theorem given below.

**THEOREM 1.1.** Let  $C$  be a nonempty convex subset of a topological vector space. Let  $B \subset C \times C$  be such that

- i) for each  $x \in C$ , the set  $\{y \in C : (x, y) \in B\}$  is closed in  $C$ ;
- ii) for each  $y \in C$  the set  $\{x \in C : (x, y) \notin B\}$  is empty or convex;
- iii)  $(x, x) \in B$  for each  $x \in C$ ; and
- iv)  $C$  has a nonempty compact convex subset  $D$  such that the set

$$A = \{y \in C : (x, y) \in B \text{ for all } x \in D\}$$

is compact.

Then there exists a point  $y_0 \in C$  such that  $C \times \{y_0\} \subseteq B$ .

**2. MAIN RESULTS.**

Now we prove our results.

**THEOREM 2.1.** Let  $C$  be a nonempty convex subset of a normed linear space  $X$ ,  $f: C \rightarrow X$  a continuous function,  $g: C \rightarrow C$  continuous, onto and almost quasi-convex function. (\*) Assume that  $C$  has a nonempty compact convex subset  $D$  such that the set

$$A = \{y \in C : \|g(x) - f(y)\| \geq \|g(y) - f(y)\| \text{ for all } x \in D\}$$

is compact.

Then there is a point  $y_0 \in C$  such that  $\|g(y_0) - f(y_0)\| = d(f(y_0), C)$ .

**PROOF.** Set

$$B = \{(x, y) \in C \times C : \|g(x) - f(y)\| \geq \|g(y) - f(y)\|\}.$$

Then the set  $\{y \in C : (x, y) \in B\}$  is closed in  $C$  since  $f$  and  $g$  are continuous. It is easy to see that  $(x, x) \in B$  for each  $x \in C$ .

We have to show that the set

$$M = \{x \in C : (x, y) \notin B\} = \{x \in C : \|g(x) - f(y)\| < \|g(y) - f(y)\|\}$$

is convex or empty.

Since  $g$  is an almost quasi-convex function, therefore  $M$  is convex.

By Theorem 1.1 we get that there is a point  $y_0 \in C$  such that

$$\|g(y_0) - f(y_0)\| = d(f(y_0), C).$$

In case the convex set  $C$  is compact we may take  $C = D$ .

**NOTE.** Condition (\*) is equivalent to the following.

Let  $D$  be a nonempty compact convex subset of  $C$ ,  $K$  be a nonempty compact subset of  $C$  such that for each  $y \in C \setminus K$  there exists an  $x_0 \in D$  such that

$$\|g(x_0) - f(y)\| < \|g(y) - f(y)\|.$$

If  $C = K = D$  and  $g$  is almost affine then we get Prolla's result stated below. Let  $C$  be a compact convex subset of a normed linear space  $X$  and  $f: C \rightarrow X$  a continuous function. Let  $g: C \rightarrow X$  be a continuous, onto and almost affine map. Then there exists a  $y_0 \in C$  such that

$$\|g(y_0) - f(y_0)\| = d(f(y_0), C).$$

**NOTE.** (i) If  $f(y_0) \in C$  then we get a coincidence result; and (ii) If  $g = I$ , an identity function, then the above result is a well-known theorem due to by Ky Fan [1]. This theorem has interesting applications in fixed point theory, approximation theory and variational problems. We give a sample application in fixed point theory.

**EXAMPLE:** Let  $C$  be a compact convex subset of a normed linear space  $X$  and  $f: C \rightarrow X$  a continuous map. If  $f(x) \neq x$ , then assume that the line segment  $[x, f(x)]$  has at least two elements of  $C$ . Then  $f$  has a fixed point.

By taking  $g = I$ , we get there is a  $y_0 \in C$  such that

$$\|y_0 - f(y_0)\| = d(f(y_0), C).$$

Now, if  $y_0 \neq f(y_0)$  then there is a  $z \in C$  such that

$$z = \lambda f(y_0) + (1 - \lambda)y_0, \quad 0 < \lambda < 1$$

and

$$\begin{aligned} \|y_0 - f(y_0)\| &\leq \|z - f(y_0)\| = \|\lambda f(y_0) + (1 - \lambda)y_0 - f(y_0)\| \\ &= (1 - \lambda) \|f(y_0) - y_0\| < \|f(y_0) - y_0\| \end{aligned}$$

a contradiction, so  $y_0 = f(y_0)$ .

We could derive several other interesting results on fixed point theorems as corollaries.

**ACKNOWLEDGEMENT.** I wish to thank the Referee for suggesting the definition of almost quasi-convexity that improved the presentation of this paper.

#### REFERENCES

1. FAN, K., Extensions of two fixed point theorems of F.E. Browder, Math. Z. **112** (1969), 234-240.
2. LIN, B.L. and SIMONS, S., Nonlinear and Convex Analysis, Proceedings in Honor of Ky Fan, Marcel Dekker, New York 1987.
3. LIN, T.C., Convex sets, fixed points, variational and minimax inequalities, Bull. Austr. Math. Soc., **34** (1986), 107-117.
4. PROLLA, J.B., Fixed point theorems for set-valued mappings and existence of best approximants, Numer. Funct. Anal. Optimiz. **5** (1982-83), 449-445.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

