

OBLIQUE INTERFACE-WAVE DIFFRACTION BY A SMALL BOTTOM DEFORMATION IN TWO SUPERPOSED FLUIDS

B. N. MANDAL

Physical and Earth Sciences Division
Indian Statistical Institute
203, Barrackpore Trunk Road
Calcutta 700035, INDIA

U. BASU

Department of Applied Mathematics
University of Calcutta
92, A P C Road
Calcutta 700009, INDIA

(Received August 5, 1993 and in revised form December 12, 1994)

ABSTRACT. The problem of diffraction of oblique interface-waves by a small bottom deformation of the lower fluid in two superposed fluids has been investigated here assuming linear theory and invoking a simplified perturbation analysis. First order corrections to the velocity potentials in the two fluids are obtained by using the Green's integral theorem in a suitable manner. The transmission and reflection coefficients are evaluated approximately. These reduce to the known results for a single fluid in the absence of the upper fluid.

KEY WORDS AND PHRASES. Oblique interface-wave diffraction, transmission and reflection coefficients, water waves

1991 AMS SUBJECT CLASSIFICATION CODE. 76C10, 76D33

1. INTRODUCTION.

A time-harmonic progressive wave train propagating on the surface of an ocean experiences no reflection if the ocean is of uniform finite depth. However, if the bottom of the ocean has a deformation, then the wave train is partially reflected by and partially transmitted over the bottom deformation. Miles [1] obtained approximately the transmission and reflection coefficients for oblique surface-waves when the bottom has a small deformation in the form of a long cylinder in the lateral direction. Mandal and Basu [2] extended this problem to include surface tension effect at the free surface.

In the present paper the oblique surface-wave diffraction problem considered in [1] for a single fluid is generalized to two superposed fluids wherein the upper fluid extends infinitely upwards and the lower fluid is of finite but nonuniform depth below the mean interface and its bottom has a small deformation in the form of a long cylinder in the lateral direction. Utilizing a simplified perturbational analysis directly to the governing partial differential equation and the boundary and other conditions describing the physical problem, the original boundary value problem (BVP) is reduced up to first order to another BVP. Solution of this BVP is then obtained by an appropriate use of Green's integral theorem to the potential functions describing the BVP and source potentials given in [3]. The first order corrections to the reflection and transmission coefficients are then evaluated from the requirements at infinity. It is verified that in the absence of the upper fluid, known results for a single fluid are recovered.

It may be mentioned here that although the source potentials in each of two superposed fluids due to various types of basic singularities are known in the literature (cf [3],[4],[5]), their application in the study of interface-waves in two superposed fluids is rather limited. In the present paper, the two-dimensional source potentials in each of two superposed fluids obtained earlier in [3] have been used suitably in the Green's integral theorem to obtain representations for the first order corrections to the velocity potentials in each of the two fluids.

2. FORMULATION OF THE PROBLEM

We consider two superposed immiscible, nonviscous, incompressible and homogeneous fluids with lower fluid of density ρ_1 occupying the region $0 \leq y \leq h + \epsilon c(x)$ and upper fluid of density $\rho_2 (< \rho_1)$ occupying the region $y \leq 0$. Here, the plane $y = 0$ is the position of the interface at rest and y -axis is taken vertically downwards into the lower fluid. There is a small deformation at the bottom of the lower fluid in the lateral direction and is described by $y = h + \epsilon c(x)$ where $c(x)$ is a bounded and continuous function of compact support so that $c(x) \rightarrow 0$ as $|x| \rightarrow \infty$, and ϵ is a small positive number characterizing the smallness of the deformation. Far away from the deformation, the lower fluid is of uniform finite depth h . The motion in each fluid is assumed to be small and irrotational so that it is described by the velocity potentials $Re\{\Phi(x, y, z)e^{-i\sigma t}\}$ and $Re\{\Psi(x, y, z)e^{-i\sigma t}\}$ in the lower and upper fluids respectively, σ being the frequency of the incoming train of progressive waves at the interface and the time dependence $e^{-i\sigma t}$ being dropped throughout the analysis. Assuming linear theory, Φ and Ψ satisfy the following coupled BVP:

$$\nabla_1^2 \Phi = 0 \quad \text{in the region} \quad 0 \leq y \leq h + \epsilon c(x), \tag{2.1}$$

$$\nabla_1^2 \Psi = 0 \quad \text{in the region} \quad y \leq 0 \tag{2.2}$$

where ∇_1^2 is the three-dimensional Laplace operator,

$$\Phi_y = \Psi_y \quad \text{on} \quad y = 0, \tag{2.3}$$

$$K\Phi + \Phi_y = s(K\Psi + \Psi_y) \quad \text{on} \quad y = 0 \tag{2.4}$$

where $s = \rho_2/\rho_1$, $K = \sigma^2/g$, g being the acceleration due to gravity,

$$\Phi_n = 0 \quad \text{on} \quad y = h + \epsilon c(x) \tag{2.5}$$

where n denotes the inward drawn normal to the bottom,

$$\nabla_1 \Psi \rightarrow 0 \quad \text{as} \quad y \rightarrow \infty. \tag{2.6}$$

A train of progressive interface-waves represented by the velocity potentials $\phi_0(x, y)e^{i\nu z}$ and $\Psi_0(x, y)e^{i\nu z}$ in the lower and upper fluids respectively where

$$\phi_0(x, y) = \frac{\cosh\alpha(h - y)}{\sinh\alpha h} e^{i\mu x}, \tag{2.7}$$

$$\Psi_0(x, y) = -e^{\alpha y + i\mu x}, \tag{2.8}$$

is obliquely incident upon the bottom deformation from negative infinity. Here α is the unique positive zero of $\Delta(k)$ where

$$\Delta(k) = K \cosh kh + \{s(K + k) - k\} \sinh kh, \tag{2.9}$$

and $\nu = \alpha \sin \theta$, $\mu = \alpha \cos \theta$ where θ characterizes the oblique incidence of the wave train. $\theta = 0$ corresponds to normal incidence. This wave train is partially reflected by and partially transmitted over the bottom deformation so that Φ and Ψ satisfy the infinity requirements

$$\begin{bmatrix} \Phi \\ \Psi \end{bmatrix} \sim \begin{cases} T \begin{bmatrix} \phi_0(x, y) \\ \psi_0(x, y) \end{bmatrix} e^{i\nu z} & \text{as } x \rightarrow \infty, \\ \begin{bmatrix} \phi_0(x, y) \\ \psi_0(x, y) \end{bmatrix} e^{i\nu z} + R \begin{bmatrix} \phi_0(-x, y) \\ \psi_0(-x, y) \end{bmatrix} e^{i\nu z} & \text{as } x \rightarrow -\infty, \end{cases} \quad (2.10)$$

where T and R denote respectively the unknown transmission and reflection coefficients

Assuming ϵ to be very small, the bottom condition (2.5) can be expressed in approximate form [2] as

$$-\Phi_y + \epsilon\{c'(x)\Phi_x - c(x)\Phi_{yy}\} + O(\epsilon^2) = 0. \quad (2.12)$$

In view of the geometry of the problems we can assume

$$\begin{aligned} \Phi(x, y, z) &= \phi(x, y)e^{i\nu z} \\ \Psi(x, y, z) &= \psi(x, y)e^{i\nu z} \end{aligned} \quad (2.13)$$

so that $\phi(x, y)$ and $\psi(x, y)$ satisfy the following BVP

$$(\nabla^2 - \nu^2)\phi = 0 \quad \text{in } 0 \leq y \leq h + \epsilon c(x), \quad (2.14)$$

$$(\nabla^2 - \nu^2)\psi = 0 \quad \text{in } y \leq 0 \quad (2.15)$$

where ∇^2 is the two-dimensional Laplace operator,

$$\phi_y = \psi_y \quad \text{on } y = 0, \quad (2.16)$$

$$K\phi + \phi_y = s(K\psi + \psi_y) \quad \text{on } y = 0, \quad (2.17)$$

$$-\phi_y + \epsilon\left\{\frac{d}{dx}(c(x)\phi_x) - \nu^2\phi\right\} + O(\epsilon^2) = 0 \quad \text{on } y = h \quad (2.18)$$

since ϕ satisfies (2.14), and

$$\nabla^2\psi \rightarrow 0 \quad \text{as } y \rightarrow -\infty. \quad (2.19)$$

Also ϕ, ψ satisfy the infinity requirements

$$\begin{bmatrix} \phi \\ \psi \end{bmatrix} \sim \begin{cases} T \begin{bmatrix} \phi_0(x, y) \\ \psi_0(x, y) \end{bmatrix} & \text{as } x \rightarrow \infty, \\ \begin{bmatrix} \phi_0(x, y) \\ \psi_0(x, y) \end{bmatrix} + R \begin{bmatrix} \phi_0(-x, y) \\ \psi_0(-x, y) \end{bmatrix} & \text{as } x \rightarrow -\infty. \end{cases} \quad (2.20)$$

3. METHOD OF SOLUTION

In view of the approximate bottom condition (2.18) coupled with the fact that an interface-wave train experiences no reflection if the lower fluid has a uniform bottom, we can assume a perturbation expansion for ϕ, ψ, T and R in terms of ϵ as

$$\begin{aligned} \phi &= \phi_0 + \epsilon\phi_1 + O(\epsilon^2), \\ \psi &= \psi_0 + \epsilon\psi_1 + O(\epsilon^2), \\ T &= 1 + \epsilon T_1 + O(\epsilon^2), \\ R &= \epsilon R_1 + O(\epsilon^2). \end{aligned} \quad (3.1)$$

Using the expansions (3.1) in (2.14) to (2.21) we find that ϕ_1, ψ_1 satisfy the following BVP

$$(\nabla^2 - \nu^2)\phi_1 = 0 \quad \text{in } 0 \leq y \leq h, \quad (3.2)$$

$$(\nabla^2 - \nu^2)\psi_1 = 0 \quad \text{in} \quad y \leq 0, \tag{3 3}$$

$$\phi_{1y} = \psi_{1y} \quad \text{on} \quad y = 0, \tag{3 4}$$

$$K\phi_1 + \phi_{1y} = s(K\phi_2 + \phi_{2y}) \quad \text{on} \quad y = 0, \tag{3 5}$$

$$\phi_{1y} = q(x) \quad \text{on} \quad y = h \tag{3 6}$$

where

$$q(x) = \frac{1}{\sinh \alpha h} \left[\nu \mu \frac{d}{dx} (c(x)e^{\mu x}) - \nu^2 c(x)e^{\mu x} \right]$$

Also, ϕ_1, ψ_1 satisfy the following infinity requirements,

$$\begin{cases} \left[\begin{matrix} \phi_1 \\ \psi_1 \end{matrix} \right] \sim \begin{cases} T_1 \begin{bmatrix} \phi_0(x, y) \\ \psi_0(x, y) \end{bmatrix} & \text{as } x \rightarrow \infty, \\ R_1 \begin{bmatrix} \phi_0(x, y) \\ \psi_0(x, y) \end{bmatrix} & \text{as } x \rightarrow -\infty, \end{cases} \end{cases} \tag{3 7}$$

$$\tag{3 8}$$

To solve the above coupled BVP, we need two-dimensional source potentials for the modified Helmholtz's equation due to a line source submerged in either of two superposed fluids wherein the lower fluid is of uniform finite depth below the mean interface $y = 0$ and the upper fluid extends infinitely upwards. When the source is submerged in the lower fluid at $(\xi, \eta) (0 < \eta < h)$, let $G(x, y; \xi, \eta)$ and $H(x, y; \xi, \eta)$ denote the source potentials in the lower and upper fluids respectively, and when the source is submerged in the upper fluid at $(\xi, \eta) (\eta < 0)$, let $\bar{G}(x, y; \xi, \eta)$ and $\bar{H}(x, y; \xi, \eta)$ denote the source potentials in the lower and upper fluids respectively. Expressions for these source potentials and their asymptotic behaviors as $|x - \xi| \rightarrow \infty$ are given in [3] and are reproduced in the Appendix after correcting the misprints.

To find $\phi_1(\xi, \eta) (0 < \eta < h)$ we apply Green's integral theorem to $\phi_1(x, y)$ and $G(x, y; \xi, \eta)$ in the region bounded externally by the lines $y = 0 (-X \leq x \leq X)$, $x = \pm X (0 \leq y \leq h)$, $y = h (-X \leq x \leq X)$ and internally by the circle C with center at (ξ, η) and radius δ , and ultimately make $X \rightarrow \infty$ and $\delta \rightarrow 0$. We then obtain

$$2\pi\phi_1(\xi, \eta) = \int_{-\infty}^{\infty} q(x)G(x, h; \xi, \eta)dx + \int_{-\infty}^{\infty} [\phi_1 G_y - G\phi_{1y}]_{y=0} dx. \tag{3 9a}$$

Again, we apply the Green's integral theorem to $\psi_1(x, y)$ and $H(x, y; \xi, \eta)$ in the region bounded externally by the lines $y = 0 (-X \leq x \leq X)$, $x = \pm X (-Y \leq y \leq 0)$, $y = Y (-X \leq x \leq X)$ and ultimately make $X, Y \rightarrow \infty$. Here we note that $H(x, y; \xi, \eta)$ has no singularity in the region. Then we find

$$0 = \int_{-\infty}^{\infty} [\psi_1 H_y - H\psi_{1y}]_{y=0} dx. \tag{3 9b}$$

Multiplying (3 9b) by s and subtracting from (3 9a) we find

$$2\pi\phi_1(\xi, \eta) = \int_{-\infty}^{\infty} q(x)G(x, h; \xi, \eta)dx + \int_{-\infty}^{\infty} [(\phi_1 G_y - G\phi_{1y}) - s(\psi_1 H_y - H\psi_{1y})]_{y=0} dx.$$

Using the conditions (3 4) and (3 5) for ϕ_1 and ψ_1 and the conditions on $y = 0$ for G and H given in the Appendix, we find that on $y = 0$,

$$\begin{aligned} \phi_1 G_y - s\psi_1 H_y &= (\phi_1 - s\psi_1)G_y = \frac{s-1}{K} \phi_{1y} G_y, \\ G\phi_{1y} - sH\psi_{1y} &= (G - sH)\phi_{1y} = \frac{s-1}{K} G_y \phi_{1y}. \end{aligned}$$

Thus the term within the square bracket in the second integral vanishes identically. Hence we obtain

$$\phi_1(\xi, \eta) = \frac{1}{2\pi} \int_{-\infty}^{\infty} q(x)G(x, h; \xi, \eta)dx, \quad 0 < \eta < h. \tag{3 10}$$

To find $\psi_1(\xi, \eta)(\eta < 0)$ we apply Green's integral theorem to $\psi_1(x, y)$ and $\bar{H}(x, y; \xi, \eta)$ in the region bounded externally by the lines $y = 0(-X \leq x \leq X)$, $x = \pm X(-Y \leq y \leq 0)$, $y = Y(-X \leq x \leq X)$ and internally by the circle C' of radius δ with center at (ξ, η) and ultimately let $X, Y \rightarrow \infty$ and $\delta \rightarrow 0$. We then find

$$2\pi \psi_1(\xi, \eta) = - \int_{-\infty}^{\infty} [\psi_1 \bar{H}_y - \bar{H} \psi_{1y}]_{y=0} dx. \tag{3 11a}$$

Again, we apply the Green's integral theorem to $\phi_1(x, y)$ and $\bar{G}(x, y; \xi, \eta)(\eta < 0)$ in the region bounded externally by the lines $y = 0(-X \leq x \leq X)$, $x = \pm X(0 \leq y \leq h)$, $y = h(-X \leq x \leq X)$ and ultimately make $X \rightarrow \infty$. Noting that \bar{G} has no singularity in the region we find

$$0 = \int_{-\infty}^{\infty} [\phi_1 \bar{G}_y - \phi_{1y} \bar{G}]_{y=0} dx + \int_{-\infty}^{\infty} q(x) \bar{G}(x, h; \xi, \eta)dx. \tag{3 11b}$$

Multiplying (3 11a) by s and adding with (3 11b) we obtain

$$2\pi s \psi_1(\xi, \eta) = \int_{-\infty}^{\infty} [(\phi_1 \bar{G}_y - \phi_{1y} \bar{G}) - s(\psi_1 \bar{H}_y - \psi_{1y} \bar{H})]_{y=0} dx + \int_{-\infty}^{\infty} q(x) \bar{G}(x, h; \xi, \eta)dx. \tag{3 12}$$

The term in the square bracket of the second integral vanishes because of the conditions satisfied by ϕ_1, ψ_1 and \bar{G}, \bar{H} at $y = 0$. Thus we find

$$\psi_1(\xi, \eta) = \frac{1}{2\pi s} \int_{-\infty}^{\infty} q(x) \bar{G}(x, h; \xi, \eta)dx, \quad \eta < 0. \tag{3 13}$$

4. EVALUATION OF T_1 AND R_1

T_1 and R_1 can be evaluated from the behavior of $\phi_1(\xi, \eta)$ or $\psi_1(\xi, \eta)$ as $\xi \rightarrow \infty$ and $-\infty$ respectively in (3.10) or (3 13). To find T_1 we note from (3.7) that

$$\phi_1(\xi, \eta) \sim T_1 \phi_0(\xi, \eta) \quad \text{as} \quad \xi \rightarrow \infty.$$

Also from (3.10) after using (A3) we find as $\xi \rightarrow \infty$

$$\phi_1(\xi, \eta) \sim \left[\frac{i \sinh \alpha h}{\mu(h + \frac{1-s}{K} \sinh^2 \alpha h)} \int_{-\infty}^{\infty} e^{-i\mu x} q(x)dx \right] \phi_0(\xi, \eta).$$

Thus

$$\begin{aligned} T_1 &= \frac{i \sinh \alpha h}{\mu(h + \frac{1-s}{K} \sinh^2 \alpha h)} \int_{-\infty}^{\infty} e^{-i\mu x} q(x)dx \\ &= - \frac{i\alpha \sec \theta}{h + \frac{1-s}{K} \sinh^2 \alpha h} \int_{-\infty}^{\infty} c(x)dx. \end{aligned} \tag{4 1}$$

It is verified that the same expression for T_1 is also obtained by noting the behavior of $\psi_1(\xi, \eta)$ as $\xi \rightarrow \infty$ in (3 7) and (3 13).

Again, to obtain R_1 , we note from (3 8)

$$\phi_1(\xi, \eta) \sim R_1 \phi_0(-\xi, \eta) \quad \text{as} \quad \xi \rightarrow -\infty$$

and using (A3) in (3.10) we find

$$\phi_1(\xi, \eta) \sim \left[\frac{i \sinh \alpha h}{\mu(h + \frac{1-s}{K} \sinh^2 \alpha h)} \int_{-\infty}^{\infty} e^{i\mu x} q(x) dx \right] \phi_0(-\xi, \eta) \quad \text{as} \quad \xi \rightarrow -\infty.$$

Thus

$$\begin{aligned} R_1 &= \frac{i \sinh \alpha h}{\mu(h + \frac{1-s}{K} \sinh^2 \alpha h)} \int_{-\infty}^{\infty} e^{i\mu x} q(x) dx \\ &= \frac{i\alpha \sec \theta \cos 2\theta}{h + \frac{1-s}{K} \sinh^2 \alpha h} \int_{-\infty}^{\infty} c(x) e^{2i\mu x} dx. \end{aligned} \tag{4.2}$$

It is again verified that the same expression for R_1 is also obtained by noting the behavior of $\psi_1(\xi, \eta)$ as $\xi \rightarrow -\infty$ in (3.8) and (3.13)

It may be noted that in the absence of the upper fluid ($s = 0$), the results of [1] for a single fluid are recovered. In that case α is the unique real positive zero of $\Delta(k) = K \cosh \alpha h - k \sinh kh$. The results for normal incidence of the wave train are obtained by putting $\theta = 0$. For $\theta = \pi/4$, R_1 vanishes independently of the bottom deformation. This was also observed by Miles [1] for a single fluid. Also, once the functional form of $c(x)$ is known, T_1 and R_1 can be obtained explicitly.

APPENDIX

(a) $G(x, y; \xi, \eta)$ and $H(x, y; \xi, \eta)$

$$\begin{aligned} &G(x, y; \xi, \eta) \text{ and } H(x, y; \xi, \eta) \text{ satisfy} \\ &(\nabla^2 - \nu^2)G = 0 \text{ in } 0 \leq y \leq h \text{ except at } (\xi, \eta) (0 < \eta < h), \\ &G \sim K_0(\nu r) \text{ as } r = \{(x - \xi)^2 + (y - \eta)^2\}^{1/2} \rightarrow 0, \\ &(\nabla^2 - \nu^2)H = 0 \text{ in } y \leq 0, \\ &G_y = H_y \text{ on } y = 0, \\ &KG + G_y = s(KH + H_y) \text{ on } y = 0, \\ &G_y = 0 \text{ on } y = h, \quad \nabla H \rightarrow 0 \text{ as } y \rightarrow -\infty, \end{aligned}$$

G, H have outgoing nature as $|x - \xi| \rightarrow \infty$. Then $G(x, y; \xi, \eta)$ and $H(x, y; \xi, \eta)$ are given by (cf [3]) after correcting some misprints

$$\begin{aligned} G(x, y; \xi, \eta) &= K_0(\nu r) - \frac{1-s}{1+s} K_0(\nu r') \\ &+ \frac{2}{1+s} \int_{\nu}^{\infty} \left[\frac{e^{-kh} \{s(K+k) - k\} (\sinh k\eta + s \cosh k\eta) \operatorname{sech} kh - k(1-s)e^{-k\eta}}{\Delta(k)} \right. \\ &\times \left. \cosh k(h-y) + \frac{e^{-kh} (\sinh k\eta + s \cosh k\eta)}{\cosh kh} \sinh ky \right] \frac{\cos\{(k^2 - \nu^2)^{1/2}(x - \xi)\}}{(k^2 - \nu^2)^{1/2}} dk, \tag{A1} \end{aligned}$$

$$\begin{aligned} H(x, y; \xi, \eta) &= \frac{2}{1+s} K_0(\nu r) + \frac{2}{1+s} \int_{\nu}^{\infty} \left[\frac{\sinh k\eta + s \cosh k\eta}{\cosh kh} e^{-kh} \right. \\ &- \left. \frac{e^{-kh} \{s(K+k) - k\} (\sinh k\eta + s \cosh k\eta) \operatorname{sech} kh - k(1-s)e^{-k\eta}}{\Delta(k)} \sinh kh \right] \\ &\times e^{ky} \frac{\cos\{(k^2 - \nu^2)^{1/2}(x - \xi)\}}{(k^2 - \nu^2)^{1/2}} dk, \tag{A2} \end{aligned}$$

where $r' = \{(x - \xi)^2 + (y + \eta)^2\}^{1/2}$ and the contour in each integral is indented below the pole at $k = \alpha$ to ensure the outgoing behavior of G and H as $|x - \xi| \rightarrow \infty$. From (A1) and (A2) it can be shown that as $|x - \xi| \rightarrow \infty$

$$G(x, y; \xi, \eta) \sim 2\pi i \frac{\cosh \alpha(h - \eta) \cosh \alpha(h - y)}{h + \frac{1-s}{K} \sinh \alpha h} \frac{e^{i(\alpha^2 - \nu^2)^{1/2}|x - \xi|}}{(\alpha^2 - \nu^2)^{1/2}}, \tag{A3}$$

$$H(x, y; \xi, \eta) \sim -2\pi i \frac{\cosh \alpha(h - \eta) \sinh \alpha h e^{\alpha y}}{h + \frac{1-s}{K} \sinh \alpha h} \frac{e^{i(\alpha^2 - \nu^2)^{1/2}|x - \xi|}}{(\alpha^2 - \nu^2)^{1/2}}. \tag{A4}$$

(b) $\bar{G}(x, y; \xi, \eta)$ and $\bar{H}(x, y; \xi, \eta)$

$\bar{G}(x, y; \xi, \eta)$ and $\bar{H}(x, y; \xi, \eta)$ satisfy

$$(\nabla^2 - \nu^2)\bar{G} = 0 \text{ in } 0 \leq y \leq h,$$

$$(\nabla^2 - \nu^2)\bar{H} = 0 \text{ in } y \leq 0 \text{ except at } (\xi, \eta) (\eta < 0),$$

$$\bar{H} \sim K_0(\nu r) \text{ as } r \rightarrow 0,$$

$$\bar{G}_y = \bar{H}_y \text{ on } y = 0,$$

$$K\bar{G} + \bar{G}_y = s(K\bar{H} + \bar{H}_y) \text{ on } y = 0,$$

$$\bar{G}_y = 0 \text{ on } y = h, \quad \nabla \bar{H} \rightarrow 0 \text{ as } y \rightarrow -\infty,$$

\bar{G}, \bar{H} have outgoing nature as $|x - \xi| \rightarrow \infty$. Then $G(x, y; \xi, \eta)$ and $\bar{H}(x, y; \xi, \eta)$ are given by (cf [3]) after correcting the misprints

$$\bar{G}(x, y; \xi, \eta) = \frac{2s}{1+s} \left[K_0(\nu r) + \int_{\nu}^{\infty} \left\{ \frac{(1-s)k + \{s(K+k) - k\} e^{-kh} \operatorname{sech} kh}{\Delta(k)} \cosh k(h-y) + \frac{e^{-kh}}{\cosh kh} \sinh ky \right\} e^{k\eta} \frac{\cos\{(k^2 - \nu^2)^{1/2}(x - \xi)\}}{(k^2 - \nu^2)^{1/2}} dk \right], \tag{A5}$$

$$\begin{aligned} \bar{H}(x, y; \xi, \eta) = & K_0(\nu r) + \frac{1-s}{1+s} K_0(\nu r') + \frac{2s}{1+s} \int_{\nu}^{\infty} \left[e^{-kh} \operatorname{sech} kh \right. \\ & \left. - \frac{(1-s)k + s(K+k) - ke^{-kh} \operatorname{sech} kh}{\Delta(k)} \sinh kh \right] \\ & \times e^{k(y+\eta)} \frac{\cos\{(k^2 - \nu^2)^{1/2}(x - \xi)\}}{(k^2 - \nu^2)^{1/2}} dk \end{aligned} \tag{A6}$$

where again the contour in each integral is indented below the pole $k = \alpha$ to ensure the outgoing behavior of \bar{G} and \bar{H} as $|x - \xi| \rightarrow \infty$. From (A5) and (A6) it can be shown that as $|x - \xi| \rightarrow \infty$

$$\bar{G}(x, y; \xi, \eta) \sim -2s\pi i \frac{e^{\alpha\eta} \sinh \alpha h \cosh \alpha(h - y)}{h + \frac{1-s}{K} \sinh^2 \alpha} \frac{e^{i(\alpha^2 - \nu^2)^{1/2}|x - \xi|}}{(\alpha^2 - \nu^2)^{1/2}}, \tag{A7}$$

$$\bar{H}(x, y; \xi, \eta) \sim 2s\pi i \frac{e^{\alpha(\eta+y)} \sinh^2 \alpha h}{h + \frac{1-s}{K} \sinh^2 \alpha} \frac{e^{i(\alpha^2 - \nu^2)^{1/2}|x - \xi|}}{(\alpha^2 - \nu^2)^{1/2}}. \tag{A8}$$

REFERENCES

- [1] MILES, J W , Oblique surface wave diffraction by a cylindrical obstacle, *Dyn. Atmos. Oceans* **6** (1981), 121-123
- [2] MANDAL, B N and BASU, U , A note on oblique water wave diffraction by a cylindrical deformation of the bottom in the presence of surface tension, *Arch. Mech.* **42** (1990), 723-727
- [3] MANDAL, B N and CHAKRABARTI, R N , Two-dimensional, source potentials in a two-fluid medium for the modified Helmholtz's equation, *Internat. J. Math. & Math. Sci.* **9** (1986), 175-184
- [4] GORGUI, M A and KASSEM, S A , Basic singularities in the theory of internal waves, *Q. Jl. Mech. Appl. Math.* **31** (1978), 31-48.
- [5] RHODES-ROBINSON, P F , On waves at an interface between two liquids, *Math. Proc. Camb. Phil. Soc.* **38** (1980), 183-191



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

