

GENERATION OF PROPER CLASSES OF SHORT EXACT SEQUENCES

ALI PANCAR

Department of Mathematics
 Karadeniz Tech. University
 61080 - Trabzon / TURKEY

(Received September 18, 1995 and in revised form April 10, 1996)

ABSTRACT. The generation of proper classes of short exact sequences of modules by subclasses is considered. The class generated by two proper classes is studied by means of some operations between these classes. These operations are investigated in details for classes of short quasi-splitting, torsion-splitting and pure exact sequences of abelian groups.

KEY WORDS AND PHRASES: Proper class of short exact sequences, generated proper class, coprojectively (coinjectively) generated class, sum of proper classes, quasi-splitting (pure, torsion splitting) short exact sequences, cotorsion groups, Ulm's subgroup.

1991 AMS SUBJECT CLASSIFICATION CODES: 18G25, 20K40.

1. INTRODUCTION

By module we mean a unitary left R -module, R being an associative ring with identity and by group we mean a \mathbf{Z} -module, i.e. an abelian group.

Let \mathcal{R} be a class of short exact sequences of modules. If a short exact sequence

$$E: 0 \longrightarrow A \xrightarrow{\alpha} B \xrightarrow{\beta} C \longrightarrow 0 \tag{1.1}$$

belongs to \mathcal{R} , α is said to be an \mathcal{R} -monomorphism and β an \mathcal{R} -epimorphism. A short exact sequence E is determined by each of the monomorphism α and epimorphism β uniquely up to isomorphism. In the case of necessity, without loss of generality, we shall regard any monomorphism to be an inclusion map. ■ stands for the end of the proof. We use the terminologies and notations of Fuchs [1] and Mac Lane [2].

DEFINITION. The class \mathcal{R} is said to be *proper*, if it satisfies the following conditions (see Buchsbaum [3], Mac Lane [2], Sklyarenko [4]):

(P-1) Along with any short exact sequence \mathcal{R} contains every one isomorphic to it.

(P-2) \mathcal{R} contains all splitting short exact sequences.

(P-3) The composite of two \mathcal{R} -monomorphisms is an \mathcal{R} -monomorphism if this composite is defined.

(P-3') The composite of two \mathcal{R} -epimorphisms is an \mathcal{R} -epimorphism if it is defined.

(P-4) If α, β are monomorphisms and $\beta \circ \alpha$ is an \mathcal{R} -monomorphism, then α is an \mathcal{R} -monomorphism.

(P-4') If γ, δ are epimorphisms and $\delta \circ \gamma$ is an \mathcal{R} -epimorphism, then δ is an \mathcal{R} -epimorphism.

Short exact sequences of the form (1.1) from \mathcal{R} give the subgroup $\text{Ext}_{\mathcal{R}}(C, A)$ of the group of extensions $\text{Ext}_{\mathbf{R}}^1(C, A)$. Throughout this paper we shall write $\text{Ext}(C, A)$ instead of $\text{Ext}_{\mathbf{R}}^1(C, A)$.

DEFINITION. Let \mathcal{E} be a class of short exact sequences. The least proper class containing \mathcal{E} is said to be *generated* by \mathcal{E} and denoted by $\langle \mathcal{E} \rangle$.

Since the intersection of any family of proper classes is proper, we have evidently

$$\langle \mathcal{E} \rangle = \bigcap \{ \mathcal{R} : \mathcal{E} \subseteq \mathcal{R}; \mathcal{R} \text{ is a proper class} \}. \tag{1.2}$$

The notion of a generated proper class is a generalization of projectively (injectively), coprojectively (coinjectively) generated, and other, proper classes (see Stenström[5], Alizade [6]). In

this paper some properties of the class $\langle \mathcal{L} \rangle$ are given in terms of \mathcal{L} and the structure of this class is investigated by means of some operations between subclasses.

2. GENERATED PROPER CLASSES.

DEFINITION. A module A is called \mathcal{L} -coprojective (\mathcal{L} -coinjective) if every short exact sequence ending (beginning) at A belongs to \mathcal{L} .

The least proper class for which every module from the class of modules \mathcal{A} is coprojective (coinjective) is denoted by $\bar{k}(\mathcal{A})$ ($\underline{k}(\mathcal{A})$). Such classes are said to be *coprojectively (coinjectively) generated* (see Alizade [7][6]).

PROPOSITION 2.1. Let \mathcal{P} be a class of modules. For each $P \in \mathcal{P}$ take a short exact sequence

$$E^P: 0 \longrightarrow K_p \longrightarrow F_p \longrightarrow P \longrightarrow 0 \tag{2.1}$$

with some projective F_p . Denote $\{ E^P : P \in \mathcal{P} \}$ by \mathcal{E}_1^P . Then $\bar{k}(\mathcal{P}) = \langle \mathcal{E}_1^P \rangle$.

PROOF. Denote $\langle \mathcal{E}_1^P \rangle$ by \mathcal{L} . Since each module $P \in \mathcal{P}$ is $\bar{k}(\mathcal{P})$ -coprojective, every E^P from \mathcal{E}_1^P belongs to $\bar{k}(\mathcal{P})$. \mathcal{L} is the least proper class which contains all E^P , therefore $\mathcal{L} \subseteq \bar{k}(\mathcal{P})$.

On the other hand, since $E^P \in \mathcal{L}$, for any module A we can write the following exact sequences:

$$\begin{array}{ccccccc} \dots & \longrightarrow & \text{Hom}(F_p, A) & \longrightarrow & \text{Hom}(K_p, A) & \longrightarrow & \text{Ext}(P, A) \longrightarrow \text{Ext}(F_p, A) \longrightarrow \\ & & \parallel & & \parallel & & \cup & \cup & \tag{2.2} \\ \dots & \longrightarrow & \text{Hom}(F_p, A) & \longrightarrow & \text{Hom}(K_p, A) & \longrightarrow & \text{Ext}_{\mathcal{L}}(P, A) & \longrightarrow & \text{Ext}_{\mathcal{L}}(F_p, A) \longrightarrow \dots \end{array}$$

$\text{Ext}_{\mathcal{L}}(F_p, A) = \text{Ext}(F_p, A) = 0$, F_p being a projective module. By 5-Lemma $\text{Ext}_{\mathcal{L}}(P, A) = \text{Ext}(P, A)$. Then P is an \mathcal{L} -coprojective module. Since $\bar{k}(\mathcal{P})$ is the least proper class for which every $P \in \mathcal{P}$ is coprojective, we have $\bar{k}(\mathcal{P}) \subseteq \mathcal{L}$. ■

The dual statement can be proved for coinjectively generated classes.

DEFINITION. Let \mathcal{L} be a class of short exact sequences. A module A said to be \mathcal{L} -projective (\mathcal{L} -injective), if for every \mathcal{L} -epimorphism (\mathcal{L} -monomorphism) $\sigma : B \rightarrow C$

$$\sigma_* : \text{Hom}(A, B) \longrightarrow \text{Hom}(A, C) \quad (\sigma^* : \text{Hom}(C, A) \longrightarrow \text{Hom}(B, A)) \tag{2.3}$$

is an epimorphism. The class of all \mathcal{L} -projective (respectively, \mathcal{L} -injective) modules is denoted by $\pi(\mathcal{L})$ (resp. $\iota(\mathcal{L})$) (see [4], [5]).

A proper class \mathcal{L} is said to be *projective*, if for every module A there is an \mathcal{L} -epimorphism from an \mathcal{L} -projective module P onto A . An *injective* class is defined dually.

PROPOSITION 2.2. Let \mathcal{L} be a projective class and \mathcal{E} be the class of all short exact sequences

$$0 \longrightarrow A \longrightarrow P \longrightarrow C \longrightarrow 0 \tag{2.4}$$

from \mathcal{L} , P being an \mathcal{L} -projective module. Then $\mathcal{L} = \langle \mathcal{E} \rangle$.

PROOF. The inclusion $\langle \mathcal{E} \rangle \subseteq \mathcal{L}$ is obvious. On the other hand, let (1.1) be an arbitrary short exact sequence from \mathcal{L} . Since \mathcal{L} is a projective class, there exists an \mathcal{L} -epimorphism $\gamma : P \rightarrow B$ with an \mathcal{L} -projective P . By (P-3') $\beta \circ \gamma$ is an \mathcal{L} -epimorphism. At the same time $\beta \circ \gamma$ is an $\langle \mathcal{E} \rangle$ -epimorphism since P is an \mathcal{L} -projective module. Therefore, β is an $\langle \mathcal{E} \rangle$ -epimorphism by (P-4'). Thus $E \in \langle \mathcal{E} \rangle$. ■

The proposition dual to proposition 2.2 can be proved for an injective class \mathcal{L}

PROPOSITION 2.3. $\pi(\mathcal{L}) = \pi(\langle \mathcal{E} \rangle)$.

PROOF. $\mathcal{L} \subseteq \langle \mathcal{E} \rangle$ by the definition of $\langle \mathcal{E} \rangle$, therefore $\pi(\mathcal{L}) \supseteq \pi(\langle \mathcal{E} \rangle)$. On the other hand, the proper class $\pi^{-1}(\{K\})$ includes \mathcal{L} for every $K \in \pi(\mathcal{L})$. Since $\langle \mathcal{E} \rangle$ is the least proper class including \mathcal{L} ,

$\langle \mathcal{E} \rangle \subseteq \pi^{-1}(\{K\})$. Therefore, K is projective for every short exact sequence from $\langle \mathcal{E} \rangle$. Then $K \in \pi(\langle \mathcal{E} \rangle)$. Thus $\pi(\mathcal{E}) \subseteq \pi(\langle \mathcal{E} \rangle)$. ■

PROPOSITION 2.4. $u(\mathcal{E}) = u(\langle \mathcal{E} \rangle)$.

The proof is dual to that of Proposition 2.3

3. SOME OPERATIONS BETWEEN PROPER CLASSES.

DEFINITION. The proper class $\mathcal{R} + \mathcal{L} = \langle \mathcal{R} \cup \mathcal{L} \rangle$ is called *the sum* of the proper classes \mathcal{R} and \mathcal{L} .

Let us denote the class of short exact sequences whose monomorphisms are $\alpha \circ \beta$, α being an \mathcal{R} -monomorphism and β being an \mathcal{L} -monomorphism, by $\mathcal{R} \circ \mathcal{L}$.

Define the class $\mathcal{R} * \mathcal{L}$ of short exact sequences by the formula

$$\text{Ext}_{\mathcal{R} * \mathcal{L}}(C, A) = \text{Ext}_{\mathcal{R}}(C, A) + \text{Ext}_{\mathcal{L}}(C, A). \tag{3.1}$$

We have the following obvious facts:

- 1) $\mathcal{R} + \mathcal{L} = \mathcal{L} + \mathcal{R}$ is a proper class;
- 2) $\mathcal{R} * \mathcal{L} = \mathcal{L} * \mathcal{R}$;
- 3) $\mathcal{R} \circ \mathcal{L} \subseteq \mathcal{L} + \mathcal{R}$ and $\mathcal{R} * \mathcal{L} \subseteq \mathcal{R} + \mathcal{L}$.

In Theorem 3.1 we present the situation where $\mathcal{R} * \mathcal{L} \subseteq \mathcal{R} \circ \mathcal{L}$.

In general $\mathcal{R} \circ \mathcal{L}$ and $\mathcal{R} * \mathcal{L}$ need not be a proper class (see Corollary 4.1) But if $\mathcal{R} \circ \mathcal{L}$ (or $\mathcal{R} * \mathcal{L}$) is proper then, clearly, $\mathcal{R} + \mathcal{L} = \mathcal{R} \circ \mathcal{L}$ ($\mathcal{R} + \mathcal{L} = \mathcal{R} * \mathcal{L}$) and we can study $\mathcal{R} + \mathcal{L}$ by means of \mathcal{R} and \mathcal{L} (as in [6]).

The following proposition can be derived from Theorem 2.1 of Kepka [8], but we give another proof of this fact which relies on Theorem 1 of Alizade [7].

PROPOSITION 3.1. If the class of modules ρ is closed under submodules and extensions (see [7], [8]), then $\bar{k}(\rho) = \mathcal{E}^{\rho} \circ \mathcal{L}_0$, \mathcal{E}^{ρ} being the class of all short exact sequences ending at modules from ρ and \mathcal{L}_0 being the class of all splitting short exact sequences.

PROOF. Take an arbitrary short exact sequence (1.1) from $\bar{k}(\rho)$. Since ρ is closed under extensions, by Theorem 1 of Alizade [7] there exist $P \in \rho$, $E_1 \in \text{Ext}(P, A)$ and a homomorphism $\gamma: C \rightarrow P$ such that $E = \gamma^*(E_1)$. Since ρ is closed under submodules, $\text{Im } \gamma \in \rho$. Therefore γ can be taken as an epimorphism. From the cohomology sequence for $0 \rightarrow K \xrightarrow{\delta} C \xrightarrow{\gamma} P \rightarrow 0$, $K = \text{Ker } \gamma$

$$\dots \rightarrow \text{Ext}(P, A) \xrightarrow{\gamma^*} \text{Ext}(C, A) \xrightarrow{\delta^*} \text{Ext}(K, A) \rightarrow \dots \tag{3.2}$$

we have $E \in \text{Im } \gamma^* = \text{Ker } \delta^*$. Then in the following commutative diagram with exact rows and columns

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 \delta^*(E): & 0 & \longrightarrow & A & \longrightarrow & B' & \longrightarrow & K & \longrightarrow & 0 \\
 & & & & & \downarrow & & \downarrow \delta & & \\
 E: & 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & 0 \\
 & & & & & \downarrow & & \downarrow \gamma & & \\
 & & & & & P & & P & & \\
 & & & & & \downarrow & & \downarrow & & \\
 & & & & & 0 & & 0 & &
 \end{array} \tag{3.3}$$

$\delta^*(E) \in \mathcal{L}_0$. On the other hand, since $0 \rightarrow B' \rightarrow B \rightarrow P \rightarrow 0 \in \mathcal{E}^{\rho}$ we have $E \in \mathcal{E}^{\rho} \circ \mathcal{L}_0$

Conversely, since $\mathcal{E}^{\rho} \subseteq \langle \mathcal{E}^{\rho} \rangle$ and $\mathcal{L}_0 \subseteq \langle \mathcal{L}_0 \rangle$, we have $\mathcal{E}^{\rho} \circ \mathcal{L}_0 \subseteq \langle \mathcal{E}^{\rho} \rangle \subseteq \bar{k}(\rho)$. ■

LEMMA 3.1. Let (1.1) be a short exact sequence, $\gamma: D \rightarrow B$ be an arbitrary homomorphism and $F = \gamma^{-1}(A)$. Then the homomorphism $\delta: D/F \rightarrow C$, induced by γ , is a monomorphism.

PROOF. If $\delta(d+F) = 0$ then $\gamma(d) \in A$. Therefore $d \in \gamma^{-1}(A) = F$ and $d+F = F$. ■

LEMMA 3.2. If \mathcal{P} is a class of modules, closed under extensions and submodules (see [7], [8]), then for $\mathcal{L} = \overline{k}(\mathcal{P})$ and any proper class \mathcal{L} , $\text{Ext}_{\mathcal{L}}(F, A)$ is a subfunctor of $\text{Ext} : \text{R-mod} \times \text{R-mod} \rightarrow \text{Ab}$, R-mod being the category of left R-modules and Ab being the category of abelian groups.

PROOF. Let $E: 0 \rightarrow A \xrightarrow{\delta} D \rightarrow F \rightarrow 0 \in \text{Ext}_{\mathcal{L}}(F, A)$ and $\gamma: A \rightarrow M$ be an arbitrary homomorphism. Since $E \in \mathcal{L} \circ \mathcal{L}$, $\delta = \alpha' \circ \beta'$, α' being an \mathcal{L} -monomorphism and β' being an \mathcal{L} -monomorphism. By Prop.3.1 α' can be written as $\alpha' = \alpha \circ \theta$, where θ is a splitting monomorphism and $P = D/\text{Im} \alpha \in \mathcal{P}$. Therefore $\theta \circ \beta' = \beta$ is an \mathcal{L} -monomorphism. Thus $\delta = \alpha \circ \beta$ and we have the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 0 & \longrightarrow & A & \xrightarrow{\beta} & B & \longrightarrow & C \longrightarrow 0 \in \mathcal{L} \\
 & & \searrow & & \downarrow & & \downarrow \\
 & 0 & \longrightarrow & M & \xrightarrow{\beta_1} & B' & \longrightarrow C \longrightarrow 0 \in \mathcal{L} \\
 & & & \downarrow \alpha & & \downarrow & \\
 E: 0 & \longrightarrow & A & \xrightarrow{\delta} & D & \longrightarrow & F \longrightarrow 0 \\
 & & \searrow & & \downarrow \alpha_1 & & \downarrow \\
 \gamma_*(E): 0 & \longrightarrow & M & \xrightarrow{\delta_1} & D' & \longrightarrow & F \longrightarrow 0 \\
 & & & \downarrow P & & \downarrow P & \\
 & & & \downarrow & & \downarrow & \\
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 & & & 0 & & 0 &
 \end{array} \tag{3.4}$$

The short exact sequence $0 \rightarrow M \xrightarrow{\beta_1} B' \rightarrow C \rightarrow 0$ belongs to \mathcal{L} as an image of the short exact sequence $0 \rightarrow A \xrightarrow{\beta} B \rightarrow C \rightarrow 0$. Then β_1 is an \mathcal{L} -monomorphism.

The short exact sequence $0 \rightarrow B' \xrightarrow{\alpha_1} D' \rightarrow P \rightarrow 0$ belongs to \mathcal{L} as an image of $0 \rightarrow B \rightarrow D \rightarrow P \rightarrow 0$. Then α_1 is an \mathcal{L} -monomorphism. Therefore $\delta_1 = \alpha_1 \circ \beta_1$ is an $\mathcal{L} \circ \mathcal{L}$ -monomorphism and $\gamma_*(E): 0 \rightarrow M \xrightarrow{\delta_1} D' \rightarrow F \rightarrow 0 \in \mathcal{L} \circ \mathcal{L}$, i.e., $\text{Ext}_{\mathcal{L}}(F, A)$ is a subfunctor with respect to the second variable.

Now let us prove functoriness with respect to the first variable. Let $\nu: N \rightarrow F$ be an arbitrary homomorphism. Then taking $C' = \nu^{-1}(C)$ and $B'' = \mu^{-1}(B)$ we have the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 0 & \longrightarrow & A & \xrightarrow{\beta} & B & \longrightarrow & C \longrightarrow 0 \in \mathcal{L} \\
 & & & \downarrow & \swarrow & \downarrow & \swarrow \\
 & 0 & \longrightarrow & A & \longrightarrow & B'' & \longrightarrow C' \longrightarrow 0 \in \mathcal{L} \\
 & & & \downarrow & & \downarrow & \\
 E: 0 & \longrightarrow & A & \xrightarrow{\delta} & D & \longrightarrow & F \longrightarrow 0 \\
 & & & \downarrow & \swarrow & \downarrow & \swarrow \\
 \nu^*(E): 0 & \longrightarrow & A & \longrightarrow & D'' & \longrightarrow & N \longrightarrow 0 \\
 & & & \downarrow & \downarrow & \downarrow & \\
 & & & \downarrow & \downarrow & \downarrow & \\
 & & & P & & P & \\
 & & & \downarrow & \downarrow & \downarrow & \\
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 & & & 0 & & 0 &
 \end{array} \tag{3.5}$$

χ is a monomorphism by Lemma 3.1. Since ρ is closed under submodules, $P' \subseteq \text{Im} \chi \in \rho$. Therefore, by (P-1) we have $0 \rightarrow B'' \rightarrow D'' \rightarrow P' \rightarrow 0 \in \bar{k}(\rho) = \rho$. Since $\text{Ext}_L(C, A)$ is a contravariant functor with respect to the first variable, $0 \rightarrow A \rightarrow B'' \rightarrow C' \rightarrow 0 \in L$. Thus

$$v^*(E): 0 \rightarrow A \rightarrow D'' \rightarrow N \rightarrow 0 \in \rho \circ L. \tag{3.6}$$

So $\text{Ext}_{\rho \circ L}$ is a subfunctor of $\text{Ext}: R\text{-mod} \times R\text{-mod} \rightarrow \text{Set}$.

To prove that $\text{Ext}_{\rho \circ L}(C, A)$ is a subgroup of $\text{Ext}(C, A)$ for every C, A , let $E_1, E_2 \in \text{Ext}_{\rho \circ L}(C, A)$. Since $E_1 + E_2 = \nabla_{\Lambda}(E_1 \oplus E_2) \Delta_C$ (see Mac Lane [2]), to prove that $E_1 + E_2 \in \rho \circ L$, it is sufficient to show that $E_1 \oplus E_2 \in \rho \circ L$. Let $E_1: 0 \rightarrow A \xrightarrow{\alpha} B_1 \rightarrow C \rightarrow 0$ and $E_2: 0 \rightarrow A \xrightarrow{\beta} B_2 \rightarrow C \rightarrow 0$. Then $\alpha = \alpha_2 \circ \alpha_1, \beta = \beta_2 \circ \beta_1$, where α_1, β_1 are L -monomorphisms and α_2, β_2 are ρ -monomorphisms. Clearly, $\alpha_1 \oplus \beta_1$ is an L -monomorphism and $\alpha_2 \oplus \beta_2$ is an ρ -monomorphism. Therefore, $\alpha \oplus \beta = (\alpha_2 \oplus \beta_2) \circ (\alpha_1 \oplus \beta_1)$ is an $\rho \circ L$ -monomorphism. So $E_1 \oplus E_2 \in \rho \circ L$.

One can easily show that $E \in \rho \circ L$ implies $-E \in \rho \circ L$. Thus $\text{Ext}_{\rho \circ L}$ is a subfunctor of $\text{Ext}: R\text{-mod} \times R\text{-mod} \rightarrow \text{Ab}$. ■

REMARK. Now to prove that $\rho \circ L$ is a proper class it remains only to show that the composite of two $\rho \circ L$ -monomorphisms is an $\rho \circ L$ -monomorphism (see Nunke [9]). By means of this fact we have obtained the description of the smallest proper class with given classes of coprojective and coinjective modules as a "composite" of coinjectively and coprojectively generated classes, which will soon be published.

Since $\rho \subseteq \rho \circ L$ and $L \subseteq \rho \circ L$, Lemma 3.2 gives the following inclusion.

THEOREM 3.1. If the class ρ of modules is closed under extensions and submodules (see [7], [8]), then $\rho * L \subseteq \rho \circ L$ for $\rho = \bar{k}(\rho)$ and for every proper class L . ■

4. SOME EXAMPLES IN THE CATEGORY OF ABELIAN GROUPS.

Let δ_0 be the class of all splitting short exact sequences, δ be the class of all pure short exact sequences and ρ be the class of all torsion splitting short exact sequences (see Fuchs [1]). For a proper class ρ put

$$\hat{\rho} = \{E : \text{for some } n \in \mathbb{Z} (n \neq 0) \ nE \in \rho\}, \tag{4.1}$$

(see Alizade [6]).

Class $\hat{\delta}_0$ of all short quasi-splitting exact sequences was introduced and studied by Walker C.P. [10]. Classes $\hat{\delta}$ and $\hat{\rho}$ were studied by Hart in [11], where these classes were denoted by \hat{A} and \hat{C} .

Clearly, $\rho + \hat{\delta}_0 \subseteq \hat{\rho}$ for every proper class ρ .

THEOREM 4.1. $\rho + \hat{\delta}_0 = \hat{\rho}$.

PROOF. Denote $\rho + \hat{\delta}_0$ by \mathcal{J} . Every $E \in \rho$ can be written as $E = 1 \cdot E$ ($n=1$), therefore $E \in \hat{\rho}$ and then $\rho \subseteq \hat{\rho}$. On the other hand, $\hat{\delta}_0 \subseteq \rho$, therefore $\hat{\delta}_0 \subseteq \hat{\rho}$. Thus $\rho + \hat{\delta}_0 \subseteq \hat{\rho}$.

Conversely, let $E: 0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0 \in \hat{\rho}$. Then $nE \in \rho$ for some $n \neq 0$. Let us denote by n the endomorphism of multiplication by n on A . Then by Lemma 52.1 in Fuchs [1] nE is the lower exact sequence in the following commutative diagram:

$$\begin{array}{ccccccc} E: & 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & 0 & \in & \hat{\rho} \\ & & & \downarrow n & & \downarrow & & \parallel & & & & \\ nE: & 0 & \longrightarrow & A & \longrightarrow & B' & \longrightarrow & C & \longrightarrow & 0 & \in & \rho. \end{array} \tag{4.2}$$

The endomorphism $n:A \rightarrow A$ can be represented in the natural way $n = \alpha \circ \sigma$, where $\sigma:A \rightarrow nA$ is epimorphism and $\alpha:nA \rightarrow A$ is inclusion map

We have the following commutative diagrams with exact rows and columns:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & A[n] & & A[n] & & \\
 & & \downarrow & & \downarrow & & \\
 E: & 0 \longrightarrow & A & \longrightarrow & B & \xrightarrow{\theta} & C \longrightarrow 0 \in \hat{\mathcal{R}} \\
 & & \downarrow \sigma & & \downarrow \gamma & & \\
 E': & 0 \longrightarrow & nA & \xrightarrow{\nu} & B_1 & \xrightarrow{\delta} & X \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & \\
 E'': & 0 \longrightarrow & nA & \xrightarrow{\nu} & B_1 & \xrightarrow{\delta} & X \longrightarrow 0 \\
 & & \downarrow \alpha & & \downarrow \beta & & \\
 nE: & 0 \longrightarrow & A & \xrightarrow{\mu} & B' & \longrightarrow & C \longrightarrow 0 \in \mathcal{R} \\
 & & \downarrow & & \downarrow & & \\
 & & A/nA & & A/nA & & \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array} \tag{4.3}$$

$$\begin{array}{ccccccc}
 & & \downarrow \alpha & & \downarrow \beta & & \\
 nE: & 0 \longrightarrow & A & \xrightarrow{\mu} & B' & \longrightarrow & C \longrightarrow 0 \in \mathcal{R} \\
 & & \downarrow & & \downarrow & & \\
 & & A/nA & & A/nA & & \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array} \tag{4.4}$$

A/nA is a bounded and hence $\hat{\delta}_0$ -coprojective group. Therefore the short exact sequence $0 \rightarrow nA \xrightarrow{\alpha} A \rightarrow A/nA \rightarrow 0$ belongs to $\hat{\delta}_0$ and since $\hat{\delta}_0 \subseteq \mathcal{I}$, it belongs to \mathcal{I} , i.e., α is a \mathcal{I} -mono-morphism. Since $nE \in \mathcal{R} \subseteq \mathcal{I}$, μ is a \mathcal{I} -monomorphism. Therefore, by (P-3), $\mu \circ \alpha = \beta \circ \nu$ is a \mathcal{I} -monomorphism. Hence, by (P-4), ν is a \mathcal{I} -monomorphism. Thus $E' \in \mathcal{I}$.

$A[n]$ is a bounded and therefore $\hat{\delta}_0$ -coinjective group. Hence the short exact sequence $0 \rightarrow A[n] \rightarrow B \rightarrow B_1 \rightarrow 0$ belongs to $\hat{\delta}_0$ and since $\hat{\delta}_0 \subseteq \mathcal{I}$, it belongs to \mathcal{I} , i.e., γ is a \mathcal{I} -epimorphism. Since $E' \in \mathcal{I}$, δ is a \mathcal{I} -epimorphism. By (P-3') $\theta = \delta \circ \gamma$ is a \mathcal{I} -epimorphism. Therefore $E \in \mathcal{I}$ and we have $\hat{\mathcal{R}} \subseteq \mathcal{I}$. ■

DEFINITION. For a subgroup A of B let $\hat{A} = \{ b \in B : mb \in A \text{ for some } m \in \mathbb{Z}, (m \neq 0) \}$

LEMMA 4.1. Let B_d be the maximal divisible subgroup and $T(B)$ be the torsion part of a group B . Then

$$\hat{B}_d = B_d + T(B) = B_d + \hat{0} \tag{4.5}$$

PROOF. The inclusion $B_d + T(B) \subseteq \hat{B}_d$ is obvious.

Let $B = B_d \oplus K$, $K \leq B$ and let $b = a + k$ be any element from \hat{B}_d ($a \in B_d, k \in K$). Then $nb \in B_d$ for some $n \neq 0$. Since B_d is divisible, $nb = na_1$ for some $a_1 \in B_d$. Therefore $na + nk = nb = na_1$. By directness of the sum $B_d \oplus K$, $nk = 0$ and hence $k \in T(B)$. So $b = a + k \in B_d + T(B)$. ■

Now we take $B = \text{Ext}(C, A)$ for any groups C, A , then $B_d = \text{Ext}_d(C, A)$ by Prop. 58.3 in Fuchs [1] and $T(B) = \text{Ext}_{\hat{\delta}_0}(C, A)$ (see [10]). Therefore by Lemma 4.1 we have

$$\text{Ext}_{\hat{\beta}}(C,A) = \text{Ext}_{\beta}(C,A) + \text{Ext}_{\hat{\beta}_0}(C,A) \tag{4.6}$$

for each group C,A, i.e., $\hat{\beta} = \beta * \hat{\beta}_0$. This equality together with Theorem 3.1 (note that $\beta = \bar{k}(\rho)$, ρ being the class of all torsion free groups (see [7]) gives us the following theorem.

THEOREM 4.2. $\beta * \hat{\beta}_0 = \beta \circ \hat{\beta}_0 = \beta + \hat{\beta}_0 = \hat{\beta}$. So every $\hat{\beta}$ -monomorphism is a composition of an $\hat{\beta}_0$ -monomorphism and a β -monomorphism. ■

For the class β the situation is worse.

LEMMA 4.2. There exists a reduced cotorsion group C, Ulm's subgroup of which is torsion free and algebraically compact group

PROOF Let J_p be the group of p-adic numbers. The p-basic subgroup $B = \langle b \rangle$ of J_p is isomorphic to the group Z of integers and J_p/B is a divisible group (see ch.20 and 32 in Fuchs [1]). Let us construct an extension K of B such that the first Ulm's subgroup K^1 of K is equal to B. Define K by means of generators

$$b_1, b_2, \dots, b_n, \dots : K = \langle b_1, b_2, \dots, b_n, \dots \rangle \tag{4.7}$$

and relations

$$b_1 = b, 2b_2 = b, \dots, nb_n = b, \dots \tag{4.8}$$

K/B is isomorphic to a direct sum of cyclic groups, hence $(K/B)^1 = 0$ and we have $K^1 \subseteq B$. On the other hand, the inclusion $B \subseteq K^1$ is obvious and thus $K^1 = B$.

Let us complete the diagram

$$\begin{array}{ccc} B & \xrightarrow{\alpha} & J_p \\ \downarrow & & \\ K & & \end{array} \tag{4.9}$$

to a pushout diagram. Then we have the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & B & \xrightarrow{\alpha} & J_p & \longrightarrow & J_p/B \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & K & \xrightarrow{\alpha'} & D & \longrightarrow & D/K \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \\ & & K/B & & D/J_p & & \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array} \tag{4.10}$$

Since α is a monomorphism, α' is also a monomorphism and $J_p/B \cong D/K$, $K/B \cong D/J_p$ (see Th 10.2 [1]).

Now let us prove that $D^1 = J_p$. Since $(D/J_p)^1 \cong (K/B)^1 = 0$ we have $D^1 \subseteq J_p$. On the other hand let $x \in J_p$ and $n \in \mathbb{Z}$ ($n \neq 0$) be arbitrary. Since J_p/B is divisible, there is $x_1 \in J_p$ such that $x - nx_1 \in B = K^1$,

therefore there is an element $k \in K$ such that $x - nx_1 = nk$. Then $x = n(x_1 + k) \in nD$. Thus we have $x \in D^1$ and $D^1 = J_p$.

Since D/J_p is isomorphic to a direct sum of finite cyclic groups, it can be embedded into a direct product of these cyclic groups, i.e., into a reduced algebraically compact group $M: D/J_p \xrightarrow{\beta} M$. By Theorem 51.3 [1]

$$\beta^*: \text{Ext}(M, J_p) \longrightarrow \text{Ext}(D/J_p, J_p) \tag{4.11}$$

is an epimorphism. Therefore there is an extension C of D such that the following diagram with exact rows and columns is commutative:

$$\begin{array}{ccccccc} 0 & \longrightarrow & J_p & \longrightarrow & D & \longrightarrow & D/J_p \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \beta \\ 0 & \longrightarrow & J_p & \longrightarrow & C & \longrightarrow & M \longrightarrow 0 \end{array} \tag{4.12}$$

$J_p = D^1 \subseteq C^1$. Since M is a reduced algebraically compact group, we have $(C/J_p)^1 \cong M^1 = 0$ by Prop.54.2 [1]. Then $C^1 \subseteq J_p$. Hence we have $C^1 = J_p$ and C is a reduced group. By Prop. 39.4 [1] J_p is an algebraically compact group. Since J_p and M are algebraically compact and hence cotorsion groups, by ch 54(D) [1] C is a reduced cotorsion group. ■

LEMMA 4.3. If the Ulm's subgroup $A = C^1$ of a reduced group C is torsion free and different from zero, then

$$A + T(C) \neq \hat{A}, \tag{4.13}$$

$T(C)$ being the torsion part of C .

PROOF. Take an element $a \notin nA$ of the reduced group A . Since $A = \bigcap_{0 \neq n \in \mathbb{Z}} nC$, there is $b \in C$ such that $nb = a$. Obviously $b \in \hat{A}$. If $b \in A + T(C)$, one can write $b = a_1 + t$ with $a_1 \in A$ and $t \in T(C)$. Then $a = nb = na_1 + nt$ and hence $nt = a - na_1 \in A \cap T(C)$. Since A is torsion free, $nt = 0$ and we have $a = na_1 \in nA$. But this contradicts the assumption. Thus $b \in \hat{A} \setminus (A + T(C))$. ■

The following theorem demonstrates that for proper classes \mathcal{R} and \mathcal{L} the class $\mathcal{R} * \mathcal{L}$ need not be proper.

THEOREM 4.3. $\mathcal{R} * \hat{\mathcal{L}}_0 \neq \hat{\mathcal{R} * \mathcal{L}}$.

PROOF. According to Lemma 4.2 there is a reduced cotorsion group C , Ulm's subgroup $A = C^1$ of which is reduced, torsion free and algebraically compact. By ch. 54(H) [1] we have

$$\text{Ext}(\mathbb{Q}/\mathbb{Z}, C) \cong C \tag{4.14}$$

By Theorem 53.3 [1]

$$\text{Ext}_s(\mathbb{Q}/\mathbb{Z}, C) = \text{Pext}(\mathbb{Q}/\mathbb{Z}, C) \cong C^1 = A. \tag{4.15}$$

On the other hand,

$$\text{Ext}_t(\mathbb{Q}/\mathbb{Z}, C) = \text{Text}(\mathbb{Q}/\mathbb{Z}, C) \cong T(C) \tag{4.16}$$

and

$$\text{Ext}_s(\mathbb{Q}/\mathbb{Z}, C) \cong \hat{A}. \tag{4.17}$$

By Lemma 4.3 we have

$$\text{Ext}_\delta(\mathbf{Q}/\mathbf{Z}, \mathbf{C}) + \text{Ext}_{\hat{\delta}}(\mathbf{Q}/\mathbf{Z}, \mathbf{C}) \neq \text{Ext}_{\hat{\delta}}(\mathbf{Q}/\mathbf{Z}, \mathbf{C}) \quad (4.18)$$

Thus $\delta * \hat{\delta}_0 \neq \hat{\delta}$. ■

COROLLARY 4.1. The class $\delta * \hat{\delta}_0$ is not proper

PROOF. Since $\text{Ext}_\delta(\mathbf{C}, \mathbf{A})$ and $\text{Ext}_{\hat{\delta}}(\mathbf{C}, \mathbf{A})$ are subgroups of $\text{Ext}_{\hat{\delta}}(\mathbf{C}, \mathbf{A})$ for any group \mathbf{C} and \mathbf{A} , evidently $\delta * \hat{\delta}_0 \subseteq \hat{\delta}$. Since $\hat{\delta} = \delta + \hat{\delta}_0$ is the least proper class including δ and $\hat{\delta}_0$, if $\delta * \hat{\delta}_0$ were a proper class we should have $\hat{\delta} \subseteq \delta * \hat{\delta}_0$ and hence $\delta * \hat{\delta}_0 = \hat{\delta}$. But this contradicts Theorem 4.3. ■

ACKNOWLEDGEMENT. The author wishes to express his gratitude to his referee for helpful criticism.

REFERENCES

- [1] FUCHS, L., "Infinite abelian groups," v.1, *Academic Press*, 1970
- [2] MAC LANE, S., "Homology," *Springer-Verlag*, 1975.
- [3] BUCHSBAUM, D., "A note on homology in categories" *Ann. of Math.* 69:1(1959), 66-74.
- [4] SKLYARENKO, E.G., "Relative homological algebra in the category of modules," *Usp. Mat. Nauk (Russian Math. Surveys)*, 33:3(1978), 85-120.
- [5] STENSTRÖM, B., "Pure submodules," *Arkiv Math.* 7 12(1967), 159-171.
- [6] ALIZADE, R.H., "On the proper classes of short exact sequences in the category of abelian groups," *Math. Notes* 40:1(1986), 3-15.
- [7] ALIZADE, R.H., "On the proper classes of Kepka," *Math. Notes* 37:2(1985), 268-273
- [8] KEPKA, T, "On one class of purities," *Comment. Math. Univ. Carol.* 14:1(1973), 139-154.
- [9] NUNKE, R.J., "Purity and subfunctors of the identity," - In: *Topics in abelian groups, Chicago*, 1963, 121-171.
- [10] WALKER, C.P. "Properties of Ext and quasi-splitting of abelian groups," *Acta Math. Acad. Sci. Hung.*, 15(1964), 157-160.
- [11] HART, N., "Two parallel homological algebras," *Acta Math. Acad. Sci. Hung.*, 25:3-4(1974), 321-327



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

