# A CLOSER LOOK AT SOME NEW IDENTITIES

**GEOFFREY B. CAMPBELL** 

Centre for Mathematics and its Applications School of Mathematical Sciences Australian National University Canberra, ACT 0200, Australia

(Received February 23, 1996)

## ABSTRACT.

We obtain infinite products related to the concept of visible from the origin point vectors. Among these is

$$\prod_{k=3}^{\prime} \left(1 - z^{k}\right)^{\varphi_{k}(k)/k} = \frac{1}{1 - z} \exp\left(\frac{z^{3}}{2(1 - z)^{2}} - \frac{1}{2}z - \frac{1}{2z(1 - z)}\right), \qquad |z| < 1,$$

in which  $\varphi_{3}(k)$  denotes for fixed k, the number of positive integer solutions of (a, b, k) = 1 where a < b < k, assuming (a, b, k) is the gcd(a, b, k)

**KEY WORDS AND PHRASES.** Combinatorial identities, Combinatorial number theory, Lattice points in specified regions, Partitions (elementary number theory)

1992 AMS SUBJECT CLASSIFICATION CODES. 05A19, 11B75, 11P21, 11P81

## 1. INTRODUCTION.

In several of the author's recent papers (see Campbell [5-9]) a class of new elementary infinite product identities was introduced. These were given the name *vpv identities*, meaning *visible (from the origin) point vector identities*, due to the fact that they involved summation over so called visible lattice points in various dimensional spaces For an introduction to the idea of visible lattice points see Apostol [2], where their distribution is calculated. The identities which occurred most frequently in Campbell [7] were often related to each other by vpv lattice sums dividing up space into radial regions from the origin. These identities when grouped were called *companion identities*, because of their interdependence. We used the operators defined by

G. B. CAMPBELL

$$\sum_{k} \sum_{j} = \sum_{k=2}^{\infty} \sum_{j \in \phi_{k}} , \qquad \qquad \prod_{k} \prod_{j} = \prod_{k=2}^{\infty} \prod_{j \in \phi_{k}} , \qquad (1 \ 1)$$

where  $\phi_k$  is the set of positive integers less than and relatively prime to k Whilst these operators are useful for the 2-D sums, as shown in the author's papers [5-6, 9], we will not require them for the higher dimensional sums considered here

Recently Mosseri [13], and Baake et al [3], have considered the so-called Optical Fourier Transform of the 2-D visible lattice points as an optical experiment The subject of lattice point methods applied to physics and chemistry is also examined in Frankel et al [10], and Glasser and Zucker [11] Some of the methods of these papers are applicable to vpv sums and products In Campbell [5-9] the following companion identities were given.

$$(1-y) \sum_{k} \sum_{j} (1-x^{j} y^{k})^{1/k} = \left(\frac{1-y}{1-xy}\right)^{1/(1-x)}, \qquad (x \neq 1)$$
(12)

$$(1 - xy) \sum_{k} \sum_{j} (1 - x^{k} y^{j})^{1/j} = (1 - xy)^{1/(1 - x)}, \qquad (1 3)$$

$$(1-y)(1-xy) \prod_{k} \prod_{j} (1-x^{j} y^{k})^{1/k} (1-x^{k} y^{j})^{1/j} = (1-y)^{1/(1-x)},$$
(14)

valid respectively for |xy| and |y| < 1, |x| and |xy| < 1, |x| and |y| < 1.

These led to ideas which link to several fields of research For example, new products over the primes were presented in Campbell [6-7,9], new Jacobi theta function identities in [6] Paper [6] also contained results on Dirichlet summations such as those connected with Ramanujan arithmetical functions, and was related to a method of Meinardus for obtaining asymptotic estimates of coefficients from infinite products (see Andrews [1], Hardy and Wright [11]) In [7], some further identities identities of this type were given and the existence of yet others was indicated In the present note we give results derived from considerations similar to those in [7] The underlying concept in the author's papers [5-9] is expressed in

LEMMA 1.1 Consider an infinite region radiating out of the origin in any Euclidean vector space The set of all lattice point vectors from the origin in that region is precisely the set of positive integer multiples of the visible point vectors (vpv's) in that region.

Immediate a priori consequences of this may be written. For example,

$$\sum_{\substack{a \ b \ge 1\\ z \ b \ge 1}} \frac{y^{a} z^{b}}{1 - y^{a} z^{b}} = \frac{z}{(1 - y)(1 - z)},$$
(1.5)

$$\sum_{\substack{a \ b = 1 \\ r \ge 0 \ b \ge 1 \\ a \ b}} \frac{y^a z^b}{1 - y^a z^b} = \sum_{k=1}^{\alpha} \frac{1 - y^k}{1 - y} z^k = \frac{z}{(1 - y)(1 - z)} - \frac{yz}{(1 - y)(1 - yz)},$$
(1.6)

valid for |y| and |z| < 1, |y| and |yz| < 1, respectively

It is straightforward to generalize these results to any number of variables For example,

$$\sum_{\substack{(a \ b \ c \ d) = 1 \\ a \ b \ c \ d) = 1}} \frac{w^a x^b y^c z^d}{1 - w^a x^b y^c z^d} = \frac{z}{(1 - w)(1 - x)(1 - y)(1 - z)},$$
(17)

582

$$\sum_{\substack{(a,b,c,d)=1\\a,b,c,d}\\a,b,c,d} \frac{w^a x^b y^c z^d}{1-w^a x^b y^c z^d} = \sum_{k=1}^{\alpha} \frac{(1-w^k)(1-x^k)(1-y^k)}{(1-w)(1-x)(1-y)} z^k , \qquad (18)$$

with (1 7) valid for |w|, |x|, |y| and |z| < 1 whilst (1 8) is true if absolute values of z, wz, xz, yz, wxz, xyz, wyz, wyz, wzyz are all less than unity

Another consequence of the above considerations is

THEOREM 1 2. Under the same conditions as (1 5) and (1 6) respectively,

$$\prod_{\substack{a,b,c\}=1\\a\,b\leq 0,c\geq 1}} (1-x^a y^b z^c)^{1/c} = (1-z)^{1/(1-x)(1-y)},$$
(19)

$$\prod_{\substack{(a,b,x)=1\\a,b\in(x)=1\\a,b\in(x)}} (1-x^a y^b z^c)^{1/\epsilon} = \left(\frac{(1-xz)(1-yz)}{(1-z)(1-xyz)}\right)^{1/(1-x)(1-y)}$$
(1.10)

This is true since logarithmic derivatives of both sides of  $(1 \ 9)$  and (1.10) yield  $(1 \ 5)$  and  $(1 \ 6)$  respectively. This establishes the theorem to within a constant factor in each product, and this constant may be determined from allowing both x and y to approach unity.

## 2. VPV IDENTITIES AND BOUNDARY LIMIT CASES.

The identities of section 1 have interesting cases in which variables other than z approach unity (often the boundary of convergence) given certain modifications In Campbell [5] we find the identity

$$\prod_{k=1}^{\infty} \left( 1 - y^k \right)^{\varphi(k)/k} = \exp\left(\frac{-y}{1 - y}\right),$$
(2 1)

where  $\varphi$  is the Euler totient function. An equivalent result was stated in Borwein and Borwein [4],

$$\prod_{k=1}^{\infty} (1 - y^k)^{\mu(k)/k} = e^{y}, \qquad (2 2)$$

where  $\mu$  is the Mobius function. We now establish

THEOREM 2.1

$$\prod_{k=3}^{m} \left(1 - z^{k}\right)^{\varphi_{1}(k)/k} = \frac{1}{1 - z} \exp\left(\frac{z^{3}}{2(1 - z)^{2}} - \frac{1}{2}z - \frac{1}{2z(1 - z)}\right), \qquad |z| < 1,$$
(2.3)

where  $\varphi_3(k)$  denotes for fixed k the number of positive integer solutions of (a, b, k) = 1 where a < b < k

The function  $\varphi_{3}(k)$  is a natural generalization of the Euler totient function  $\varphi$  which is the number of positive integers less than and relatively prime to k

In the recent paper by the author [7], it was shown that for absolute values of z, yz, xyz, all less than unity,

$$\prod_{\substack{(a,b,t)=1\\a,b,t-1\\a,t,b-1,t-2}} (1-x^{a}y^{b}z^{c})^{-1/t}$$

$$= \frac{(1-yz)^{(1-x)(1-y)}(1-xyz)^{(1-x)(1-y)}}{(1-z)^{(1-x)(1-y)}} \exp\left(\frac{-xyz(2xy-\frac{1}{2}yz-\frac{1}{2}xyz-z)}{(1-x)(1-y)(1-xy)}\right)$$
(2.4)

If we arrange suitably for  $x \to 1$ ,  $y \to 1$ , in this we should expect to obtain (2.3) There is no difficulty with the left side, but the right side seems to present formal (if not conceptual) barriers A way around this is to construct (2.3) by a method offering a geometrical viewpoint For suitable conditions the logarithm of the left side of (2.3) equals

$$\sum_{k=3}^{\infty} \frac{\varphi_{3}(k)}{k} \log(1-z^{k})$$

$$= \sum_{k=1}^{7} \sum_{\substack{a \in b \\ a \neq b \\ a \neq b \neq 1}} 1^{ak} 1^{bk} z^{kr} (kc)^{-1}$$

$$= \sum_{\substack{a \in b \\ a \neq b \neq 1}} 1^{a} 1^{b} z^{r} c^{-1}$$

$$= \frac{z^{3}}{3} (1^{2}1^{1}) + \frac{z^{4}}{4} (\frac{1^{3}1^{2} + 1^{3}1^{1}}{+1^{2}1^{1}}) + \frac{z^{5}}{5} (\frac{1^{4}1^{1} + 1^{4}1^{2} + 1^{4}1^{1}}{+1^{2}1^{1}}) + \dots$$

$$= \frac{z^{3}}{3} \cdot 1 + \frac{z^{4}}{4} \cdot 3 + \frac{z^{5}}{5} \cdot 6 + \frac{z^{6}}{6} \cdot 10 + \frac{z^{7}}{7} \cdot 15 + \dots$$

$$= \log(\frac{1}{1-z}) - \frac{1}{2z(1-z)} - \frac{1}{2}z + \frac{z^{3}}{2(1-z)^{2}}.$$

From here, exponentiation gives the right side of (2 3) which proves the theorem

Further examples of the vpv identities reduce to new results An infinite product example given in Campbell [7] is for |x| and |y| < 1,

$$\prod_{\substack{(a,b),\ 1\\a,b'\in Z^*}} \left(\frac{1}{1-x^a y^b}\right)^{\frac{1}{a^m b^*}} = \exp\left\{ \left(\sum_{j=1}^n \frac{x^j}{j^m}\right) \left(\sum_{k=1}^n \frac{y^k}{k^n}\right) \right\} \quad \text{where} \quad m+n=1.$$
(2.5)

Suppose now we select x = y, and  $m = n = \frac{1}{2}$  Then we have the result valid for |x| < 1

$$(1 - x^{1+1})^{-\sqrt{(1)(1)}} (1 - x^{1+2})^{-2\sqrt{(1)(2)}} (1 - x^{1+3})^{-2\sqrt{(1)(3)}} (1 - x^{2+3})^{-2\sqrt{(2)(3)}} (1 - x^{1+4})^{-2\sqrt{(1)(4)}} (1 - x^{3+4})^{-2\sqrt{(3)(1)}} \cdots$$

$$= \exp\left\{\left(\frac{x}{\sqrt{1}} + \frac{x^{2}}{\sqrt{2}} + \frac{x^{3}}{\sqrt{3}} + \dots\right)^{2}\right\}$$

$$= (1 - x^{2})^{a_{1}} (1 - x^{3})^{a_{1}} (1 - x^{1})^{a_{4}} (1 - x^{5})^{a_{5}} \cdots$$
(2.6)

where  $a_2 = -\sqrt{(1)(1)}$ ,  $a_3 = -2\sqrt{(1)(2)}$ ,  $a_4 = -2\sqrt{(1)(3)}$ ,  $a_5 = -2(\sqrt{(1)(4)} + \sqrt{(2)(3)})$ , ... in which the factors in the products under each square root are coprime and sum to the suffix i of  $a_i$ 

## 3. A DIRICHLET SERIES VERSION OF A GENERAL VPV THEOREM.

In the author's paper [7] we find the fundamental identity in vpv theory, THEOREM 3.1 If i = 1, 2, 3, ..., n then for each  $|x_i| < 1$ , and  $b_i \in C$ , with  $\sum_{i=1}^{n} b_i = 1$ ,

$$\prod_{\substack{(a_1,a_2,\ldots,a_r)=1\\a_i,Z^*}} \left(\frac{1}{1-x_1^{a_1}x_2^{a_2}\cdots x_n^{a_r}}\right)^{a_1^{b_1}a_2^{b_2}\cdots a_n^{b_r}} = \exp\left\{\prod_{i=1}^n \left(\sum_{k=1}^r \frac{x_i^j}{j^{b_i}}\right)\right\}.$$
 (3.1)

This result came from lemma 1 1 much in the manner shown above for theorem 1.2 A similar line of reasoning applies also to Dirichlet summations over the vpv lattices We now state THEOREM 3.2 If i = 1, 2, ..., n then for each  $b_i \in C$ ,  $\Re b_i > 1$ ,

$$\sum_{\substack{(a_1,a_2,\ldots,a_n)=1\\a_1\in Z^*}}\frac{1}{a_1^{b_1}a_2^{b_2}\cdots a_n^{b_n}} = \frac{\zeta(b_1)\zeta(b_2)\cdots \zeta(b_n)}{\zeta(b_1+b_2+\cdots+b_n)}.$$
(3.2)

This result appears also to be fundamental in some sense, but is not seen in any of the literature, except for the case with n = 2 given recently in Campbell [6] The latter case is so near the surface it is probably known. The proof of (3.2) follows trivially from applying lemma 1.1 to the left side series after multiplying both sides by

$$\sum_{a:\mathbb{Z}^*} \frac{1}{a^{(b_1+b_2+\cdots+b_n)}} = \zeta(b_1+b_2+\cdots+b_n).$$
(3.3)

It may of course be argued that (3.2) is simply a limiting case of (3.1) Indeed generalization of the Euler totient function much in the style of our  $\varphi_3(k)$  will give the generating Dirichlet series via (3.2)

ACKNOWLEDGEMENTS. The author thanks the referee for providing important comments, Australian National University for continued association, and Professor George E Andrews for continued moral support

#### **REFERENCES**

- 1 ANDREWS, G E <u>The Theory of Partitions</u>, Encyclopedia of Mathematics, Vol 2, Addison Wesley, 1976
- 2 APOSTOL, T Introduction to Analytic Number Theory, Springer-Verlag, New York, 1976
- 3 BAAKE, M, GRIMM, U, and WARRINGTON, D H Some remarks on the visible points of a lattice, J Phys A Vol 27, No 8, 1994, 2669-2674
- 4 BORWEIN, J., and BORWEIN, P Pi and the AGM., John Wiley and Sons, New York, 1986
- 5 CAMPBELL, G. B A new class of infinite product, and Euler's totient, <u>Int J Math & Math</u> <u>Sci.</u>, Vol 17, No 4, 1994, 417-422
- 6 CAMPBELL, G B Dirichlet summations and products over primes, <u>Int J Math & Math Sci.</u>, Vol 16, No 2, 1993, 359-372
- 7 CAMPBELL, G. B. Infinite products over visible lattice points, <u>Int J Math & Math Sci</u>, Vol 17, No 4, 1994, 637-654.
- 8 CAMPBELL, G B Infinite products over hyperpyramid lattices, <u>Int J Math & Math Sci</u>, (to appear)
- 9 CAMPBELL, G B Multiplicative functions over Riemann zeta function products, <u>J Ramanujan</u> Soc 7 No 1, 1992, 52-63
- 10 FRANKEL, N E, GLASSER, M L., HUGHES, B D, and NINHAM, B W Mobius, Mellin, and Mathematical Physics, <u>Physica A, 186</u>, 1992, 441-481.

- 11 GLASSER, M L, and ZUCKER, I J Lattice Sums, <u>Th Chem\_Adv Persp</u>, Vol 5, 1980, 67-139
- 12 HARDY, G H, and WRIGHT, E M <u>An Introduction to the Theory of Numbers</u>, Oxford University Press, Clarendon, London, 1971
- 13 MOSSERI, R Visible points in a lattice, <u>J Phys A25</u>, 1992, L25-29



Advances in **Operations Research** 



**The Scientific** World Journal







Hindawi

Submit your manuscripts at http://www.hindawi.com



Algebra



Journal of Probability and Statistics



International Journal of Differential Equations





Complex Analysis





Mathematical Problems in Engineering



Abstract and Applied Analysis



Discrete Dynamics in Nature and Society



International Journal of Mathematics and Mathematical Sciences





Journal of **Function Spaces** 



International Journal of Stochastic Analysis

