L_1 SPACES FAIL A CERTAIN APPROXIMATIVE PROPERTY

AREF KAMAL

Department of Mathematics and Computer Sciences
U.A.E. University
P.O. Box 17551
Al-Ain, UNITED ARAB EMIRATES

(Received May 4, 1994)

ABSTRACT. In this paper the author studies some cases of Banach space that does not have the property P_1 . He shows that if $X = \ell_1$ or $L_1(\mu)$ for some non-purely atomic measure μ , then X does not have the property P_1 . He also shows that if $X = \ell_\infty$ or $C(Q)$ for some infinite compact Hausdorff space Q , then X^* does not have the property P_1 .

KEY WORDS AND PHRASES: Property P_1 , classical Banach spaces ℓ_1 , $L_1(\mu)$, ℓ_1^n , compact width
1991 AMS SUBJECT CLASSIFICATION CODES: 41A65.

1. INTRODUCTION

The Banach space X is said to have the property P_1 , if for each $\epsilon > 0$ and each $r > 0$, there is $\delta > 0$, such that for each x and y in X , there is $z \in \overline{B(x, \epsilon)}$ satisfying that for each θ with $0 < \theta < \delta$

$$B(x, r + \delta) \cap B(y, r + \theta) \subseteq B(z, r + \theta)$$

where $B(x, r)$ is the open ball of radius r and centered at x , and $\overline{B(x, r)}$ is its clouser u

The property P_1 plays an important role in approximation theory, and many authors used it. This property appears in approximation by compact operators, simultaneous approximation and other areas (see for example Roversi [1], Lau [2], Mach [3] and Kamal [4]). Mach [3] showed that if X is uniformly convex then it has the property P_1 [3], and that if $X = C(Q)$, or $X = B(Q)$ then X has the property P_1 [4]. Mach [4, page 259] asked if the space $L_1(\mu)$ has the property P_1 .

In this paper the author studies some cases of normed linear space X , for which X does not have the property P_1 . In section 2, it is shown that if $X = \ell_1$ then X does not have the property P_1 , and in section 3, it is shown that if μ is a non-purely atomic measure, then $L_1(\mu)$ does not have the property P_1 . These two results give a negative answer for the question of Mach [4]. In section 3, it will be shown also that if $X = (\ell_\infty)^*$, or $X = (C(Q))^*$, where Q is an infinite compact Hausdorff space, then X does not have the property P_1 .

In this paper ℓ_1 is the Banach space of all real sequences $x = \{x_i\}$ satisfying that $\sum |x_i| < \infty$, together with the norm $\|x\| = \sum |x_i|$. Also ℓ_1^n is the Banach space of all real n -tuples $x = (x_1, x_2, \dots, x_n)$ together with the norm $\|x\| = \sum_{i=1}^n |x_i|$.

2. ℓ_1 DOES NOT HAVE THE PROPERTY P_1

The proof of the fact that ℓ_1 does not have the property P_1 depends on the behavior of the property P_1 in ℓ_1^n . In Lemma 2.3, it will be shown that if $\epsilon > 0$ is fixed, and δ_n corresponds to ϵ for $X = \ell_1^n$ in Lemma 2.1, then $\delta_n \rightarrow 0$ when $n \rightarrow \infty$, so using the fact that ℓ_1^n is a norm-one-complemented subspace of ℓ_1 , it will be shown in Theorem 2.4, that ℓ_1 does not have the property P_1 .

LEMMA 2.1. If the Banach space X has the property P_1 then for each $\epsilon > 0$, there is $\delta > 0$ such that for each $y \in X$, there is $z \in \overline{B(0, \epsilon)}$ such that if $0 < \theta < \delta$ then

$$B(0, 1 + \delta + \theta) \cap B(y, 1 + \theta) \subseteq B(z, 1 + \theta).$$

PROOF. Let $r = 1$ and let $\epsilon > 0$ be given. By the definition of the property P_1 there is $\delta > 0$ such that for each x and y in X , there is $z \in B(x, \epsilon)$ satisfying the following; for each θ' such that $0 < \theta' < \delta'$

$$B(x, 1 + \delta') \cap B(y, 1 + \theta') \subseteq B(z, 1 + \theta').$$

Let $x = 0$ and $\delta = 1/2 \delta'$, then for all θ satisfying $0 < \theta < \delta$;

$$B(0, 1 + \delta + \theta) \cap B(y, 1 + \theta) \subseteq B(0, 1 + \delta') \cap B(y, 1 + \theta) \subseteq B(z, 1 + \theta).$$

LEMMA 2.2. Let $n \geq 3$ be a positive integer, let $\delta > 0$ be given and let $(z_1, \dots, z_n)$ be an n -tuple of real numbers

If $\sum_{i=1}^n z_i \geq \delta$, and for each $i \leq n-1$

$$z_1 + \dots + z_{i-1} - z_i + z_{i+1} + \dots + z_n \leq -\delta,$$

then $\sum_{i=1}^n |z_i| \geq (2n-3)\delta$.

PROOF. For each $i = 1, 2, \dots, n$, let $y_i = z_1 + \dots + z_{i-1} - z_i + z_{i+1} + \dots + z_n$, then $\sum_{i=1}^n y_i = (n-2) \sum_{i=1}^n z_i$, thus

$$y_n = (n-2) \sum_{i=1}^n z_i - \sum_{i=1}^{n-1} y_i.$$

Therefore

$$\sum_{i=1}^n |z_i| \geq |y_n| \geq (n-2)\delta + (n-1)\delta = (2n-3)\delta.$$

LEMMA 2.3. Let $n \geq 3$ be a positive integer and let $\delta > 0$ be a given real number such that $(2n-5)\delta \leq 1$. Then the element $x_n = (\delta, \delta, \dots, \delta, -(n-2)\delta)$ in ℓ_1^n satisfies the following conditions

(1) $\forall \theta$ such that $\theta > 0$

$$B(0, 1 + \delta + \theta) \cap B(x_n, 1 + \theta) \neq \phi.$$

(2) If $z \in \ell_1^n$ and for each θ with $0 < \theta < \delta$;

$$B(0, 1 + \delta + \theta) \cap B(x_n, 1 + \theta) \subseteq B(z, 1 + \theta),$$

then $\|z\| \geq (2n-3)\delta$

PROOF. Let $\{e'_i\}_{i=1}^n$ be the standard basis in ℓ_1^n , that is $e'_i = (x_1^i, x_2^i, \dots, x_n^i)$, where $x_j^i = 1$ if $i = j$ and $x_j^i = 0$ if $i \neq j$ and let

$$x_n = \delta \sum_{i=1}^{n-1} e'_i - (n-2)\delta e'_n = (\delta, \delta, \dots, \delta, -(n-2)\delta) \in \ell_1^n.$$

Then $\|x_n\| = (n-1)\delta + (n-2)\delta = (2n-3)\delta \leq 1 + 2\delta \leq 2 + \delta$, therefore for each θ such that $0 < \theta < \delta$

$$B(0, 1 + \delta + \theta) \cap B(x_n, 1 + \theta) \neq \phi.$$

Assume that $z = (z_1, z_2, \dots, z_n) \in \ell_1^n$ is such that for each θ with $0 < \theta < \delta$

$$B(0, 1 + \delta + \theta) \cap B(x_n, 1 + \theta) \subseteq B(z, 1 + \theta).$$

It will be shown that

- (1) $\sum_{i=1}^n z_i \geq \delta$, and
- (2) for each $i \leq n-1$

$$z_1 + \dots + z_{i-1} - z_i + z_{i+1} + \dots + z_n \leq -\delta.$$

If these are true then by Lemma 2.2, $\|z\| = \sum_{i=1}^n |z_i| \geq (2n-3)\delta$

- (1) Assume that $z_1 + \dots + z_n < \delta$. Let

$$\begin{aligned} y &= \delta \sum_{i=1}^{n-2} e'_i + \frac{1+\delta}{2} e'_{n-1} + \left[\frac{1-(2n-5)}{2} \right] e'_n \\ &= \left(\delta, \dots, \delta, \frac{1+\delta}{2}, \frac{1-(2n-5)}{2} \right) \in \ell_1^n. \end{aligned}$$

Then

$$\|y\| = (n-2)\delta + \frac{1+\delta}{2} + \frac{1-(2n-5)\delta}{2} = 1 + \delta.$$

On the other hand

$$\begin{aligned} \|y - x_n\| &= \left| \frac{1+\delta}{2} - \delta \right| + \left| \frac{1-(2n-5)\delta}{2} + (n-2)\delta \right| \\ &= 1. \end{aligned}$$

Thus, for each θ such that $0 < \theta < \delta$,

$$y \in B(0, 1 + \delta + \theta) \cap B(x_n, 1 + \theta).$$

But

$$\begin{aligned} \|y - z\| &= \sum_{i=1}^n |y_i - z_i| \geq \left| \sum_{i=1}^n y_i - \sum_{i=1}^n z_i \right| = \left| 1 + \delta - \sum_{i=1}^n z_i \right| \\ &= 1 + \left(\delta - \sum_{i=1}^n z_i \right) \\ &> 1, \end{aligned}$$

so for any $\theta < \left(\delta - \sum_{i=1}^n z_i \right)$, $y \notin B(z, 1 + \theta)$.

- (2) Assume that for a certain $i_0 \leq n-1$

$$z_1 + \dots + z_{i_0-1} - z_{i_0} + z_{i_0+1} + \dots + z_n > -\delta.$$

Let

$$y = \left(\frac{1+\delta}{2} \right) e'_{i_0} - \left(\frac{1+\delta}{2} \right) e'_n = \left(0, 0, \dots, 0, \frac{1+\delta}{2}, 0, \dots, 0, -\left(\frac{1+\delta}{2} \right) \right) \in \ell_1^n.$$

i_0 -th term

Then

$$\|y\| = 1 + \delta,$$

and

$$\begin{aligned}
\|y - x_n\| &= (n-2)\delta + \left| \frac{1+\delta}{2} - \delta \right| + \left| \frac{1+\delta}{2} + (n-2)\delta \right| \\
&= (n-2)\delta + \frac{1-\delta}{2} + \frac{1-(2n-5)\delta}{2} \\
&= 1.
\end{aligned}$$

Thus, for each θ such that $0 < \theta < \delta$, $y \in B(0, 1 + \delta + \theta) \cap B(x_n, 1 + \theta)$. But

$$\begin{aligned}
\|y - z\| &= \sum_{i=1}^n |y_i - z_i| \\
&\geq |(y_1 + \dots + y_{i_0-1} - y_{i_0} + y_{i_0+1} + \dots + y_n) - (z_1 + \dots + z_{i_0-1} - z_{i_0} + z_{i_0+1} + \dots + z_n)| \\
&= |-1 - \delta - (z_1 + \dots + z_{i_0-1} - z_{i_0} + z_{i_0+1} + \dots + z_n)| \\
&= |1 + [\delta + (z_1 + \dots + z_{i_0-1} - z_{i_0} + z_{i_0+1} + \dots + z_n)]| \\
&> 1.
\end{aligned}$$

Thus, for some $\theta > 0$, $y \notin B(z, 1 + \theta)$.

THEOREM 2.4. ℓ_1 does not have the property P_1

PROOF. It will be shown that for each $\delta > 0$, there is $x_\delta \in \ell_1$, such that if $z \in \ell_1$ and for all θ with $0 < \theta < \delta$ it is true that $B(0, 1 + \delta + \theta) \cap B(x_\delta, 1 + \theta) \subseteq B(z, 1 + \theta)$, then $\|z\| > \frac{1}{2}$. Let $\{e_i\}_{i=1}^\infty$ be the standard basis in ℓ_1 , and let $\delta > 0$ be given. If $\delta > 1$ then for each $\theta > 0$

$$B(0, 1 + 1 + \theta) \cap B(x_1, 1 + \theta) \subseteq B(0, 1 + \delta + \theta) \cap B(x_1, 1 + \theta).$$

Thus one can take x_1 to be x_δ . So without loss of generality one may assume that $\delta \leq 1$

Let $n \geq 3$ be a positive integer satisfying $(2n-5)\delta \leq 1$ and $(2n-3)\delta > \frac{1}{2}$, and let x_n be as in Lemma 2.3. Define

$$x_\delta = \delta \sum_{i=1}^{n-1} e_i - (n-2)\delta e_n = (\delta, \delta, \dots, \delta, -(n-2)\delta, 0, 0, \dots) \in \ell_1.$$

Then $\|x_\delta\| = \|x_n\| \leq 2 + \delta$, thus

$$B(0, 1 + \delta + \theta) \cap B(x_\delta, 1 + \theta) \neq \emptyset \quad \text{for } 0 < \theta < \delta.$$

Let $P_n : \ell_1 \rightarrow \ell_1^n$ be the mapping defined by $P_n(\{x_i\}_{i=1}^\infty) = \{x_i\}_{i=1}^n$. By the construction of x_δ its image under P_n is the element x_n .

Assume that for some $z \in \ell_1$

$$B(0, 1 + \delta + \theta) \cap B(x_\delta, 1 + \theta) \subseteq B(z, 1 + \theta) \quad 0 < \theta < \delta,$$

then in ℓ_1^n

$$B(0, 1 + \delta + \theta) \cap B(x_n, 1 + \theta) \subseteq B(P_n(z), 1 + \theta) \quad 0 < \theta < \delta,$$

Thus by Lemma 2.3 $\|P_n(z)\| \geq (2n-3)\delta > \frac{1}{2}$. Therefore

$$\|z\| \geq \|P_n(z)\| > \frac{1}{2}.$$

3. OTHER SPACES THAT DO NOT HAVE THE PROPERTY P_1

The subspace Y of X is called a norm-one-complemented subspace of X if there is a linear projection $P : X \rightarrow Y$ satisfying that $\|P\| = 1$. If A is a subset of X , and $x \in X$ then

$$d(x, A) = \inf\{\|x - y\|; y \in A\},$$

and if B is another subset of X , then the deviation of A from B is defined by

$$\delta(A, B) = \sup\{d(x, B); x \in A\}.$$

The compact width of A in X is defined by

$$a(A, X) = \inf\{\delta(A, K); K \text{ is a compact subset of } X\}.$$

The compact width is said to be attained if there is a compact subset K of X satisfying that $a(A, X) = \delta(A, K)$

In this section it will be shown that if $X = (C(Q))^*$, where Q is an infinite compact Hausdorff space, $X = (\ell_\infty)^*$, or $X = L_1(\mu)$ where μ is non-purely atomic measure, then X does not have the property P_1 .

The proof of the following proposition is elementary

PROPOSITION 3.1. Let X be a Banach space that has the property P_1 , and let Y be a closed subspace of X . If Y is a norm-one-complemented subspace of X , then Y has the property P_1

COROLLARY 3.2. If μ is non-purely atomic measure then $L_1(\mu)$ does not have the property P_1

PROOF. By Feder [5, Theorem 2], $L_1[0, 1]$ has a subset A for which the compact width $a(A, L_1[0, 1])$ is not attained, thus by Kamal [6, Theorem 4.3] $L_1[0, 1]$ does not have the property P_1 , but by Lacy [7, sec 8], $L_1[0, 1]$ is a norm-one-complemented subspace of $L_1(\mu)$, therefore by Proposition 3.1, $L_1(\mu)$ does not have the property P_1 .

NOTE 3.3. Theorem 2.4 together with Corollary 3.2 give a negative answer to the question of Mach [4, page 259].

COROLLARY 3.4. If $X = \ell_\infty$ or $X = C(Q)$ for some compact infinite Hausdorff space Q . Then X^* does not have the property P_1

PROOF. If $X = \ell_\infty$ then ℓ_1 is a norm-one-complemented subspace of X^* , and if $X = C(Q)$ then by Kamal [8, Lemma 3.2], ℓ_1 is a norm-one-complemented subspace of X^* , in both cases one concludes by Proposition 3.1 that X^* does not have the property P_1 .

REFERENCES

- [1] ROVERSI, M., Best approximation of bounded functions by continuous functions, *J. Approx. Theory* **41** (1984), 135-148
- [2] LAU, K., Approximation by continuous vector valued functions, *Studia Math.* **68** (1980), 291-298
- [3] MACH, J., Best simultaneous approximation of bounded functions with values in certain Banach spaces, *Math. Ann.* **240** (1979), 157-164.
- [4] MACH, J., On the existence of best simultaneous approximation, *J. Approx. Theory* **25** (1979), 258-265.
- [5] FERDER, M., On certain subset of $L_1[0, 1]$ and non-existence of best approximation in some spaces of operators, *J. Approx. Theory* **29** (1980), 170-177.
- [6] KAMAL, A., On certain diameters of bounded sets, *J. Port. Math.* **51** (1994), 321-333
- [7] LACY, H.E., *The Isometric Theory of Classical Banach Spaces*, Springer-Verlag, 1974
- [8] KAMAL, A., On proximality and sets of operators III, Approximation by finite rank operators on spaces of continuous functions, *J. Approx. Theory* **47** (1986), 156-171.