

### THE DIOPHANTINE EQUATION

$$x^2 + 3^m = y^n$$

S. AKHTAR ARIF and FADWA S. ABU MURIEFAH

Department of Mathematics  
Girls College of Education  
Al-Riyadh, SAUDI ARABIA

(Received March 11, 1996 and in revised form May 31, 1996)

**ABSTRACT.** The object of this paper is to prove the following

**THEOREM.** Let  $m$  be odd. Then the diophantine equation  $x^2 + 3^m = y^n$ ,  $n \geq 3$  has only one solution in positive integers  $x, y, m$  and the unique solution is given by  $m = 5 + 6M$ ,  $x = 10 \cdot 3^{3M}$ ,  $y = 7 \cdot 3^{2M}$  and  $n = 3$ .

**KEY WORDS AND PHRASES:** Diophantine equation.

**1992 AMS SUBJECT CLASSIFICATION CODES:** 11D41.

#### INTRODUCTION

It is well known that there is no general method for determining all integral solutions  $x$  and  $y$  for a given diophantine equation  $ax^2 + bx + c = dy^n$ , where  $a, b, c$  and  $d$  are integers,  $a \neq 0$ ,  $b^2 - 4ac \neq 0$ ,  $d \neq 0$ , but we know that it has only a finite number of solutions when  $n \geq 3$ . This was first shown by Thue [1]

The first result for the title equation for general  $n$  is due to Lebesgue [2] who proved that when  $m = 0$  there is no solution, for  $m = 1$ , Nagell [3] has proved that it has no solution and in 1993 Cohn [4] has given another proof for this case.

The proof of the theorem is divided into two main cases  $(3, x) = 1$  and  $3|x$ . It is sufficient to consider  $x$  a positive integer.

To prove the theorem we need the following

**LEMMA** (Nagell [5]). The equation  $3x^2 + 1 = y^n$ , where  $n$  is an odd integer  $\geq 3$  has no solution in integers  $x$  and  $y$  for  $y$  odd and  $\geq 1$ .

**PROOF OF THEOREM.** Suppose  $m = 2k + 1$ . Since the result is known for  $m = 1$  we shall assume that  $k > 0$ . The case when  $x$  is odd, can be easily eliminated since  $y^n \equiv 0 \pmod{8}$ , so we assume that  $x$  is even.

**CASE 1: Let  $(3, x) = 1$ .** First let  $n$  be odd, then there is no loss of generality in considering  $n = p$  an odd prime. Thus  $x^2 + 3^{2k+1} = y^p$ . Then from [6, Theorem 1] we have only two possibilities and they are

$$x + 3^k \sqrt{-3} = (a + b\sqrt{-3})^p \tag{1}$$

where  $y = a^2 + 3b^2$  and

$$x + 3^k \sqrt{-3} = \left( \frac{a + b\sqrt{-3}}{2} \right)^3, \quad a \equiv b \equiv 1 \pmod{2} \tag{2}$$

where  $y = \frac{a^2 + 3b^2}{4}$ , for some rational integers  $a$  and  $b$ .

In (1) since  $y = a^2 + 3b^2$  and  $y$  is odd so only one of  $a$  or  $b$  is odd and the other is even. Equating imaginary parts we get

$$3^k = b \sum_{r=0}^{\frac{p-1}{2}} \binom{p}{2r+1} a^{p-2r-1} (-3b^2)^r.$$

So  $b$  is odd. Since 3 does not divide the term inside  $\sum$  we get  $b = \pm 3^k$ . Hence

$$\pm 1 = \sum_{r=0}^{\frac{p-1}{2}} \binom{p}{2r+1} a^{p-2r-1} (-3^{2k+1})^r.$$

This is equation (1) in [6], and Lemmas 4 and 5 in [6] show that both the signs are impossible. Hence (1) gives rise to no solutions.

Now consider equation (2). By equating imaginary parts we obtain

$$8 \cdot 3^k = b(3a^2 - 3b^2). \quad (3)$$

If  $b = \pm 1$  in (3) we get

$$\pm 8 \cdot 3^k = 3a^2 - 3.$$

The case  $k = 1$  can be easily eliminated, so suppose  $k > 1$  then

$$\pm 8 \cdot 3^{k-1} = a^2 - 1.$$

This equation has the only solution  $a = \pm 5$ ,  $k = 2$  and so  $y = \frac{a^2+3b^2}{4} = (25+3)/4 = 7$ . Hence from (2)  $x = \left| \frac{a^2-9ab^2}{8} \right| = 10$

If  $b = \pm 3^\lambda$ ,  $0 < \lambda < k$ , then (3) becomes  $\pm 8 \cdot 3^{k-\lambda-1} = a^2 - 3^{2\lambda}$ , and this is not possible modulo 3 if  $k - \lambda - 1 > 0$ . So  $k - \lambda - 1 = 0$ , that is  $\pm 8 = a^2 - 3^{2(k-1)}$ , and we can reject the positive sign modulo 3. So we have  $a^2 - 3^{2(k-1)} = -8$ , which has the only solution  $a = \pm 1$ ,  $k = 2$  and  $x = 10$ . Finally if  $b = \pm 3^k$  then  $\pm 8 = 3a^2 - 3^{2k+1}$ , and this is not true modulo 3.

Now if  $n$  is even, then from the above it is sufficient to consider  $n = 4$ , hence  $(y^2+x)(y^2-x) = 3^{2k+1}$ . Since  $(3, x) = 1$ , we get

$$y^2 + x = 3^{2k+1} \quad \text{and} \quad y^2 - x = 1,$$

by adding these two equations we get  $2y^2 = 3^{2k+1} + 1$ , which is impossible modulo 3.

**CASE 2.** Let  $3|x$ . Then of course  $3|y$ . Suppose that  $x = 3^u X$ ,  $y = 3^\nu Y$  where  $u > 0$ ,  $\nu > 0$  and  $(3, X) = (3, Y) = 1$ . Then  $3^{2u} X^2 + 3^{2k+1} = 3^{n\nu} Y^n$ . There are three possibilities.





1.  $2u = \min(2u, 2k+1, n\nu)$ . Then by cancelling  $3^{2u}$  we get  $X^2 + 3^{2(k-u)+1} = 3^{m\nu-2u} Y^n$ , and considering this equation modulo 3 we deduce that  $n\nu - 2u = 0$ , then  $x^2 + 3^{2(k-u)+1} = Y^n$ , with  $(3, X) = 1$ . If  $k - u = 0$ , this equation has no solution [3,4] and if  $k - u > 0$ , as proved above this equation has a solution only if  $k - u = 2$  and  $n = 3$ , so  $n\nu = 3\nu = 2u$  that is  $3|u$ , let  $u = 3M$  then  $k = 2 + 3M$  and  $m = 5 + 6M$ . So this equation has a solution only if  $m = 5 + 6M$  and the solution is given by  $X = 10$ ,  $Y = 7$ . Hence the solution of our title equation is  $x = 10 \cdot 3^u = 10 \cdot 3^{3M}$  and  $y = 7 \cdot 3^\nu = 7 \cdot 3^{2M}$ .

2.  $2k+1 = \min(2u, 2k+1, n\nu)$ . Then  $3^{2u-2k-1} X^2 + 1 = 3^{n\nu-2k-1} Y^n$  and considering this equation modulo 3 we get  $n\nu - 2k - 1 = 0$ , so  $n$  is odd and  $3(3^{u-k-1} X)^2 + 1 = Y^n$ , by the lemma this equation has no solution.

3.  $n\nu = \min(2u, 2k+1, n\nu)$ . Then  $3^{2u-n\nu} X^2 + 3^{2k+1-n\nu} = Y^n$  and this is possible modulo 3 only if  $2u - n\nu = 0$  or  $2k+1 - n\nu = 0$  and both of these cases have already been discussed. This concludes the proof.

## REFERENCES

- [1] THUE, A., Über die Unlösbarkeit der Gleichung  $ax^2 + bx + c = dy^n$  in grossen ganzen Zahlen  $x$  und  $y$ , *Arch. Math. og Naturvidenskab, Kristiania*, Bd XXXIV (1916), 1-6.
- [2] LEBESGUE, V.A., Sur l'impossibilité en nombres entières de l'équation  $x^m = y^2 + 1$ , *Nouvelles Ann. des Math.* 9 (1) (1850), 178-181.
- [3] NAGELL, T., Sur l'impossibilité de quelques équations à deux indéterminées, *Norsk Math. Forenings Skrifter*, Kristiania, Ser. I, 13 (1923), 65-82.
- [4] COHN, J.H.E., The diophantine equation  $x^2 + 3 = y^n$ , *Glasgow Math. J.*, 35 (1993), 203-206.
- [5] NAGELL, T., Contributions to the theory of a category of diophantine equations of the second degree with two unknown, *Nova Acta Reg. Soc. Upsal.*, Ser IV, 16 (1955), 1-38.
- [6] COHN, J.H.E., The diophantine equation  $x^2 + c = y^n$ , *Acta Arith.* 65 (4) (1993), 367-381



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

