

NONWANDERING SETS OF MAPS ON THE CIRCLE

SEUNG WHA YEOM, KYUNG JIN MIN, and SEONG HOON CHO

(Received 30 April 1997 and in revised form 25 July 1997)

ABSTRACT. Let f be a continuous map of the circle S^1 into itself. And let $R(f)$, $\Lambda(f)$, $\Gamma(f)$, and $\Omega(f)$ denote the set of recurrent points, ω -limit points, γ -limit points, and nonwandering points of f , respectively. In this paper, we show that each point of $\Omega(f) \setminus \overline{R(f)}$ is one-side isolated, and prove that

- (1) $\Omega(f) \setminus \Gamma(f)$ is countable and
- (2) $\Lambda(f) \setminus \Gamma(f)$ and $\overline{R(f)} \setminus \Gamma(f)$ are either empty or countably infinite.

Keywords and phrases. Nonwandering point, recurrent point, one-side isolated.

1991 Mathematics Subject Classification. 58F.

1. Introduction. Let I be the unit interval, S^1 the circle, and X a topological space. And let $C^0(X, X)$ denote the set of continuous maps from X into itself. For any $f \in C^0(X, X)$, let $P(f)$, $R(f)$, $\Lambda(f)$, $\Gamma(f)$, and $\Omega(f)$ denote the set of periodic points, recurrent points, ω -limit points, γ -limit points and nonwandering points of f , respectively.

For any $f \in C^0(I, I)$, in 1980, Z. Nitecki [6] has proved that each point of $\Omega(f) \setminus \overline{P(f)}$ is isolated in $\Omega(f)$ if f is piecewise monotone and is not flat on any subinterval of I . In 1984, J. C. Xiong [7] has proved that each point of $\Omega(f) \setminus \overline{P(f)}$ is one-side isolated in $\Omega(f)$, for a continuous self map of interval I . And, in 1988, J. C. Xiong [9] also showed that $\Omega(f) \setminus \Gamma(f)$ is countable and that $\Lambda(f) \setminus \Gamma(f)$ and $\overline{P(f)} \setminus \Gamma(f)$ are either empty or countably infinite.

In this paper, we obtain the following similar results for maps of the circle:

THEOREM 1.1. *Let $f \in C^0(S^1, S^1)$. Then each point of $\Omega(f) \setminus \overline{R(f)}$ is one-side isolated in $\Omega(f)$.*

THEOREM 1.2. *Let $f \in C^0(S^1, S^1)$. Then*

- (1) $\Omega(f) \setminus \Gamma(f)$ is countable.
- (2) $\Lambda(f) \setminus \Gamma(f)$ and $\overline{R(f)} \setminus \Gamma(f)$ are either empty or countably infinite.

2. Preliminaries and definitions. Let X be a compact metric space and $f \in C^0(X, X)$. For any positive integer n , we define f^n inductively by $f^1 = f$ and $f^{n+1} = f \circ f^n$. Let f^0 denote the identity map of X . The *forward orbit* $\text{Orb}(x)$ of $x \in X$ is the set $\{f^k(x) \mid k = 0, 1, 2, \dots\}$. Usually, the forward orbit of x is simply called the *orbit* of x .

A point $x \in X$ is called a *periodic point* of f if, for some positive integer n , $f^n(x) = x$. The period of x is the least such integer n . We denote the set of periodic points of f by $P(f)$. A point $x \in X$ is called a *recurrent point* of f if there exists a sequence $\{n_i\}$ of positive integers with $n_i \rightarrow \infty$ such that $f^{n_i}(x) \rightarrow x$. We denote the set of recurrent

points of f by $R(f)$. A point $x \in X$ is called a *nonwandering point* of f if, for every neighborhood U of x , there exists a positive integer m such that $f^m(U) \cap U \neq \emptyset$. We denote the set of nonwandering points of f by $\Omega(f)$.

A point $y \in X$ is called an ω -*limit point* of x if there exists a sequence $\{n_i\}$ of positive integers with $n_i \rightarrow \infty$ such that $f^{n_i}(x) \rightarrow y$. We denote the set of ω -limit points of x by $\omega(x)$. Define $\Lambda(f) = \bigcup_{x \in X} \omega(x)$. A point $y \in X$ is called an α -*limit point* of x if there exists a sequence $\{n_i\}$ of positive integers with $n_i \rightarrow \infty$ and a sequence $\{y_i\}$ of points such that $f^{n_i}(y_i) = x$ and $y_i \rightarrow y$. The symbol $\alpha(x)$ denotes the set of α -limit points of x . A point $y \in X$ is called a γ -*limit point* of x if $y \in \omega(x) \cap \alpha(x)$. The symbol $\gamma(x)$ denotes the set of γ -limit points of x and $\Gamma(f) = \bigcup_{x \in X} \gamma(x)$.

Let R be the set of reals and Z be the set of integers. Formally, we think of the circle S^1 as R/Z and use $\pi: R \rightarrow R/Z$ to denote the canonical projection. In fact, the map $\pi: R \rightarrow S^1$ is an example of a covering map since it wraps R around S^1 without doubling back (i.e., without critical points). To study the dynamics of the circle map, it is helpful to use a *lifting*. Let f be a continuous map on the circle. We say that a continuous map F from R into itself is a lifting of f if $f \circ \pi = \pi \circ F$. We use the following notations throughout this paper.

Let $a, b \in S^1$ with $a \neq b$, and let $A \in \pi^{-1}(a), B \in \pi^{-1}(b)$ with $|A - B| < 1$ and $A < B$. Then we write $\pi((A, B)), \pi([A, B]), \pi([A, B])$ and $\pi((A, B])$ to denote the open, closed, and half-open arcs from a counterclockwise to b , respectively, and we denote it by $(a, b), [a, b], [a, b)$, and $(a, b]$. For $x, y \in [a, b]$ with $a \neq b$, let $X \in \pi^{-1}(x), Y \in \pi^{-1}(y)$ with $X, Y \in [A, B]$, then we define for $x, y \in [a, b], x > y$ if and only if $X > Y$. Let C be a subset of a closed arc $[a, b]$, then we define $\sup C = \pi(\sup(\pi^{-1}(C) \cap [A, B]))$ and $\inf C = \pi(\inf(\pi^{-1}(C) \cap [A, B]))$.

In particular, for $a, b, c \in S^1, a < b < c$ means that b lies in the open arc (a, c) , that is, $b \in (a, c)$.

Let X be I or S^1 and $Y \subset X$. Let $x \in Y$. A point $x \in X$ is said to be *left-sided isolated* (resp., *right-sided isolated*) in Y if, for some $\epsilon > 0$, $(x - \epsilon, x) \cap Y = \emptyset$ (resp., $(x, x + \epsilon) \cap Y = \emptyset$). A point x is said to be *one-side isolated* in Y if x is either left-side or right-side isolated in Y , and a point x which is both a right-sided and a left-sided isolated in Y is said to be *isolated* in Y .

Let $x \in S^1$ and $f \in C^0(S^1, S^1)$ be given. Then we use the symbols $\omega_+(x)$ (resp., $\omega_-(x)$) to denote the set of all points $y \in S^1$ such that there exists a sequence $\{n_i\}$ of positive integers with $n_i \rightarrow \infty$ such that $f^{n_i}(x) \rightarrow y$ and $y < \dots < f^{n_i}(x) < \dots < f^{n_2}(x) < f^{n_1}(x)$ (resp. $f^{n_1}(x) < f^{n_2}(x) < \dots < f^{n_i}(x) < \dots < y$). It is clear that if $x \notin P(f)$, then $\omega(x) = \omega_+(x) \cup \omega_-(x)$. Define $\Lambda_+(f) = \bigcup_{x \in S^1} \omega_+(x)$ and $\Lambda_-(f) = \bigcup_{x \in S^1} \omega_-(x)$.

Also, we use the symbols $\alpha_+(x)$ (resp. $\alpha_-(x)$) to denote the set of all points $y \in S^1$ such that there exists a sequence $\{n_i\}$ of positive integers with $n_i \rightarrow \infty$ and a sequence $\{x_i\}$ of points such that $x_i \rightarrow y, f^{n_i}(x_i) = x$ for every $i > 0$ and $y < \dots < x_i < \dots < x_2 < x_1$ (resp. $x_1 < x_2 < \dots < x_i < \dots < y$). It is clear that if $x \notin P(f)$, then $\alpha(x) = \alpha_+(x) \cup \alpha_-(x)$.

Define $\gamma_+(x) = \omega_+(x) \cap \alpha_+(x)$ and $\gamma_-(x) = \omega_-(x) \cap \alpha_-(x)$. Also, we define $\Gamma_+(f) = \bigcup_{x \in S^1} \gamma_+(x)$ and $\Gamma_-(f) = \bigcup_{x \in S^1} \gamma_-(x)$.

Let Y be an arc in S^1 and let \bar{Y} denote the closure of Y as usual. A point $y \in S^1$ is

called a *right-sided* (resp., *left-sided*) *accumulation point* of Y if, for any $z \in S^1$, $(y, z) \cap Y \neq \emptyset$ (resp. $(z, y) \cap Y \neq \emptyset$).

The right-side closure \bar{Y}_+ (resp. left-side closure \bar{Y}_-) is the union of Y and the set of right-sided (resp. left-sided) accumulation points of Y . A point which is both a right-sided and a left-sided accumulation point of Y is called a *two-sided accumulation point* of Y .

3. Main results. The following lemmas are founded in [3].

LEMMA 3.1. *Let $f \in C^0(S^1, S^1)$ and $x \in \Omega(f)$. Then we have $x \in \alpha(x)$.*

LEMMA 3.2. *Let $f \in C^0(S^1, S^1)$ and $I = [a, b]$ be an arc for some $a, b \in S^1$ with $a \neq b$, and let $I \cap P(f) = \emptyset$.*

- (a) *Suppose that there exists $x \in I$ such that $f(x) \in I$ and $x < f(x)$. Then*
 - (i) *if $y \in I, x < y$, and $f(y) \notin [y, b]$, then $[x, y]$ f -covers $[f(x), b]$, and*
 - (ii) *if $y \in I, y < x$, and $f(y) \notin [y, b]$, then $[y, x]$ f -covers $[f(x), b]$.*
- (b) *Suppose that there exists $x \in I$ such that $f(x) \in I$ and $x > f(x)$. Then*
 - (i) *if $y \in I, x < y$, and $f(y) \notin [a, y]$, then $[x, y]$ f -covers $[a, f(x)]$, and*
 - (ii) *if $y \in I, y < x$, and $f(y) \notin [a, y]$, then $[y, x]$ f -covers $[a, f(x)]$.*

LEMMA 3.3. *Let $f \in C^0(S^1, S^1)$. Then we have*

$$P(f) \subset R(f) \subset \Gamma(f) \subset \overline{R(f)} \subset \Lambda(f) \subset \Omega(f) \subset CR(f). \quad (1)$$

The following lemma is due to [5]

LEMMA 3.4. *Let $f \in C^0(S^1, S^1)$, and let $K \subset S^1$ with $f(K) \subset K$. If $x \in \Omega(f) \setminus K$, then $f^n(x) \notin K^\circ$ for any $n \geq 1$.*

The idea of the proof of the following lemma is due to [7].

LEMMA 3.5. *Let $f \in C^0(S^1, S^1)$, and let $K \subset S^1$ have only finitely many connected components and $f(K) = K$. Then we have $\bar{K} \setminus K \subset P(f)$.*

PROOF. By continuity of f , we have $f(\bar{K}) \subset \overline{f(K)}$. And by the compactness of \bar{K} , $f(\bar{K}) \subset S^1$ is closed. Thus, $\overline{f(K)} \subset \overline{f(\bar{K})} = f(\bar{K})$. Therefore, $f(\bar{K}) = \overline{f(K)} = \bar{K}$. Hence, for each $x \in \bar{K} \setminus K$, there exists $x' \in \bar{K} \setminus K$ such that $f(x') = x$, i.e., $f(\bar{K} \setminus K) = \bar{K} \setminus K$. By the finiteness of $\bar{K} \setminus K$, $\bar{K} \setminus K \subset P(f)$. \square

PROPOSITION 3.6. *Let $f \in C^0(S^1, S^1)$. Suppose that $x \in \Omega(f) \setminus \overline{R(f)}$.*

- (1) *If $x \in \alpha_+(x)$, then there exists $z \in S^1$ such that $f^i(z, x) \cap (z, x) = \emptyset$ for all $i \geq 1$.*
- (2) *If $x \in \alpha_-(x)$, then there exists $u \in S^1$ such that $f^i(x, u) \cap (x, u) = \emptyset$ for all $i \geq 1$.*

PROOF. We only need to prove part (1). There exists $a, b \in S^1$ such that $x \in (a, b)$ and $(a, b) \cap \text{Orb}(x) = \emptyset$. Let $V = (a, x)$ and let $W = \bigcup_{i=0}^{\infty} f^i(V)$. Then $x \in \bar{W}$. Since $x \in \alpha_+(x)$, there exist a positive integer m and a point $y \in (x, b)$ such that $f^m(y) = x$. By Lemma 3.2,

$$[x, y] f^m\text{-covers } [a, x]. \quad (2)$$

We claim that $x \notin W$. To show this, suppose that $x \in W$. Then there exist a positive integer j and a point $x_0 \in (a, x)$ such that $f^j(x_0) = x$. By Lemma 3.2,

$$[x_0, x] f^j\text{-covers } [x, b]. \quad (3)$$

In particular, $[x_0, x] f^j\text{-covers } [x, y]$.

By (2),

$$[x_0, x] f^j\text{-covers } [x_0, y]. \quad (4)$$

Thus,

$$[x_0, x] f^{j+m}\text{-covers itself}, \quad (5)$$

and, hence, f^{j+m} has a periodic point in (a, b) , a contradiction. Hence, we have $x \in \overline{W} \setminus W$.

Assume that the proposition is false, i.e., for each $z \in (a, x)$, there is some $i \geq 1$ such that $(z, x) \cap f^i(z, x) \neq \emptyset$. Note that $V \subset f(W)$. Because, for each $y' \in V$, there is some $i \geq 1$ such that $(y', x) \cap f^i(y', x) \neq \emptyset$. There exists $x_0 \in (y', x)$ such that $f^i(x_0) \in (y', x)$. By Lemma 3.2, either

$$[x_0, x] f^i\text{-covers } [f^i(x_0), b] \quad \text{or} \quad [x_0, x] f^i\text{-covers } [a, f^i(x_0)]. \quad (6)$$

Particularly, either

$$[x_0, x] f^i\text{-covers } [x, b] \quad \text{or} \quad [x_0, x] f^j\text{-covers } [a, f^i(x_0)]. \quad (7)$$

If

$$[x_0, x] f^i\text{-covers } [x, b], \quad (8)$$

then

$$[x_0, x] f^j\text{-covers } [x, y]. \quad (9)$$

By (2),

$$[x, y] f^m\text{-covers } [x_0, x]. \quad (10)$$

Hence,

$$[x_0, x] f^{i+m}\text{-covers itself}. \quad (11)$$

Thus, f^{j+m} has a periodic point in (a, b) . This is a contradiction. Therefore,

$$[x_0, x] f^j\text{-covers } [a, f^i(x_0)]. \quad (12)$$

Thus, $y' \in f^i(x_0, x) \subset f^i(V) \subset f(W)$ since $y' \in (a, f^i(x_0))$. Thus, for each $i = 1, 2, 3, \dots, l-1$, $f^i(V) \cap f^{l+i}(V) \neq \emptyset$, and $f^{l+i}(V) \cap f^{2l+i}(V) \neq \emptyset, \dots$. Therefore, $U_i = \bigcup_{m=0}^{\infty} f^{ml+i}(V)$ is connected and $W = \bigcup_{i=0}^{l-1} U_i$ has only finitely many connected components. Now, by Lemma 3.5, $x \in \overline{W} \setminus W \subset P(f)$. This is in contradiction with the assumption of this proposition. \square

The following theorem follows immediately from the proposition.

THEOREM 3.7. *Let $f \in C^0(S^1, S^1)$. Then each point of $\Omega(f) \setminus \overline{R(f)}$ is one-side isolated in $\Omega(f)$.*

COROLLARY 3.8. *Let $f \in C^0(S^1, S^1)$. Then $\Omega(f) \setminus \overline{R(f)}$ is countable which is nowhere dense in S^1 .*

The following proposition is found in [1].

PROPOSITION 3.9. *Let $f \in C^0(S^1, S^1)$. Then we have*

- (1) $\overline{R(f)}_+ \setminus R(f) \subset \Lambda(f)_+$.
- (2) $\overline{R(f)}_- \setminus R(f) \subset \Lambda(f)_-$.

PROPOSITION 3.10. *Let $f \in C^0(S^1, S^1)$. Then we have $\overline{R(f)}_+ \cap \overline{R(f)}_- \setminus R(f) \subset \Gamma(f)$.*

PROOF. If $P(f) = \phi$, then we have the desired results since $\overline{R(f)} = \Gamma(f)$ [2]. Suppose that $P(f) \neq \phi$. Let $z \in \overline{R(f)}_+ \cap \overline{R(f)}_- \setminus R(f)$. Then there exist $a, b \in S^1$ with $a < b$ such that $z \in (a, b)$ and $(a, b) \cap \text{Orb}(z) = \phi$. By Proposition 3.9, $z \in \Lambda(f)_+ \cap \Lambda(f)_-$. Then there exist y_1, y_2 such that $a < y_1 < z < y_2 < b$ with $z \in \omega(y_1) \cap \omega(y_2)$. Since $\overline{P(f)} = \overline{R(f)}$ [4], $z \in \overline{P(f)}_+ \cap \overline{P(f)}_- \setminus P(f)$. Then there exists u_i of periodic point of f with $a < y_1 < u_1 < u_2 < \dots < z$ and $u_i \rightarrow z$. Let p_i be the period of u_i with respect to f . Then $f^{p_i}(u_i) = u_i$ for all $i \geq 1$. Then either $[u_i, z]$ f^{p_i} -covers $[a, u_i]$ or $[u_i, z]$ f^{p_i} -covers $[u_i, b]$.

We may assume that, for infinitely many i , either

$$[u_i, z] \text{ } f^{p_i}\text{-covers } [a, u_i] \quad \text{or} \quad [u_i, z] \text{ } f^{p_i}\text{-covers } [u_i, b]. \quad (13)$$

Then we consider two cases.

CASE I. $[u_i, z]$ f^{p_i} -covers $[a, u_i]$ for infinitely many i . There exists $z_i \in [u_i, z]$ such that $f^{p_i}(z_i) = y_1$. Since $u_i \rightarrow z$, $z_i \rightarrow z$. Thus, $z \in \alpha(y_1)$ and, hence, $z \in \omega(y_1) \cap \alpha(y_1) \subset \Gamma(f)$.

CASE II. $[u_i, z]$ f^{p_i} -covers $[u_i, b]$ for infinitely many i . There exists $z'_i \in [u_i, z]$ such that $f^{p_i}(z'_i) = y_1$. Since $u_i \rightarrow z$, $z'_i \rightarrow z$. Thus, $z \in \alpha(y_2)$ and, hence, $z \in \omega(y_2) \cap \alpha(y_2) \subset \Gamma(f)$. \square

The idea of the proof of the following lemma is due to [8].

LEMMA 3.11. *Let $f \in C^0(S^1, S^1)$ and $Y \subset S^1$. Then $\overline{Y} \setminus (\overline{Y}_+ \cap \overline{Y}_-)$ is countable.*

PROOF. For each $y \in \overline{Y}_+ \setminus \overline{Y}_-$, there is some $u_y \in S^1$ such that $(u_y, y) \cap Y = \phi$. The family of $\{(u_y, y) \mid y \in \overline{Y}_+ \setminus \overline{Y}_-\}$ is countable because it is disjoint. Hence, $\overline{Y}_+ \setminus \overline{Y}_-$ is countable. Similarly, $\overline{Y}_- \setminus \overline{Y}_+$ is also countable. Therefore,

$$\overline{Y} \setminus (\overline{Y}_+ \cap \overline{Y}_-) = (\overline{Y}_+ \setminus \overline{Y}_-) \cup (\overline{Y}_- \setminus \overline{Y}_+) \quad (14)$$

is countable. \square

THEOREM 3.12. *Let $f \in C^0(S^1, S^1)$. Then*

- (1) $\Omega(f) \setminus \Gamma(f)$ is countable.
- (2) $\Lambda(f) \setminus \Gamma(f)$ and $\overline{R(f)} \setminus \Gamma(f)$ are either empty or countably infinite.

PROOF. (1) We know that $\overline{R(f)} \setminus (\overline{R(f)}_+ \cap \overline{R(f)}_-)$ is countable by Lemma 3.11. By Proposition 3.10, $\overline{R(f)} \setminus \Gamma(f)$ is also countable. By Corollary 3.8, $\Omega(f) \setminus \overline{R(f)}$ is countable. Hence, $\Omega(f) \setminus \Gamma(f)$ is countable.

(2) It is easy to prove that $f(\omega(x)) = \omega(x)$ and $f(\overline{R(f)}) = \overline{R(f)}$ for $x \in S^1$. Hence, $f(\Lambda(f)) = \Lambda(f)$. Suppose that $\Lambda(f) \setminus \Gamma(f) \neq \emptyset$ (resp., $\overline{R(f)} \setminus \Gamma(f) \neq \emptyset$). Then we take $z_1 \in \Lambda(f) \setminus \Gamma(f)$ (resp., $z_1 \in \overline{R(f)} \setminus \Gamma(f)$). We can take $z_2 \in \Lambda(f) \setminus \Gamma(f)$ (resp., $z_2 \in \overline{R(f)} \setminus \Gamma(f)$) such that $z_1 = f(z_2)$. Continuing this process, we can take $z_i \in \Lambda(f) \setminus \Gamma(f)$ (resp., $z_i \in \overline{R(f)} \setminus \Gamma(f)$) such that $z_i = f(z_{i+1})$ for all $i = 1, 2, \dots$. Since $z_i \notin \Gamma(f)$ for all $i \geq 1$, the points z_1, z_2, \dots are pairwise disjoint. Hence, $\Lambda(f) \setminus \Gamma(f)$ (resp., $\overline{R(f)} \setminus \Gamma(f)$) is infinite and, hence, $\Lambda(f) \setminus \Gamma(f)$ (resp., $\overline{R(f)} \setminus \Gamma(f)$) is countably infinite. \square

ACKNOWLEDGEMENT. This research was supported by fund of SNUT.

REFERENCES

- [1] J. S. Bae, S. H. Cho, K. J. Min, and S. K. Yang, *Nonwandering points and the attracting centre of a map on the circle*, to appear.
- [2] ———, *Nonwandering points of a map on the circle*, J. Korean Math. Soc. **33** (1996), no. 4, 1115–1121. MR 97m:58061. Zbl 871.58031.
- [3] J. S. Bae, S. H. Cho, and S. K. Yang, *Relations between nonwandering points and turning points on the circle*, Bull. Korean Math. Soc. **32** (1995), no. 1, 57–66. MR 96b:58028. Zbl 836.54026.
- [4] J. S. Bae and S. K. Yang, $\bar{P} = \bar{R}$ for maps of the circle, Bull. Korean Math. Soc. **24** (1987), 151–157. Zbl 718.54047.
- [5] ———, ω -limit sets for maps of the circle, Bull. Korean Math. Soc. **25** (1988), no. 2, 233–242. MR 90m:58164. Zbl 681.54020.
- [6] Z. Nitecki, *Periodic and limit orbits and the depth of the center for piecewise monotone interval maps*, Proc. Amer. Math. Soc. **80** (1980), no. 3, 511–514. MR 81j:58068. Zbl 478.58019.
- [7] J. C. Xiong, *Nonwandering sets of continuous interval self-maps*, Kexue Tongbao (English Ed.) **29** (1984), no. 11, 1431–1433. MR 86j:58131. Zbl 575.54034.
- [8] ———, *Sets of recurrent points of continuous maps of the interval*, Proc. Amer. Math. Soc. **95** (1985), no. 3, 491–494. MR 87d:58114. Zbl 591.54031.
- [9] ———, *The attracting centre of a continuous self-map of the interval*, Ergodic Theory Dynamical Systems **8** (1988), no. 2, 205–213. MR 90b:58131. Zbl 658.58032.

YEOM AND MIN: DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY OF TECHNOLOGY, NOWON-GU, SEOUL, 139-743, KOREA

CHO: DEPARTMENT OF MATHEMATICS, HANSEO UNIVERSITY, CHUNGNAM, SEOSAN, 356-820, KOREA

