ON THE SHARP CONSTANT FOR STARLIKENESS

CHEN KEYING

(Received 24 February 2000)

ABSTRACT. We obtain a sharp constant of the sufficient condition for p -valently starlikeness, which had been studied by Nunokawa (1991), Obradović and Owa (1989), and Li (1993).

2000 Mathematics Subject Classification. Primary 30C45.

1. Introduction. Let $A(p)$ denote the class of functions of the form

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n \quad (p \in \mathbb{N}) \quad (1.1)$$

which are analytic in $U = \{z : |z| < 1\}$. A function $f(z)$ in $A(p)$ is said to be p -valently starlike if and only if

$$\Re \left[\frac{zf'(z)}{f(z)} \right] > 0 \quad \text{in } U. \quad (1.2)$$

Let $S(p)$ denote the subclass of $A(p)$ consisting of all functions $f(z)$ which are p -valently starlike in U (cf. [1]). For a function $g(z)$ in $A(p)$, the interesting problem is to find the best constant A such that $g(z)$ is in $S(p)$ whenever

$$\left| 1 + \frac{zg^{(p+1)}(z)}{g^{(p)}(z)} \right| < A \left| \frac{zg^{(p)}(z)}{g^{(p-1)}(z)} \right| \quad \text{in } U. \quad (1.3)$$

In 1989, Obradović and Owa [6] obtained that $A = 5/4$ for the case of $p = 1$. For the general case, Nunokawa [5] gained that $A = \log 4$. Recently, Li [2] improved these results and obtained that $A = 3/2$. In this paper, we will solve this problem completely and give the sharp constant $A = 1.80898\dots$, where A is the unique solution of the equation

$$xe^{1/(x^2-1)} = x + 1. \quad (1.4)$$

For proving our result, we should recall the concept of subordination between analytic functions. Given two analytic functions $f(z)$ and $F(z)$, the function $f(z)$ is said to be subordinate to $F(z)$ if $F(z)$ is univalent in U , $f(0) = F(0)$, and $f(U) \subset F(U)$. We denote this subordination by $f(z) \prec F(z)$ (see [7]).

Suppose that $h(z)$ is analytic in U , and that $\Phi(z)$ is analytic in an appropriate domain D , we consider the following first-order differential subordination

$$\beta + zp'(z)\Phi(p(z)) \prec h(z), \quad (1.5)$$

where $p(z)$ is analytic in $\mathbf{U}$, β is a complex constant. Changing the “ $<$ ” of (1.5) to “ $=$ ”, we get the corresponding first-order differential equation

$$\beta + zp'(z)\Phi(p(z)) = h(z). \quad (1.6)$$

2. Main results. Our results rest on the following lemma, which is the special case of [3, Theorem 3].

LEMMA 2.1. *Suppose that $h(z)$ is a starlike function in $\mathbf{U}$, $\Phi(z)$ is analytic in the domain $\mathbf{D}$ and $p(z)$, $q(z)$ are two analytic functions in $\mathbf{U}$. If $p(z)$ satisfies the relation (1.5), $q(z)$ is a univalent solution of the corresponding equation (1.6) and $p(0) = q(0)$, then $p(z) < q(z)$.*

THEOREM 2.2. *Let $g(z) \in A(p)$, and suppose that*

$$\left| 1 + \frac{zg^{(p+1)}(z)}{g^{(p)}(z)} \right| < A \left| \frac{zg^{(p)}(z)}{g^{(p-1)}(z)} \right| \quad \text{in } \mathbf{U}, \quad (2.1)$$

where the constant A is given by (1.4). Then $g(z) \in S(p)$ and the result is sharp.

PROOF. Let

$$f(z) = \frac{g^{(p-1)}(z)}{p!}. \quad (2.2)$$

Then $f(z) \in A(1)$. From the assumption (2.1), $f(z)$ satisfies

$$\left| 1 + \frac{zf''(z)}{f'(z)} \right| < A \left| \frac{zf'(z)}{f(z)} \right| \quad \text{in } \mathbf{U}. \quad (2.3)$$

By putting $p(z) = zf'(z)/f(z)$, equation (2.3) can be rewritten as

$$\left| 1 + \frac{zp'(z)}{p^2(z)} \right| < A. \quad (2.4)$$

Let $\varphi(z) = A(1 + Az)/(A + z)$ for $z \in \mathbf{U}$. Obviously $\varphi(z)$ is a conformal mapping from $\mathbf{U}$ to $\Omega = \{w : |w| < A\}$ and $\varphi(0) = 1$. Combining (2.4) with the definition of subordination, we obtain

$$1 + \frac{zp'(z)}{p^2(z)} < \frac{A(1 + Az)}{A + z}. \quad (2.5)$$

Setting

$$q(z) = \frac{1}{1 + (A^2 - 1)\log A/(A + z)}, \quad (2.6)$$

we have

$$1 + \frac{zq'(z)}{q^2(z)} = \frac{A(1 + Az)}{A + z} \quad (2.7)$$

and $p(0) = q(0) = 1$. As $A > 1$, we can choose a uniform analytic branch of $\log(A + z)$ such that $q(z)$ is univalent on this branch. By taking the real part of the denominator of $q(z)$ and combining (1.4), we conclude that

$$\Re \left[1 + (A^2 - 1)\log \frac{A}{A + z} \right] > 1 + (A^2 - 1)\log \frac{A}{A + 1} = 0. \quad (2.8)$$

It follows that $\Re[q(z)] > 0$, so $q(z)$ is analytic and univalent. Let $\mathbf{D} = \mathbb{C} \setminus \{0\}$, $\Phi(z) = 1/z^2$, $\beta = 1$, and $h(z) = A(1 + Az)/(A + z)$, where $\mathbb{C}$ is the complex plane. It is clear that $h(z)$ is a starlike function. From [Lemma 2.1](#), we deduce that $p(z) \prec q(z)$. Hence

$$\Re \left[\frac{zf'(z)}{f(z)} \right] = \Re[p(z)] \geq \min_{|z|=r<1} \Re[q(z)] > 0. \quad (2.9)$$

This is equivalent to

$$\Re \left[\frac{zg^{(p)}(z)}{g^{(p-1)}(z)} \right] = \Re \left[\frac{zf'(z)}{f(z)} \right] > 0 \quad \text{in } \mathbf{U}. \quad (2.10)$$

From [\[4, Theorem 5\]](#), we have

$$\Re \left[\frac{zg'(z)}{g(z)} \right] > 0 \quad \text{in } \mathbf{U}. \quad (2.11)$$

This proves $g(z) \in S(p)$.

For any $A_1 > A = 1.80898\dots$, we get a function $q_1(z)$ by replacing A in [\(2.6\)](#) with A_1 and choosing an appropriate branch of $\log(A_1 + z)$. We can easily observe that the real part of $q_1(z)$ is not always positive. Through the relations $q_1(z) = zf'(z)/f(z)$ and $f(z) = g^{(p-1)}(z)/p!$, we can construct an analytic function $g(z)$ which belongs to $A(p)$ and satisfies [\(2.1\)](#), but it is not in $S(p)$. This completes the proof. $\square$

Taking $p = 1$ in [Theorem 2.2](#), we easily have the following corollary.

COROLLARY 2.3. *If $f(z) \in A(1)$ and it satisfies the condition*

$$\left| 1 + \frac{zf''(z)}{f'(z)} \right| < A \left| \frac{zf'(z)}{f(z)} \right| \quad \text{in } \mathbf{U}, \quad (2.12)$$

where the constant A is given by [\(1.4\)](#), then $f(z)$ is univalent and starlike in $\mathbf{U}$.

The problem that Nunokawa proposed in [\[5\]](#) has been solved completely, but the converse proposition of [Theorem 2.2](#) is not true. We find a simple example $f(z) = z/(1 - z)$ which belongs to $S(1)$, but it does not satisfy [\(2.12\)](#). The following theorem is better than [\(2.1\)](#) because it includes at least this example.

THEOREM 2.4. *Let $g(z) \in A(p)$, and suppose that*

$$\left| 1 + \frac{zg^{(p+1)}(z)}{g^{(p)}(z)} - \frac{zg^{(p)}(z)}{g^{(p-1)}(z)} \right| < \left| \frac{zg^{(p)}(z)}{g^{(p-1)}(z)} \right| \quad \text{in } \mathbf{U}. \quad (2.13)$$

Then $g(z) \in S(p)$.

PROOF. Let

$$f(z) = \frac{g^{(p-1)}(z)}{p!}. \quad (2.14)$$

Then $f(z) \in A(1)$. From the assumption [\(2.13\)](#), $f(z)$ satisfies

$$\left| 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right| < \left| \frac{zf'(z)}{f(z)} \right| \quad \text{in } \mathbf{U}. \quad (2.15)$$

By setting $p(z) = zf'(z)/f(z)$, equation (2.15) can be rewritten as

$$\left| \frac{zp'(z)}{p^2(z)} \right| < 1. \quad (2.16)$$

From the definition of subordination, we obtain

$$\frac{zp'(z)}{p^2(z)} < z. \quad (2.17)$$

Let $q(z) = 1/(1-z)$, we observe that $zq'(z)/q^2(z) = z$, $p(0) = q(0) = 1$, and $\Re[q(z)] > 0$. From Lemma 2.1, we know that $p(z) < 1/(1-z)$. Therefore

$$\Re \left[\frac{zf'(z)}{f(z)} \right] = \Re[p(z)] \geq \min_{|z|=r<1} \Re[q(z)] > 0. \quad (2.18)$$

This is equivalent to

$$\Re \left[\frac{zg^{(p)}(z)}{g^{(p-1)}(z)} \right] = \Re \left[\frac{zf'(z)}{f(z)} \right] > 0 \quad \text{in } U. \quad (2.19)$$

From [4, Theorem 5], we have

$$\Re \left[\frac{zg'(z)}{g(z)} \right] > 0 \quad \text{in } U. \quad (2.20)$$

This completes the proof. $\square$

Taking $p = 1$ in Theorem 2.4, we obviously have the following corollary.

COROLLARY 2.5. *If $f(z) \in A(1)$ and it satisfies the condition*

$$\left| 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right| < \left| \frac{zf'(z)}{f(z)} \right| \quad \text{in } U, \quad (2.21)$$

then $f(z) \in S(1)$.

ACKNOWLEDGEMENTS. I wish to express my gratitude to Professor Hu Ke and Professor Fang Ainong for their guidance, advice, and encouragement in my work, past and present. I am also grateful to the referee for his valuable advice.

This research was supported by China NSF (Grant No. 19531060) and Doctor Spot Foundation (Grant No. 97024811).

REFERENCES

- [1] A. W. Goodman, *On the Schwarz-Christoffel transformation and p -valent functions*, Trans. Amer. Math. Soc. **68** (1950), 204–223. [MR 11,508d](#). [Zbl 037.05502](#).
- [2] J. L. Li, *On a criterion of starlikeness*, Math. Japon. **38** (1993), no. 5, 897–899. [MR 94j:30012](#). [Zbl 786.30010](#).
- [3] S. S. Miller and P. T. Mocanu, *On some classes of first-order differential subordinations*, Michigan Math. J. **32** (1985), no. 2, 185–195. [MR 86h:30046](#). [Zbl 575.30019](#).
- [4] M. Nunokawa, *On the theory of multivalent functions*, Tsukuba J. Math. **11** (1987), no. 2, 273–286. [MR 89d:30013](#). [Zbl 639.30014](#).

- [5] ———, *On certain multivalent functions*, Math. Japon. **36** (1991), no. 1, 67–70.
[MR 92b:30017](#). [Zbl 718.30010](#).
- [6] M. Obradović and S. Owa, *A criterion for starlikeness*, Math. Nachr. **140** (1989), 97–102.
[MR 90i:30020](#). [Zbl 676.30009](#).
- [7] C. Pommerenke, *Univalent Functions*. With a chapter on quadratic differentials by Gerd Jensen, *Studia Mathematica/Mathematische Lehrbücher*, vol. 25, Vandenhoeck & Ruprecht, Göttingen, 1975. [MR 58#22526](#). [Zbl 298.30014](#).

CHEN KEYING: DEPARTMENT OF APPLIED MATHEMATICS, SHANGHAI JIAOTONG UNIVERSITY,
SHANGHAI 200240, CHINA

E-mail address: kychen801@mail11.sjtu.edu.cn