ROUGH MARCINKIEWICZ INTEGRAL OPERATORS

HUSSAIN AL-QASSEM and AHMAD AL-SALMAN

(Received 16 January 2001)

ABSTRACT. We study the Marcinkiewicz integral operator $M_{\mathcal{P}}f(x) = (\int_{-\infty}^{\infty} |\int_{|y|\leq 2t} f(x - \mathcal{P}(y))(\Omega(y)/|y|^{n-1}) dy|^2 dt/2^{2t})^{1/2}$, where $\mathcal{P}$ is a polynomial mapping from $\mathbb{R}^n$ into $\mathbb{R}^d$ and Ω is a homogeneous function of degree zero on $\mathbb{R}^n$ with mean value zero over the unit sphere S^{n-1} . We prove an L^p boundedness result of $M_{\mathcal{P}}$ for rough Ω .

2000 Mathematics Subject Classification. 42B20, 42B15, 42B25.

1. Introduction. Let $\mathbb{R}^n$, $n \geq 2$ be the n -dimensional Euclidean space and S^{n-1} be the unit sphere in $\mathbb{R}^n$ equipped with the induced Lebesgue measure. Consider the Marcinkiewicz integral operator

$$\mu f(x) = \left(\int_{-\infty}^{\infty} |\mathbf{F}_t(x)|^2 \frac{dt}{2^{2t}} \right)^{1/2}, \quad (1.1)$$

where

$$\mathbf{F}_t(x) = \int_{|x-y|\leq 2t} f(y) \frac{\Omega(x-y)}{|x-y|^{n-1}} dy, \quad (1.2)$$

and Ω is a homogeneous function of degree zero which has the following properties:

$$\Omega \in L^1(S^{n-1}), \quad \int_{S^{n-1}} \Omega(y') d\sigma(y') = 0. \quad (1.3)$$

When $\Omega \in \text{Lip}_{\alpha}(S^{n-1})$, ($0 < \alpha \leq 1$), Stein proved the L^p boundedness of $\mu(f)$ for all $1 < p \leq 2$. Subsequently, Benedek, Calderón, and Panzone proved the L^p boundedness of $\mu(f)$ for all $1 < p < \infty$ under the condition $\Omega \in C^1(S^{n-1})$ (see [2]).

The authors of [3] were able to prove the following result for the more general class of operators

$$\mu_P f(x) = \left(\int_{-\infty}^{\infty} |\mathbf{F}_{P,t}(x)|^2 \frac{dt}{2^{2t}} \right)^{1/2}, \quad (1.4)$$

where

$$\mathbf{F}_{P,t}(x) = \int_{|y|\leq 2t} f(x - P(|y|)y') \frac{\Omega(y)}{|y|^{n-1}} dy \quad (1.5)$$

and P is a real-valued polynomial on $\mathbb{R}$ and satisfies $P(0) = 0$.

THEOREM 1.1 (see [3]). *Let $\alpha > 0$, and $\Omega \in V_{\alpha}(n)$. Then the operator μ_P is bounded in $L^p(\mathbb{R}^n)$ for $(2\alpha + 2)/(2\alpha + 1) < p < 2 + 2\alpha$.*

In [1], Al-Salman and Pan studied the singular integral operator

$$\mathbf{T}_{\Omega, \mathcal{P}} f(x) = \text{p.v.} \int_{\mathbb{R}^n} f(x - \mathcal{P}(y)) \frac{\Omega(y')}{|y|^n} dy, \quad (1.6)$$

where $\mathcal{P} = (P_1, \dots, P_d): \mathbb{R}^n \rightarrow \mathbb{R}^d$ is a polynomial mapping, $d \geq 1$, $n \geq 2$. The authors of [1] proved that $\mathbf{T}_{\Omega, \mathcal{P}}$ is bounded in $L^p(\mathbb{R}^d)$ whenever $(2 + 2\alpha)/(1 + 2\alpha) < p < 2 + 2\alpha$ and $\Omega \in W_\alpha(n)$. Here $W_\alpha(n)$ is a subspace of $L^1(\mathbb{S}^{n-1})$ and its definition as well as the definition of $V_\alpha(n)$ will be reviewed in Section 2. It was shown in [1] that $W_\alpha(n) = V_\alpha(n)$, if $n = 2$ and it is a proper subspace of $V_\alpha(n)$ if $n \geq 3$.

Our purpose in this paper is to study the L^p boundedness of the class of operators

$$M_{\mathcal{P}} f(x) = \left(\int_{-\infty}^{\infty} |\mathbf{F}_{\mathcal{P}, t}(x)|^2 \frac{dt}{2^{2t}} \right)^{1/2}, \quad (1.7)$$

where

$$\mathbf{F}_{\mathcal{P}, t}(x) = \int_{|y| \leq 2^t} f(x - \mathcal{P}(y)) \frac{\Omega(y')}{|y|^{n-1}} dy. \quad (1.8)$$

Our main result in this paper is the following theorem.

THEOREM 1.2. *Let $\alpha > 0$, and $\Omega \in W_\alpha(n)$. Then the operator $M_{\mathcal{P}}$ is bounded in $L^p(\mathbb{R}^d)$ for $(2\alpha + 2)/(2\alpha + 1) < p < 2 + 2\alpha$. The bound of $M_{\mathcal{P}} f$ is independent of the coefficients of $\{P_j\}$.*

By [1, Theorem 3.1] and Theorem 1.2 we have the following corollary.

COROLLARY 1.3. *Let $\alpha > 0$, $\Omega \in V_\alpha(2)$ and $\mathcal{P}: \mathbb{R}^2 \rightarrow \mathbb{R}^d$. Then $M_{\mathcal{P}}$ is bounded in $L^p(\mathbb{R}^d)$ for $(2\alpha + 2)/(2\alpha + 1) < p < 2 + 2\alpha$. The bound of $M_{\mathcal{P}}$ is independent of the coefficients of $\{P_j\}$.*

2. Preparation. We start this section by recalling the following definition from [1].

DEFINITION 2.1. For $\alpha > 0$, $N \geq 1$, let $\tilde{\mathcal{V}}(n, N) = \bigcup_{m=1}^N \mathcal{V}(n, m)$ and let $W_\alpha(N, n)$ be the subspace of $L^1(\mathbb{S}^{n-1})$ defined by

$$W_\alpha(N, n) = \left\{ \Omega \in L^1(\mathbb{S}^{n-1}) : \int_{\mathbb{S}^{n-1}} \Omega(y') d\sigma(y') = 0, M_\alpha(N, n) < \infty \right\}, \quad (2.1)$$

where

$$\begin{aligned} M_\alpha(N, n) &= \max \left\{ \int_{\mathbb{S}^{n-1}} |\Omega(y')| \left(\log \frac{1}{|P(y')|} \right)^{1+\alpha} d\sigma(y') : P \in \tilde{\mathcal{V}}(n, N) \text{ with } \|P\| = 1 \right\}. \end{aligned} \quad (2.2)$$

For $\alpha > 0$, we define $W_\alpha(n)$ to be

$$W_\alpha(n) = \bigcap_{N=1}^{\infty} W_\alpha(N, n). \quad (2.3)$$

Also, for $\alpha > 0$, we define $V_\alpha(n)$ by $V_\alpha(n) = W_\alpha(1, n)$ (see [6]).

Here $\mathcal{V}(n, m)$ is the space of all real-valued homogeneous polynomials on $\mathbb{R}^n$ with degree equal to m and with norm $\|\cdot\|$ defined by

$$\left\| \sum_{|\alpha|=m} a_\alpha \mathbf{y}^\alpha \right\| = \sum_{|\alpha|=m} |a_\alpha|. \quad (2.4)$$

Now we need to recall the following results.

LEMMA 2.2 (see van der Corput [7]). *Suppose ϕ and ψ are real-valued and smooth in (a, b) , and that $|\phi^{(k)}(t)| \geq 1$ for all $t \in (a, b)$. Then the inequality*

$$\left| \int_a^b e^{-i\lambda\phi(t)} \psi(t) dt \right| \leq C_k |\lambda|^{-1/k} \left[|\psi(b)| + \int_a^b |\psi'(t)| dt \right], \quad (2.5)$$

holds when

(i) $k \geq 2$, or

(ii) $k = 1$ and ϕ' is monotonic.

The bound C_k is independent of a, b, ϕ , and λ .

LEMMA 2.3 (see [7]). *Let $\mathcal{P} = (P_1, \dots, P_d)$ be a polynomial mapping from $\mathbb{R}^n$ into $\mathbb{R}^d$. Let $\deg(\mathcal{P}) = \max_{1 \leq j \leq d} \deg(P_j)$. Suppose $\Omega \in L^1(\mathbb{S}^{n-1})$ and*

$$\mu_{\Omega, \mathcal{P}} f(x) = \sup_{h>0} \left| \frac{1}{h^n} \int_{|y|<h} f(x - \mathcal{P}(y)) \Omega(y') dy \right|. \quad (2.6)$$

Then for every $1 < p \leq \infty$, there exists a constant $C_p > 0$ which is independent of Ω and the coefficients of $\{P_j\}$ such that

$$\|\mu_{\Omega, \mathcal{P}} f\|_p \leq C_p \|\Omega\|_{L^1(\mathbb{S}^{n-1})} \|f\|_p \quad (2.7)$$

for every $f \in L^p(\mathbb{R}^d)$.

To each polynomial mapping $\mathcal{P} = (P_1, \dots, P_d) : \mathbb{R}^n \rightarrow \mathbb{R}^d$ with

$$\deg \mathcal{P} = \max_{1 \leq j \leq d} \deg P_j = N, \quad d \geq 1, \quad n \geq 2, \quad (2.8)$$

we define a family of measures

$$\{\mathfrak{g}_t^l, \lambda_t^l : l = 0, 1, \dots, N, \quad t \in \mathbb{R}\} \quad (2.9)$$

as follows.

For $1 \leq j \leq d$, $0 \leq l \leq N$ let $P_j = \sum_{|\alpha| \leq N} C_{j\alpha} \mathbf{y}^\alpha$ and let $Q^l = (Q_1^l, \dots, Q_d^l)$ where $Q_j^l = \sum_{|\alpha| \leq l} C_{j\alpha} \mathbf{y}^\alpha$.

Now for $0 \leq l \leq N$ and $t \in \mathbb{R}$, let $\mathfrak{g}_t^l$ and λ_t^l be the measures defined in the Fourier transform side by

$$\begin{aligned} (\mathfrak{g}_t^l)^\wedge(\xi) &= \int_{|y'| \leq 2^t} e^{-2\pi i \xi \cdot Q^l(y')} \frac{\Omega(y')}{|y'|^{n-1}} \frac{dy'}{2^t}, \\ (\lambda_t^l)^\wedge(\xi) &= \int_{|y'| \leq 2^t} e^{-2\pi i \xi \cdot Q^l(y')} \frac{|\Omega(y')|}{|y'|^{n-1}} \frac{dy'}{2^t}. \end{aligned} \quad (2.10)$$

The maximal functions $(\mathfrak{I}^l)^*$ defined by

$$(\mathfrak{I}^l)^*(f)(x) = \sup_{t \in \mathbb{R}} |\lambda_t^l * f(x)|, \quad (2.11)$$

for $l = 0, 1, \dots, N$.

For later purposes, we need the following definition.

DEFINITION 2.4. For each $1 \leq l \leq N$, let $N_l = |\{\alpha \in \mathbb{N}^n : |\alpha| = l\}|$ and let $\{\alpha \in \mathbb{N}^n : |\alpha| = l\} = \{\alpha_1, \dots, \alpha_{N_l}\}$. For each $1 \leq l \leq N$, define the linear transformations $L_l^{\alpha_j} : \mathbb{R}^d \rightarrow \mathbb{R}$ and $L_l : \mathbb{R}^d \rightarrow \mathbb{R}^{N_l}$ by

$$\begin{aligned} L_l^{\alpha_j}(\xi) &= \sum_{i=1}^d (C_{i, \alpha_j} y^{\alpha_j}) \xi_i, \quad j = 1, \dots, N_l, \\ L_l(\xi) &= (L_l^{\alpha_1}(\xi), \dots, L_l^{\alpha_{N_l}}(\xi)). \end{aligned} \quad (2.12)$$

To simplify the proof of our result we need the following lemma.

LEMMA 2.5. Let $\{\sigma_t^l : l = 0, 1, \dots, N, t \in \mathbb{R}\}$ be a family of measures such that $\sigma_t^0 = 0$ for all $t \in \mathbb{R}$. Let $D_l : \mathbb{R}^n \rightarrow \mathbb{R}^d$, $l = 0, 1, \dots, N$ be linear transformations. Suppose that for all $t \in \mathbb{R}$ and $l = 0, 1, \dots, N$, then

$$\begin{aligned} \|\sigma_t^l\| &\leq C(l), \\ |(\sigma_t^l)\hat{(\xi)}| &\leq C \frac{M_\alpha}{(\log[c2^{lt}|D_l(\xi)|])^{1+\alpha}}, \\ |(\sigma_t^l)\hat{(\xi)} - (\sigma_t^{l-1})\hat{(\xi)}| &\leq C2^{lt}|D_l(\xi)|. \end{aligned} \quad (2.13)$$

Then there exists a family of measures $\{v_t^l : l = 1, \dots, N\}_{t \in \mathbb{R}}$ such that

$$\begin{aligned} \|v_t^l\| &\leq C(l), \\ |(v_t^l)\hat{(\xi)}| &\leq C \frac{M_\alpha}{(\log[c2^{lt}|D_l(\xi)|])^{1+\alpha}}, \\ |(v_t^l)\hat{(\xi)}| &\leq C2^{lt}|D_l(\xi)|, \\ \sigma_t^N &= \sum_{l=1}^N v_t^l. \end{aligned} \quad (2.14)$$

PROOF. By [5, Lemma 6.1], for each $l = 1, \dots, N$ choose two nonsingular linear transformations

$$A_l : \mathbb{R}^{r(l)} \rightarrow \mathbb{R}^d, \quad B_l : \mathbb{R}^d \rightarrow \mathbb{R}^d, \quad (2.15)$$

such that

$$|A_l \pi_{r(l)}^d B_l(\xi)| \leq |D_l(\xi)| \leq N |A_l \pi_{r(l)}^d B_l(\xi)|, \quad \xi \in \mathbb{R}^d, \quad (2.16)$$

where $r(l) = \text{rank}(D_l)$ and $\pi_{r(l)}^d$ is the projection operator from $\mathbb{R}^d$ into $\mathbb{R}^{r(l)}$.

Now choose $\eta \in C_0^\infty(\mathbb{R})$ such that $\eta(t) = 1$ for $|t| \leq 1/2$ and $\eta(t) = 0$ for $|t| \geq 1$. Let $\varphi(t) = \phi(t^2)$ and let

$$\begin{aligned} (v_t^l)^{\wedge}(\xi) &= (\sigma_t^l)^{\wedge}(\xi) \prod_{l < j \leq N} \varphi(|2^{tj} A_j \pi_{r(j)}^d B_j(\xi)|) \\ &\quad - (\sigma_t^{l-1})^{\wedge}(\xi) \prod_{l-1 < j \leq N} \varphi(|2^{tj} A_j \pi_{r(j)}^d B_j(\xi)|) \end{aligned} \quad (2.17)$$

with the convention $\prod_{j \in \emptyset} a_j = 1$, $1 \leq l \leq N$.

Hence, one can easily see that $\{\sigma_t^l : l = 1, \dots, N, t \in \mathbb{R}\}$ is the desired family of measures. $\square$

Now for the boundedness of the maximal functions $(\mathfrak{g}^l)^*$, $l = 0, 1, \dots, N$, we have the following lemma whose proof is an easy consequence of [Lemma 2.3](#), polar coordinates and Hölder's inequality:

LEMMA 2.6. *For $l = 1, \dots, N$ and $p \in (1, \infty)$, there exists a constant $C_{p,l}$ which is independent of the coefficients of the polynomial components of the mapping Q^l such that*

$$\|(\mathfrak{g}^l)^* f\|_p \leq C_{p,l} \|f\|_p. \quad (2.18)$$

3. Boundedness of some square functions. For a nonnegative C^∞ radial function Φ on $\mathbb{R}^n$ with

$$\text{supp}(\Phi) \subset \left\{x \in \mathbb{R}^n : \frac{1}{2} \leq |x| \leq 2\right\}, \quad \int_0^\infty \frac{\Phi(t)}{t} dt = 1, \quad (3.1)$$

and for a linear transformation $\mathbf{L} : \mathbb{R}^n \rightarrow \mathbb{R}^d$, define the functions ψ_t , $t \in \mathbb{R}$ by $\hat{\psi}_t(y) = \Phi(2^t \mathbf{L}(y))$.

For a family of measures $\{\sigma_t\}_{t \in \mathbb{R}}$, real number u and $l \in \mathbb{N}$, let $\mathbf{J}_u^l(f)$ be the square function defined by

$$\mathbf{J}_u^l(f)(x) = \left(\int_{-\infty}^\infty |\sigma_t * \psi_{l(t+u)} * f(x)|^2 dt \right)^{1/2}. \quad (3.2)$$

For such a square function we have the following theorem.

THEOREM 3.1. *If $\{\sigma_t\}_{t \in \mathbb{R}}$ is a family of measures such that the corresponding maximal function*

$$\sigma^*(f)(x) = \sup_{t \in \mathbb{R}} |\sigma_t * f(x)| \quad (3.3)$$

is bounded on $L^p(\mathbb{R}^d)$ for every $1 < p < \infty$, then

$$\|\mathbf{J}_u^l(f)\|_{L^p(\mathbb{R}^d)} \leq C_{p,l} \sqrt{\|\sigma^*\|_{(p/2)'} \sup_{t \in \mathbb{R}} \|\sigma_t\|} \|f\|_{L^p(\mathbb{R}^d)} \quad (3.4)$$

for every $1 < p < \infty$. Here $C_{p,l}$ is a constant that depends only on p and the dimension of the underlying space.

PROOF. If $\sup_{t \in \mathbb{R}} \|\sigma_t\| = \infty$, then the inequality holds trivially. Thus we may assume that $\sup_{t \in \mathbb{R}} \|\sigma_t\| < \infty$. In this case we follow a similar argument as in [4]. Let $p > 2$ and $q = (p/2)'$. Choose a nonnegative function $v \in L_+^q$ with $\|v\|_q = 1$ such that

$$\|\mathbf{J}_u^l(f)\|_p^2 = \int_{\mathbb{R}^d} \left(\int_{-\infty}^{\infty} |\sigma_t * \psi_{l(t+u)} * f(x)|^2 dt \right) v(x) dx. \quad (3.5)$$

Thus it is easy to see that

$$\begin{aligned} \|\mathbf{J}_u^l(f)\|_p^2 &\leq \sup_{t \in \mathbb{R}} \|\sigma_t\| \int_{-\infty}^{\infty} \int_{\mathbb{R}^d} |\psi_{l(t+u)} * f(z)|^2 \sigma^*(v)(-z) dz dt \\ &\leq \sup_{t \in \mathbb{R}} \|\sigma_t\| \int_{\mathbb{R}^d} [g(f)]^2(z) \sigma^*(v)(-z) dz, \end{aligned} \quad (3.6)$$

where

$$g(f)(x) = \left(\int_{-\infty}^{\infty} |\psi_{l(t+u)} * f(x)|^2 dt \right)^{1/2}. \quad (3.7)$$

Now since $\int_{\mathbb{R}^d} \psi_t(x) dx = 0$, it is well known that

$$\|g(f)\|_p \leq C_p \|f\|_p \quad \forall 1 < p < \infty \quad (3.8)$$

with constant C_p that depends only on p and the dimension of the underlying space.

Thus by (3.6) and Hölder's inequality we have

$$\begin{aligned} \|\mathbf{J}_u^l(f)\|_p^2 &\leq \sup_{t \in \mathbb{R}} \|\sigma_t\| \|g(f)\|_p^2 \|\sigma^*(u)\|_q \\ &\leq C_p^2 \sup_{t \in \mathbb{R}} \|\sigma_t\| \|\sigma^*\|_{(p/2)'} \|f\|_p^2. \end{aligned} \quad (3.9)$$

Hence our result follows by taking the square root on both sides. The case $p < 2$ follows by duality. $\square$

4. Proof of the main theorem. Let $\alpha > 0$, $\Omega \in W_\alpha(n)$. Let $\mathcal{P} = (P_1, \dots, P_d)$ be a polynomial mapping from $\mathbb{R}^n$ into $\mathbb{R}^d$ with $\deg \mathcal{P} = \max_{1 \leq j \leq d} \deg P_j = N$, where $d \geq 1$ and $n \geq 2$. For $0 \leq l \leq N$ let N_l , Q^l , ν_t^l , λ_t^l , and L_l be as in Section 3.

The first step in our proof is to show that each $\mathcal{G}_t^l$, $l = 1, \dots, N$ satisfies the hypotheses of Lemma 2.5, that is,

$$\|\mathcal{G}_t^l\| \leq C(l), \quad (4.1)$$

$$|(\mathcal{G}_t^l)^\wedge(\xi)| \leq C \frac{M_\alpha}{(\log[c2^{lt} |L_l(\xi)|])^{1+\alpha}}, \quad (4.2)$$

$$|(\mathcal{G}_t^l)^\wedge(\xi) - (\mathcal{G}_t^{l-1})^\wedge(\xi)| \leq C2^{lt} |L_l(\xi)|. \quad (4.3)$$

One can easily see that (4.1) holds trivially. Using the cancellation property of Ω , it is easy to see that (4.3) holds. Thus, we need only to verify (4.2). To see that, we notice that

$$|(\mathcal{G}_t^l)^\wedge(\xi)| \leq \int_{\mathbb{S}^{n-1}} |\Omega(y')| \left| \int_0^1 e^{-2\pi i \xi \cdot Q^l(2^l r y')} dr \right| d\sigma(y'). \quad (4.4)$$

Now the quantity $\xi \cdot Q^l(2^{tl}r\gamma')$ can be written in the form

$$\xi \cdot Q^l(2^{tl}r\gamma') = 2^{tl}r^l \lambda G^l(\gamma') + \xi \cdot R(2^t r \gamma'), \quad (4.5)$$

where Q^l is a homogeneous polynomial of degree l with $\|G^l\| = 1$, R is a polynomial of degree at most $l-1$ in the variable r ,

$$\lambda = \sum_{j=1}^{N_l} |L_l^{\alpha_j}(\xi)| \geq N_l |L_l(\xi)| \quad (4.6)$$

and $\alpha_1, \dots, \alpha_{N_l}$ are the constants that appeared in [Section 2](#). Thus by van der Corput lemma, we have

$$\left| \int_0^1 e^{-2\pi i \xi \cdot Q^l(2^t r \gamma')} dr \right| \leq C \min \left\{ 1, (2^{tl} |L_l(\xi)| |G^l(\gamma')|)^{-1/l} \right\} \quad (4.7)$$

and hence

$$\left| \int_0^1 e^{-2\pi i \xi \cdot Q^l(2^t r \gamma')} dr \right| \leq C \frac{[\log(c |G^l(\gamma')|^{-1})]^{1+\alpha}}{(\log[c 2^{tl} |L_l(\xi)|])^{1+\alpha}}, \quad (4.8)$$

where C is a constant independent of t and ξ . Since $\Omega \in W_\alpha(n)$, the estimate [\(4.2\)](#) follows.

By [Lemma 2.5](#), there exists a family of measures $\{\nu_t^l : l = 1, \dots, N, t \in \mathbb{R}\}$ such that

$$\|\nu_t^l\| \leq C(l), \quad (4.9)$$

$$|(\nu_t^l)^\wedge(\xi)| \leq C \frac{M_\alpha}{(\log[c 2^{lt} |L_l(\xi)|])^{1+\alpha}}, \quad (4.10)$$

$$|(\nu_t^l)^\wedge(\xi)| \leq C 2^{lt} |L_l(\xi)|, \quad (4.11)$$

$$\mathfrak{g}_t^N = \sum_{l=1}^N \nu_t^l. \quad (4.12)$$

Also by [Lemma 2.6](#) and the definition of ν_t^l (see the proof of [Lemma 2.5](#)), we have

$$\|(\nu^l)^* f\|_p \leq C_{p,l} \|f\|_p \quad \forall 1 < p < \infty. \quad (4.13)$$

Now one can easily see that

$$2^{-t} \mathbf{F}_{\mathcal{P},t}(x) = \mathfrak{g}_t^N * f(x) = \sum_{l=1}^N \nu_t^l * f(x). \quad (4.14)$$

Therefore,

$$\|M_{\mathcal{P}} f\|_p \leq \sum_{l=1}^N \|M_{\mathcal{P}}^l f\|_p, \quad (4.15)$$

where

$$M_{\mathcal{P}}^l f(x) = \left(\int_{-\infty}^{\infty} |\nu_t^l * f(x)|^2 dt \right)^{1/2}. \quad (4.16)$$

Thus to show the boundedness of $M_\Phi f$, it suffices to show that

$$\|M_\Phi^l f\|_p \leq C_{p,l} \|f\|_p \quad (4.17)$$

for $p \in ((2+2\alpha)/(1+2\alpha), 2+2\alpha)$, and for all $l = 1, \dots, N$.

To show (4.17), we proceed as follows: let Φ and ψ_t be as in Section 3. Then

$$\begin{aligned} M_\Phi^l f(x) &= \log 2^l \left(\int_{-\infty}^{\infty} \left| \int_{-\infty}^{\infty} v_t^l * \psi_{l(t+u)} * f(x) du \right|^2 dt \right)^{1/2} \\ &\leq \log 2^l \int_{-\infty}^{\infty} S_u^l f(x) du, \end{aligned} \quad (4.18)$$

where

$$S_u^l f(x) = \left(\int_{-\infty}^{\infty} |v_t^l * \psi_{l(t+u)} * f(x)|^2 dt \right)^{1/2}. \quad (4.19)$$

Now by (4.13) and Theorem 3.1, we have

$$\|S_u^l f\|_p \leq C_p \|f\|_p \quad (4.20)$$

for all $p \in (1, \infty)$ and for $l = 1, \dots, N$ which in turn implies that

$$\int_{-1}^1 \|S_u^l f\|_p du \leq 2C_p \|f\|_p \quad \forall p \in (1, \infty). \quad (4.21)$$

On the other hand, if $u \geq 1$, by the estimate (4.11) we have

$$\begin{aligned} \|S_u^l f\|_2^2 &= \int_{\mathbb{R}^d} \int_{-\infty}^{\infty} |v_t^l * \psi_{l(t+u)} * f(x)|^2 dt dx \\ &= \int_{-\infty}^{\infty} \int_{\mathbb{R}^d} (\Phi(2^{lt+lu} \mathbf{L}_l(\xi)))^2 |(\hat{v}_t^l)(\xi)|^2 |\hat{f}(\xi)|^2 d\xi dt \\ &\leq 2^{2l-2lu} \int_{\mathbb{R}^d} |\hat{f}(\xi)|^2 \left(\int_{\log(1/2^l |\mathbf{L}_l(\xi)|) - u}^{\log(2^l/|\mathbf{L}_l(\xi)|) - u} dt \right) d\xi \\ &= 2 \log 2^l 2^{2l-2lu} \|f\|_2^2. \end{aligned} \quad (4.22)$$

Thus

$$\|S_u^l f\|_2 \leq \sqrt{2 \log 2^l} 2^{l-lu} \|f\|_2. \quad (4.23)$$

By interpolating between (4.20) and (4.23) we get

$$\|S_u^l f\|_p \leq C_{p,l} 2^{\theta l - \theta l u} \|f\|_p \quad (4.24)$$

for all $1 < p < \infty$ and for some $\theta = \theta(p) > 0$. Hence we have

$$\int_1^\infty \|S_u^l f\|_p du \leq C_p \|f\|_p \quad \text{for } p \in (1, \infty). \quad (4.25)$$

Finally, if $u < -1$, by the estimate (4.10) and similar argument as in the case of $u \leq 1$, we get

$$\|S_u^l f\|_2 \leq C_l(|u|)^{-1-\alpha} \|f\|_2. \quad (4.26)$$

By interpolating between (4.26) and any $p \in (1, \infty)$ in (4.20), we get that, if $p \in ((2+2\alpha)/(1+2\alpha), 2+2\alpha)$ there exists $\beta > 0$ such that

$$\|S_u^l f\|_p \leq C_p(|u|)^{-\beta} \|f\|_p, \quad (4.27)$$

which implies that

$$\int_{-\infty}^{-1} \|S_u^l f\|_p du \leq C_p \|f\|_p \quad (4.28)$$

for $p \in ((2+2\alpha)/(1+2\alpha), 2+2\alpha)$.

Hence by combining (4.18), (4.21), (4.25), and (4.28) we get (4.17). $\square$

ACKNOWLEDGEMENT. The publication of this paper was supported by Yarmouk University Research Council.

REFERENCES

- [1] A. Al-Salman and Y. Pan, *Singular integrals with rough kernels*, to appear in *Canad. Math. Bull.*
- [2] A. Benedek, A. P. Calderón, and R. Panzone, *Convolution operators on Banach space valued functions*, *Proc. Nat. Acad. Sci. U.S.A.* **48** (1962), 356–365. [MR 24#A3479](#). [Zbl 103.33402](#).
- [3] J. Chen, D. Fan, and Y. Pan, *A note on a Marcinkiewics integral operator*, to appear in *Math. Nachr.*
- [4] J. Duoandikoetxea and J. L. Rubio de Francia, *Maximal and singular integral operators via Fourier transform estimates*, *Invent. Math.* **84** (1986), no. 3, 541–561. [MR 87f:42046](#). [Zbl 568.42012](#).
- [5] D. Fan and Y. Pan, *Singular integral operators with rough kernels supported by subvarieties*, *Amer. J. Math.* **119** (1997), no. 4, 799–839. [MR 99c:42029](#). [Zbl 0899.42002](#).
- [6] L. Grafakos and A. Stefanov, *L^p bounds for singular integrals and maximal singular integrals with rough kernels*, *Indiana Univ. Math. J.* **47** (1998), no. 2, 455–469. [MR 99i:42019](#). [Zbl 913.42014](#).
- [7] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Mathematical Series, vol. 43, Princeton University Press, New Jersey, 1993. [MR 95c:42002](#). [Zbl 821.42001](#).

HUSSAIN AL-QASSEM: DEPARTMENT OF MATHEMATICS, YARMOUK UNIVERSITY, IRBID, JORDAN
E-mail address: husseink@yu.edu.jo

AHMAD AL-SALMAN: DEPARTMENT OF MATHEMATICS, YARMOUK UNIVERSITY, IRBID, JORDAN
E-mail address: alsalman@yu.edu.jo