PERIODIC SOLUTIONS OF PERIODIC GENERALIZED FOOD LIMITED MODEL

YONGKUN LI

(Received 14 December 1999)

ABSTRACT. By using the continuation theorem of coincidence degree theory, the existence of positive periodic solutions for a periodic generalized food limited model with state dependent delays and distributed delays is studied, respectively.

2000 Mathematics Subject Classification. Primary 34K13.

1. Introduction. Gopalsamy, Kulenović, and Ladas [4] studied the existence of periodic solutions of the following periodic food limited model:

$$\frac{dN(t)}{dt} = r(t)N(t) \left[\frac{K(t) - N(t - m\omega)}{K(t) + c(t)r(t)N(t - m\omega)} \right],\tag{1.1}$$

where K, r, c are continuous positive ω -periodic functions, m is a positive integer. Also, since as it was indicated by Freedman and Wu [1] that it would be of interesting to study the existence of periodic solutions for population models with periodic delays; and since at present, there are only a few papers which have been published on the existence of periodic solutions of state dependent delay differential equations (see [6] and references therein). Our main purpose in this paper is—by using the Mawhin's continuation theorem of coincidence degree theory [2, 10]—to establish the existence of positive periodic solutions for the periodic generalized food limited model

$$\frac{dN(t)}{dt} = r(t)N(t) \left[\frac{K(t) - N(t) - \sum_{j=1}^{n} a_j(t)N(t - \tau_j(t, N(t)))}{1 + r(t) \sum_{j=1}^{n} b_j(t)N(t - \sigma_j(t, N(t)))} \right]^{\theta}$$
(1.2)

with state dependent delays and

$$\frac{dN(t)}{dt} = r(t)N(t) \left[\frac{K(t) - N(t) - \sum_{j=1}^{n} a_j(t) \int_{-l_j}^{0} N(t+s) d\eta_j(s)}{1 + r(t) \sum_{j=1}^{n} b_j(t) \int_{-h_j}^{0} N(t+s) d\mu_j(s)} \right]^{\theta}$$
(1.3)

with distributed delays, where K > 0, r > 0, $a_j \ge 0$, $b_j \ge 0$ (j = 1, 2, ..., n) are continuous ω -periodic functions, $\tau_j, \sigma_j \in C(R^2, R)$ (j = 1, 2, ..., n) are ω -periodic with respect to their first arguments, respectively, l_j, h_j (j = 1, 2, ..., n) are positive constants, η_j, μ_j (j = 1, 2, ..., n) are nondecreasing with respect to their arguments, respectively, and θ is a positive odd number. As a special case of (1.2), we also obtain

YONGKUN LI

the existence of periodic solutions of the following generalized food limited model with periodic delays

$$\frac{dN(t)}{dt} = r(t)N(t) \left[\frac{K(t) - N(t) - \sum_{j=1}^{n} a_j(t)N(t - \tau_j(t))}{1 + r(t) \sum_{j=1}^{n} b_j(t)N(t - \sigma_j(t))} \right]^{\theta},$$
(1.4)

where K > 0, r > 0, $a_j \ge 0$, $b_j \ge 0$, τ_j , σ_j (j = 1, 2, ..., n) are continuous ω -periodic functions, and θ is a positive odd number. For the ecological sense of (1.4), we refer to [3] and the references therein. For some work concerning the existence of periodic solutions of functional differential equations which was done by using the Mawhin's coincidence degree theory we refer to [5, 7, 8, 9].

2. Main results. In this section, by using the Mawhin continuation theorem we show the existence of at least one positive periodic solution of (1.2) and (1.3). To do so, we first make some preparations.

Let X, Y be real Banach spaces, $L : Dom L \subset X \to Y$ a Fredholm mapping of index zero, and $P : X \to X$, $Q : Y \to Y$ continuous projectors such that Im P = Ker L, Ker Q = Im L, and $X = Ker L \bigoplus Ker P$, $Y = Im L \bigoplus Im Q$. Denote by L_P the restriction of L to $Dom L \cap Ker P$, $K_P : Im L \to Ker P \cap Dom L$ the inverse (to L_P), and $J : Im Q \to Ker L$ an isomorphism of Im Q onto Ker L.

For convenience, we introduce the continuation theorem [2, page 40] as follows.

LEMMA 2.1. Let $\Omega \subset X$ be an open bounded set and $N : X \to Y$ be a continuous operator which is L-compact on $\overline{\Omega}$ (i.e., $QN : \overline{\Omega} \to Y$ and $K_P(I-Q)N : \overline{\Omega} \to Y$ are compact). Assume

(i) for each $\lambda \in (0,1)$, $x \in \partial \Omega \cap \text{Dom}L$, $Lx \neq \lambda Nx$;

(ii) for each $x \in \partial \Omega \cap \text{Ker } L$, $QNx \neq 0$, and $\text{deg}\{JQN, \Omega \cap \text{Ker } L, 0\} \neq 0$. Then Lx = Nx has at least one solution in $\overline{\Omega} \cap \text{Dom } L$.

In what follows, we use the notation

$$\bar{u} = \frac{1}{\omega} \int_0^\omega u(t) \, dt, \qquad |u|_0 = \max_{t \in [0,\omega]} |u(t)|, \tag{2.1}$$

where u is a continuous ω -periodic function.

We are now in a position to state and prove our first main result.

THEOREM 2.2. Equation (1.2) has at least one positive ω -periodic solution.

PROOF. Consider the following equation

$$\frac{dx(t)}{dt} = r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_j(t) \exp\left(x(t - \tau_j(t, e^{x(t)}))\right)}{1 + r(t) \sum_{j=1}^{n} b_j(t) \exp\left(x(t - \sigma_j(t, e^{x(t)}))\right)} \right]^{\theta},$$
(2.2)

where $K, r, a_j, b_j, \tau_j, \sigma_j$ (j = 1, = 2, ..., n) and θ are the same as those in (1.2). It is easy to see that if (2.2) has an ω -periodic solution $x^*(t)$, then $N^*(t) = \exp(x^*(t))$ is a positive ω -periodic solution of (1.2). So, to complete the proof, it suffices to show that (2.2) has an ω -periodic solution. In order to use the continuation theorem of coincidence degree theory to establish the existence of ω -periodic solution of (2.2), we take $X = Y = \{x(t) \in C(R,R) : x(t+\omega) = x(t)\}$. With the norm $|\cdot|_0$, X is a Banach space. Set

$$L: \text{Dom}L\bigcap X, \qquad Lx = \frac{dx}{dt},$$
 (2.3)

where $\text{Dom} L = \{x(t) \in C^1(R, R)\}$, and

$$N: X \longrightarrow X, \qquad Nx = r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_j(t) \exp\left(x(t - \tau_j(t, e^{x(t)}))\right)}{1 + r(t) \sum_{j=1}^{n} b_j(t) \exp\left(x(t - \sigma_j(t, e^{x(t)}))\right)} \right]^{\theta}.$$
 (2.4)

Define two projectors P and Q as

$$Px = Qx = \frac{1}{\omega} \int_0^\omega x(t) dt, \quad x \in X.$$
(2.5)

Clearly, Ker L = R, Im $L = \{x \in X : \int_0^{\omega} x(t) dt = 0\}$ is closed in X and dim Ker L = codim Im L = 1. Hence, L is a Fredholm mapping of index zero. Furthermore, through an easy computation, we find that the inverse K_P of L_P has the form

$$K_P : \operatorname{Im} L \longrightarrow \operatorname{Dom} L \big(\operatorname{Ner} P,$$

$$K_P(y) = \int_0^t x(s) \, ds - \frac{1}{\omega} \int_0^\omega \int_0^u x(s) \, ds \, du, \quad t \in [0, \omega].$$
(2.6)

Notice that

$$QN: X \to X,$$

$$QNx = \frac{1}{\omega} \int_0^{\omega} \left\{ r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^n a_j(t) \exp\left(x(t - \tau_j(t, e^{x(t)}))\right)}{1 + r(t) \sum_{j=1}^n b_j(t) \exp\left(x(t - \sigma_j(t, e^{x(t)}))\right)} \right]^{\theta} \right\} dt.$$
(2.7)

We find

$$K_{P}(I-Q)N: X \to X,$$

$$x \mapsto \int_{0}^{t} \left\{ r(s) \left[\frac{K(s) - e^{x(s)} - \sum_{j=1}^{n} a_{j}(s) \exp\left(x\left(s - \tau_{j}(s, e^{x(s)})\right)\right)}{1 + r(s) \sum_{j=1}^{n} b_{j}(s) \exp\left(x\left(s - \sigma_{j}(s, e^{x(s)})\right)\right)} \right]^{\theta} \right\} ds$$

$$- \frac{1}{\omega} \int_{0}^{\omega} \int_{0}^{u} \left\{ r(s) \left[\frac{K(s) - e^{x(s)} - \sum_{j=1}^{n} a_{j}(s) \exp\left(x\left(s - \tau_{j}(s, e^{x(s)})\right)\right)}{1 + r(s) \sum_{j=1}^{n} b_{j}(s) \exp\left(x\left(s - \sigma_{j}(s, e^{x(s)})\right)\right)} \right]^{\theta} \right\} ds du$$

$$- \left(\frac{t}{\omega} - \frac{1}{2} \right) \int_{0}^{\omega} \left\{ r(s) \left[\frac{K(s) - e^{x(s)} - \sum_{j=1}^{n} a_{j}(s) \exp\left(x\left(s - \tau_{j}(s, e^{x(s)})\right)\right)}{1 + r(s) \sum_{j=1}^{n} b_{j}(s) \exp\left(x\left(s - \sigma_{j}(s, e^{x(s)})\right)\right)} \right]^{\theta} \right\} ds.$$
(2.8)

Obviously, QN and $K_P(I - Q)N$ are continuous by the Lebesgue theorem and $QN(\bar{\Omega})$, $K_P(I - Q)N(\bar{\Omega})$ are relatively compact for any open bounded set $\Omega \subset X$.

YONGKUN LI

Therefore, *N* is *L*-compact on $\overline{\Omega}$ for any open bounded set $\Omega \subset X$. Corresponding to the operator equation $Lx = \lambda Nx$, $\lambda \in (0,1)$, we have

$$\frac{dx(t)}{dt} = \lambda \left\{ r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_j(t) \exp\left(x(t - \tau_j(t, e^{x(t)}))\right)}{1 + r(t) \sum_{j=1}^{n} b_j(t) \exp\left(x(t - \sigma_j(t, e^{x(t)}))\right)} \right]^{\theta} \right\}, \quad \lambda \in (0, 1).$$
(2.9)

Suppose that $x(t) \in X$ is a solution of (2.9) for some $\lambda \in (0,1)$. Choose $t_1 \in [0, \omega]$ such that

$$x(t_1) = \max_{t \in [0,\omega]} x(t).$$
 (2.10)

Then it is clear that

$$x'(t_1) = 0. (2.11)$$

In view of this and (2.9), we obtain

$$r(t_1)\left\{\left[\frac{K(t_1) - e^{x(t_1)} - \sum_{j=1}^n a_j(t_1) \exp\left(x(t_1 - \tau_j(t_1, e^{x(t_1)}))\right)}{1 + r(t_1) \sum_{j=1}^n b_j(t_1) \exp\left(x(t_1 - \sigma_j(t_1, e^{x(t_1)}))\right)}\right]^{\theta}\right\} = 0$$
(2.12)

which implies that

$$K(t_1) - e^{x(t_1)} - \sum_{j=1}^n a_j(t_1) \exp\left(x(t_1 - \tau_j(t_1, e^{x(t_1)}))\right) = 0.$$
(2.13)

Thus

$$e^{x(t_1)} < K(t_1) \le |K|_0, \quad x(t_1) < \ln |K|_0.$$
 (2.14)

Integrating (2.9) over $[0, \omega]$, we obtain

$$\int_{0}^{\omega} \left\{ r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_{j}(t) \exp\left(x\left(t - \tau_{j}(t, e^{x(t)})\right)\right)}{1 + r(t) \sum_{j=1}^{n} b_{j}(t) \exp\left(x\left(t - \sigma_{j}(t, e^{x(t)})\right)\right)} \right]^{\theta} \right\} dt = 0$$
(2.15)

which implies that there exist a positive constant A_1 and a point $t_0 \in [0, \omega]$ such that

$$x(t_0) > -A_1.$$
 (2.16)

It follows from (2.9) and (2.14) that

$$\int_{0}^{\omega} \left| \frac{dx(t)}{dt} \right| dt = \int_{0}^{\omega} \left| r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_{j}(t) \exp\left(x\left(t - \tau_{j}(t, e^{x(t)})\right)\right)}{1 + r(t) \sum_{j=1}^{n} b_{j}(t) \exp\left(x\left(t - \sigma_{j}(t, e^{x(t)})\right)\right)} \right]^{\theta} \right| dt$$

$$< \int_{0}^{\omega} \left| r(t) \left[K(t) - e^{x(t)} - \sum_{j=1}^{n} a_{j}(t) \exp\left(x\left(t - \tau_{j}(t, e^{x(t)})\right)\right) \right]^{\theta} \right| dt$$

$$< \omega \tilde{r} (|K|_{0})^{\theta} \left[2 + \sum_{j=1}^{n} |a_{j}|_{0} \right]^{\theta} \stackrel{\text{def}}{=} A_{2}.$$
(2.17)

By this and (2.16), we obtain

$$x(t) = x(t_0) + \int_{t_0}^t x'(t) dt \ge x(t_0) - \int_0^\omega |x'(t)| dt > -(A_1 + A_2).$$
(2.18)

Denote by $A = \max\{|\ln |K|_0|, A_1 + A_2\} + A_3$, here $A_3 > 0$ is taken sufficiently large such that $K(t) - e^A - e^A \sum_{j=1}^n a_j(t) < 0$ and $K(t) - e^{-A} - e^{-A} \sum_{j=1}^n a_j(t) > 0$ for $t \in [0, \omega]$ and take $\Omega = \{x(t) \in X : |x|_0 < A\}$, then it is clear that Ω satisfies condition (i) in Lemma 2.1. When $x \in \partial \Omega \cap R$, x is a constant with |x| = A. Hence,

$$QNx = \frac{1}{\omega} \int_0^{\omega} \left\{ r(t) \left[\frac{K(t) - e^x - e^x \sum_{j=1}^n a_j(t)}{1 + r(t)e^x \sum_{j=1}^n b_j(t)} \right]^{\theta} \right\} dt \neq 0.$$
(2.19)

Furthermore, take $J = I : \text{Im } Q \to \text{Ker } L$, $x \mapsto x$, by a straightforward computation, we find

$$\deg[JQN, \operatorname{Ker} L \cap \Omega, 0] \neq 0. \tag{2.20}$$

According to Lemma 2.1, we have completed the proof.

Next, immediately, from Theorem 2.2, we have the following corollary.

COROLLARY 2.3. Equation (1.4) has at least one positive ω -periodic solution.

REMARK 2.4. In (1.4), when $b_j(t) = 0$ for all $t \in [0, \omega]$ and j = 1, 2, ..., n, equation (1.4) becomes the well-known logistic equation with several delays.

Finally, similar to the proof of Theorem 2.2, one can prove the following theorem.

THEOREM 2.5. Equation (1.3) has at least one positive ω -periodic solution.

PROOF. The proof is similar to the one of Theorem 2.2, to complete the proof, it suffices to show that the following equation:

$$\frac{dx(t)}{dt} = r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_j(t) \int_{-l_j}^{0} e^{x(t+s)} d\eta_j(s)}{1 + r(t) \sum_{j=1}^{n} b_j(t) \int_{-h_j}^{0} e^{x(t+s)} d\mu_j(s)} \right]^{\theta},$$
(2.21)

has an ω -periodic solution, where $r, K, \theta, a_j, b_j, l_j, h_j, \eta_j, \mu_j, j = 1, 2, ..., n$ are the same as those in (1.3). To this end, set

$$Nx = r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_j(t) \int_{-l_j}^{0} e^{x(t+s)} d\eta_j(s)}{1 + r(t) \sum_{j=1}^{n} b_j(t) \int_{-h_j}^{0} e^{x(t+s)} d\mu_j(s)} \right]^{\theta}$$
(2.22)

and *L*,*P*,*Q*,*X* are the same as those in the proof of Theorem 2.2. Corresponding to the operator equation $Lx = \lambda x$, $\lambda \in (0, 1)$, we have

$$\frac{dx(t)}{dt} = \lambda \left\{ r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_j(t) \int_{-l_j}^{0} e^{x(t+s)} d\eta_j(s)}{1 + r(t) \sum_{j=1}^{n} b_j(t) \int_{-h_j}^{0} e^{x(t+s)} d\mu_j(s)} \right]^{\theta} \right\}, \quad \lambda \in [0, \omega].$$
(2.23)

Suppose that $x(t) \in X$ is a solution of (2.23) for a certain $\lambda \in (0, 1)$. Choose $t^1 \in [0, \omega]$ such that

$$x(t^1) = \max_{t \in [0,\omega]} x(t).$$
 (2.24)

YONGKUN LI

Then

$$x'(t^1) = 0. (2.25)$$

By this and (2.23), we have

$$r(t^{1})\left[\frac{K(t^{1}) - e^{x(t^{1})} - \sum_{j=1}^{n} a_{j}(t^{1}) \int_{-l_{j}}^{0} e^{x(t^{1}+s)} d\eta_{j}(s)}{1 + r(t^{1}) \sum_{j=1}^{n} b_{j}(t^{1}) \int_{-h_{j}}^{0} e^{x(t^{1}+s)} d\mu_{j}(s)}\right]^{\theta} = 0,$$
(2.26)

which implies that

$$K(t^{1}) - e^{x(t^{1})} - \sum_{j=1}^{n} a_{j}(t^{1}) \int_{-l_{j}}^{0} e^{x(t^{1}+s)} d\eta_{j}(s) = 0.$$
(2.27)

Therefore,

$$e^{x(t^1)} < K(t^1) \le |K|_0, \quad x(t^1) < \ln |K|_0.$$
 (2.28)

Integrating (2.23) over $[0, \omega]$, we have

$$\int_{0}^{\omega} \left\{ r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_{j}(t) \int_{-l_{j}}^{0} e^{x(t+s)} d\eta_{j}(s)}{1 + r(t) \sum_{j=1}^{n} b_{j}(t) \int_{-h_{j}}^{0} e^{x(t+s)} d\mu_{j}(s)} \right]^{\theta} \right\} dt = 0,$$
(2.29)

which implies that there exist a positive constant B_1 and a point $t^0 \in [0, \omega]$ such that

$$x(t^0) > -B_1. (2.30)$$

In view of (2.23) and (2.28), we obtain

$$\int_{0}^{\omega} \left| \frac{dx(t)}{dt} \right| dt = \int_{0}^{\omega} \left| r(t) \left[\frac{K(t) - e^{x(t)} - \sum_{j=1}^{n} a_{j}(t) \int_{-l_{j}}^{0} e^{x(t+s)} d\eta_{j}(s)}{1 + r(t) \sum_{j=1}^{n} b_{j}(t) \int_{-h_{j}}^{0} e^{x(t+s)} d\mu_{j}(s)} \right]^{\theta} \right| dt$$

$$< \int_{0}^{\omega} \left| r(t) \left[K(t) - e^{x(t)} - \sum_{j=1}^{n} a_{j}(t) \int_{-l_{j}}^{0} e^{x(t+s)} d\eta_{j}(s) \right]^{\theta} \right| dt \qquad (2.31)$$

$$< \omega \bar{r} |K|_{0}^{\theta} \left[2 + \sum_{j=1}^{n} |a_{j}|_{0} \right]^{\theta} \stackrel{\text{def}}{=} B_{2}.$$

According to this and (2.30), we have

$$x(t) = x(t^{0}) + \int_{t^{0}}^{t} x'(t) dt \ge x(t^{0}) - \int_{0}^{\omega} |x'(t)| dt > -B_{1} - B_{2}.$$
 (2.32)

From this and (2.28) it follows that

$$|x(t)|_{0} < \max\{|\ln|K|_{0}|, B_{1} + B_{2}\}.$$
(2.33)

The rest of the proof is similar to the one of Theorem 2.2 and will be omitted. \Box

REMARK 2.6. In (1.3), when $b_j = 0$ for all j = 1, 2, ..., n, equation (1.3) becomes the well-known logistic equation with several delays.

ACKNOWLEDGEMENT. This work was partially supported by the ABF of Yunnan Province of China.

REFERENCES

- [1] H. I. Freedman and J. H. Wu, Periodic solutions of single-species models with periodic delay, SIAM J. Math. Anal. 23 (1992), no. 3, 689–701. MR 93e:92012. Zbl 764.92016.
- [2] R. E. Gaines and J. L. Mawhin, Coincidence Degree, and Nonlinear Differential Equations, Lecture Notes in Mathematics, vol. 568, Springer-Verlag, Berlin, New York, 1977. MR 58#30551. Zbl 339.47031.
- [3] K. Gopalsamy, Stability and Oscillations in Delay Differential Equations of Population Dynamics, Mathematics and its Applications, vol. 74, Kluwer Academic Publishers Group, Dordrecht, 1992. MR 93c:34150. Zbl 752.34039.
- [4] K. Gopalsamy, M. R. S. Kulenović, and G. Ladas, Environmental periodicity and time delays in a "food-limited" population model, J. Math. Anal. Appl. 147 (1990), no. 2, 545-555. MR 91f:92020. Zbl 701.92021.
- [5] J. K. Hale and J. Mawhin, Coincidence degree and periodic solutions of neutral equations, J. Differential Equations 15 (1974), 295-307. MR 49#780. Zbl 274.34070.
- [6] J. K. Hale and S. M. Verduyn Lunel, Introduction to Functional-differential Equations, Applied Mathematical Sciences, vol. 99, Springer-Verlag, New York, 1993. MR 94m:34169. Zbl 787.34002.
- [7] L. Y. Kun, Periodic solution of a periodic neutral delay equation, J. Math. Anal. Appl. 214 (1997), no. 1, 11-21. MR 99j:34101. Zbl 894.34075.
- [8] Y. Li, Periodic solutions of a periodic delay predator-prey system, Proc. Amer. Math. Soc. 127 (1999), no. 5, 1331-1335. MR 99i:34101. Zbl 917.34057.
- [9] J. Mawhin, Periodic solutions of some vector retarded functional differential equations, J. Math. Anal. Appl. 45 (1974), 588-603. MR 48#11725. Zbl 275.34070.
- [10]., Topological Degree Methods in Nonlinear Boundary Value Problems, CBMS Regional Conference Series in Mathematics, no. 40, American Mathematical Society, Providence, R.I., 1979, Expository lectures from the CBMS Regional Conference held at Harvey Mudd College, Claremont, Calif., June 9-15, 1977. MR 80c:47055. Zbl 414.34025.

YONGKUN LI: DEPARTMENT OF MATHEMATICS, YUNNAN UNIVERSITY, KUNMING, YUNNAN 650091, CHINA

E-mail address: yklie@ynu.edu.cn

271



Advances in **Operations Research**



The Scientific World Journal







Hindawi

Submit your manuscripts at http://www.hindawi.com



Algebra



Journal of Probability and Statistics



International Journal of Differential Equations





Complex Analysis





Mathematical Problems in Engineering



Abstract and Applied Analysis



Discrete Dynamics in Nature and Society



International Journal of Mathematics and Mathematical Sciences





Journal of **Function Spaces**



International Journal of Stochastic Analysis

