

EXPLICIT SOLUTION FOR AN INFINITE DIMENSIONAL GENERALIZED INVERSE EIGENVALUE PROBLEM

KAZEM GHANBARI

(Received 28 March 2000 and in revised form 7 September 2000)

ABSTRACT. We study a generalized inverse eigenvalue problem (GIEP), $Ax = \lambda Bx$, in which A is a semi-infinite Jacobi matrix with positive off-diagonal entries $c_i > 0$, and $B = \text{diag}(b_0, b_1, \dots)$, where $b_i \neq 0$ for $i = 0, 1, \dots$. We give an explicit solution by establishing an appropriate spectral function with respect to a given set of spectral data.

2000 Mathematics Subject Classification. 65F18, 47B36.

1. Introduction. Inverse eigenvalue problems, that concern the reconstruction of matrices from a prescribed set of spectral data, are very important subclass of inverse problems that arise in mathematical modeling. There have been challenging work in this area in the last 20 years. Inverse eigenvalue problems mostly are connected to the theory of orthogonal polynomials. An inverse eigenvalue problem appears to be more challenging whence the objective matrix is a specifically structured matrix. Jacobi matrices are among those interesting structured matrices concerned with three term recurrence relations. In this paper we study the generalized inverse eigenvalue problem $Ax = \lambda Bx$, in which A is a semi-infinite Jacobi matrix of the form

$$A = \begin{pmatrix} a_0 & c_0 & 0 & \cdots & \cdots & \cdots \\ c_0 & a_1 & c_1 & 0 & \cdots & \cdots \\ 0 & c_1 & a_2 & c_2 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}, \quad (1.1)$$

$B = \text{diag}(b_0, b_1, \dots)$, where $b_i \neq 0$, and $c_i > 0$, for $i = 0, 1, \dots$. Using the definition of positive definite sequence given by Ahiezer and Krein [1], we establish an explicit formula for recovering the Jacobi matrix A via a given set of spectral data and a diagonal matrix B . We use the usual approach of dealing with this type of problems, that is, the orthogonal polynomials. In this paper, we only discuss the infinite dimensional version of GIEP. The finite dimensional case of this problem has been studied by many authors, see for example [4] for the classical case $Ax = \lambda x$, and [2, 3] for the generalized case $Ax = \lambda Bx$.

2. Three term recurrence relation and spectral function. Let $y_n(\lambda)$ be a solution of

$$\begin{aligned} c_n y_{n+1}(\lambda) &= (b_n \lambda - a_n) y_n(\lambda) - c_{n-1} y_{n-1}(\lambda), \quad n = 0, 1, \dots, \\ y_{-1}(\lambda) &= 0, \quad c_{-1} y_0(\lambda) = 1, \end{aligned} \quad (2.1)$$

where $\{a_n\}_{n \geq 0}$ is an arbitrary sequence of real numbers, $\{b_n\}_{n \geq 0}$, and $\{c_n\}_{n \geq 0}$ are sequences of nonzero real numbers with $c_n > 0$, for $n = 0, 1, \dots$. A complex number λ is said to be an *eigenvalue* of the recurrence relation (2.1) if there is a nonzero solution $y(\lambda) = (y_0(\lambda), y_1(\lambda), \dots)$ satisfying the recurrence relation (2.1). In this case $y(\lambda) = (y_0(\lambda), y_1(\lambda), \dots)^T$ is the *eigenvector* corresponding to λ . If λ_r is an eigenvalue and $y^{(r)} = (y_0(\lambda_r), y_1(\lambda_r), \dots)^T$ is the corresponding eigenvector, then the number

$$\rho_r = \left\{ \sum_{n=0}^{\infty} b_n |y_n(\lambda_r)|^2 \right\} \tag{2.2}$$

is called the *normalization constant*. The set of all eigenvalues and normalization constants of the recurrence relation (2.1) are denoted by $\sigma(A, B)$ and $\rho(A, B)$, respectively.

DEFINITION 2.1. A spectral function $\tau(\lambda)$ associated with the recurrence formula (2.1) is to be nondecreasing, right continuous, satisfying the boundedness requirement

$$\int_{-\infty}^{\infty} \lambda^{2n} d\tau(\lambda) < \infty, \quad n = 0, 1, \dots, \tag{2.3}$$

for all n , and the orthogonality

$$\int_{-\infty}^{\infty} y_p(\lambda) y_q(\lambda) d\tau(\lambda) = b_p^{-1} \delta_{pq}, \quad p, q = 0, 1, \dots \tag{2.4}$$

THEOREM 2.2. *There is at least one spectral function for the recurrence relation (2.1).*

PROOF. See [2, Theorem 5.2.1]. □

THEOREM 2.3. *For every λ , the recurrence relation (2.1) has at least one nontrivial solution of summable square, in the sense that,*

$$\sum_0^{\infty} b_n |y_n(\lambda)|^2 < \infty. \tag{2.5}$$

PROOF. See [2, Theorems 5.4.2, 5.6.1]. □

DEFINITION 2.4. Let $\{y_n(\lambda)\}_{n \geq 0}$ be an arbitrary solution of the recurrence relation (2.1). If

$$\sum_0^{\infty} b_n |y_n(\lambda)|^2 < \infty \tag{2.6}$$

for some $\lambda \in \mathbb{C}$, then we say that the limit-circle case holds for the recurrence relation (2.1).

THEOREM 2.5. *Suppose that the limit circle case holds for one value $\lambda = \lambda_0 \in \mathbb{C}$. Then this is true for all $\lambda \in \mathbb{C}$.*

PROOF. See [2, Theorem 5.6.1]. □

3. Eigenvalues in the limit-circle case. Assuming the limit-circle case to hold, we take it all solutions of (2.1) are summable square in the sense of Theorem 2.3, and uniformly so in any finite λ -region. We therefore have that

$$\sum_0^\infty b_n |y_n(\lambda)|^2 < \eta(\lambda), \tag{3.1}$$

where $\eta(\lambda)$ is some function which is bounded for bounded λ . Adopting some fixed real μ as an eigenvalue, we define the eigenvalues as the zeros of

$$(\lambda - \mu) \sum_0^\infty b_n y_n(\lambda) y_n(\mu). \tag{3.2}$$

By (3.1), the function (3.2) is an entire function of λ . It does not vanish identically, since its derivative is not zero when $\lambda = \mu$. Hence its zeros will have no finite limit. Moreover, these zeros will be real, being the limits of the zeros

$$(\lambda - \mu) \sum_0^m b_n y_n(\lambda) y_n(\mu) = c_{m-1} \{y_m(\lambda) y_{m-1}(\mu) - y_m(\mu) y_{m-1}(\lambda)\}, \tag{3.3}$$

the zeros of the later are real [1, Theorem 4.3.1], and the zeros of (3.2) are the limits of the zeros of (3.3), by Rouché's theorem.

Having defined the eigenvalues λ_r as the roots of

$$\lim_{m \rightarrow \infty} c_{m-1} \{y_m(\lambda) y_{m-1}(\mu) - y_m(\mu) y_{m-1}(\lambda)\} = 0, \tag{3.4}$$

we may define as a spectral function a step function whose jumps are at the λ_r and are of amount

$$\frac{1}{\rho_r} = \left\{ \sum_0^\infty b_n |y_n(\lambda_r)|^2 \right\}^{-1}. \tag{3.5}$$

More formally, we can express this spectral function as follows

$$\tau(\lambda) = \sum_{\lambda_r \leq \lambda} \frac{1}{\rho_r}. \tag{3.6}$$

REMARK 3.1. In general, we may have continuous spectrum for the recurrence relation (2.1). If we consider the discrete Schrödinger operator

$$Hy_n = y_{n+1} + y_{n-1} + a_n y_n, \tag{3.7}$$

where, $\lim |a_n| = 0$, as $n \rightarrow \infty$, then H has continuous spectrum (cf. [5]).

REMARK 3.2. Note that we can display the recurrence relation (2.1) as

$$A\mathcal{Y} = \lambda B\mathcal{Y}, \tag{3.8}$$

where

$$A = \begin{pmatrix} a_0 & c_0 & 0 & \cdots & \cdots & \cdots \\ c_0 & a_1 & c_1 & 0 & \cdots & \cdots \\ 0 & c_1 & a_2 & c_2 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}, \tag{3.9}$$

$$B = \text{diag}(b_0, b_1, \dots), \tag{3.10}$$

$$\mathcal{Y}(\lambda) = (\mathcal{Y}_0(\lambda), \mathcal{Y}_1(\lambda), \dots)^T. \tag{3.11}$$

LEMMA 3.3. Let A and B be the matrices given by (3.9) and (3.10), and let $\mathcal{Y}(\lambda)$ be the vector given by (3.11). Let $\{\lambda_r\}_{r \geq 0}$ and $\{\rho_r\}_{r \geq 0}$ be the sequence of eigenvalues and the normalization constants, respectively. Then

(i) If $i \neq j$ then $\mathcal{Y}(\lambda_i)$ is orthogonal to $\mathcal{Y}(\lambda_j)$ in the sense that

$$\sum_{p=0}^{\infty} b_p \mathcal{Y}_p(\lambda_i) \mathcal{Y}_p(\lambda_j) = \rho_i \delta_{ij}. \tag{3.12}$$

(ii) If $p, q \geq 0$ then

$$\sum_{r=0}^{\infty} \mathcal{Y}_p(\lambda_r) \mathcal{Y}_q(\lambda_r) \rho_r^{-1} = b_p^{-1} \delta_{pq}. \tag{3.13}$$

The property (ii) is called the dual orthogonality.

PROOF. See [2, page 133]. □

4. Infinite dimensional GIEP

PROBLEM 4.1. Let $B = \text{diag}(b_0, b_1, \dots)$ be a given diagonal real matrix, where $b_i \neq 0$ for $i = 0, 1, \dots$, and let $\{\lambda_i\}$ and $\{\rho_i\}$ be real numbers such that $\lambda_i \rho_i > 0$, for $i = 0, 1, \dots$. We find a positive definite semi-infinite Jacobi matrix A that satisfies (3.8) with

(i) $\sigma(A, B) = \{\lambda_0, \lambda_1, \dots\}$,

(ii) $\rho(A, B) = \{\rho_0, \rho_1, \dots\}$.

We call **Problem 4.1** infinite dimensional GIEP. It is easy to see that if there is a solution for GIEP then $\lambda_i \rho_i > 0$, for $i = 0, 1, \dots$. For, if $\lambda_i \in \sigma(A, B)$, and $\mathcal{Y}^{(i)}$ is the corresponding eigenvector, then

$$0 < \mathcal{Y}^{(i)T} A \mathcal{Y}^{(i)} = \lambda_i \mathcal{Y}^{(i)T} B \mathcal{Y}^{(i)} = \lambda_i \rho_i, \quad \text{for } i = 0, 1, \dots \tag{4.1}$$

LEMMA 4.2. Equation (2.4) with $p \neq q$ is equivalent to

$$\int_{-\infty}^{\infty} \mathcal{Y}_p(\lambda) \lambda^q d\tau(\lambda) = 0, \quad 0 \leq q \leq p - 1. \tag{4.2}$$

PROOF. See [2, Theorem 4.6.1]. □

DEFINITION 4.3. Let $\tau(\lambda)$ be the spectral function defined by (3.6). The scalars

$$\mu_j = \int_{-\infty}^{\infty} \lambda^j d\tau(\lambda), \quad j = 0, 1, \dots \tag{4.3}$$

are called the moments of $\tau(\lambda)$.

Note that formula (4.3) is equivalent to

$$\mu_j = \sum_0^{\infty} \frac{\lambda_r^j}{\rho_r}, \quad j = 0, 1, \dots, \tag{4.4}$$

and since $\lambda_r \rho_r > 0$, the odd moments μ_{2n+1} are all positive.

DEFINITION 4.4. Let $\tau(\lambda)$ be the spectral function defined by (3.6), and let μ_j be the moments of $\tau(\lambda)$ given by (4.4). We define M_n and Δ_n by

$$M_n = \begin{pmatrix} \mu_0 & \mu_1 & \cdots & \mu_{n-1} \\ \mu_1 & \mu_2 & \cdots & \mu_n \\ \vdots & \vdots & \vdots & \vdots \\ \mu_{n-1} & \mu_n & \cdots & \mu_{2n-2} \end{pmatrix}, \tag{4.5}$$

$$\Delta_n = \det(M_n), \quad \Delta_0 = 1.$$

LEMMA 4.5. Let A_{ij} be the determinant of the matrix obtained from M_n by deleting row i and column j . If $n = 1$ we set $A_{11} = 1$. Then

$$\begin{aligned} \Delta_{n+1} &= \mu_n \sum_{k=1}^n (-1)^k \mu_{n-1+k} A_{1k} \\ &+ \mu_{n+1} \sum_{k=1}^n (-1)^{k+1} \mu_{n-1+k} A_{2k} + \cdots \\ &+ \mu_{n+r} \sum_{k=1}^n (-1)^{r+k} \mu_{n-1+k} A_{r+1+k} + \cdots \\ &+ \mu_{2n-1} \sum_{k=1}^n (-1)^{n-1+k} \mu_{n-1+k} A_{nk} + \mu_{2n} \Delta_n. \end{aligned} \tag{4.6}$$

PROOF. The proof follows by induction on n . □

5. Positive definite sequences. In this section we use the concept of positive definite sequences given by Aheizer and Krein [1] to prove that the sequence of the moments of the spectral function $\tau(\lambda)$ is a positive definite sequence.

DEFINITION 5.1. Let $J = (a, b)$ ($-\infty \leq a < b \leq \infty$) be an interval in \mathbb{R} . An infinite sequence $\{s_k\}_{k \geq 0}$ is said to be a *positive definite* sequence on J if for every nonnegative polynomial $R_n(\lambda) = p_0 + p_1\lambda + p_2\lambda^2 + \dots + p_n\lambda^n$ in J , the sequence $\{s_k\}_{k \geq 0}$ satisfies the property

$$\sum_{j=0}^n p_j s_j > 0, \quad \text{for } n = 0, 1, 2, \dots \tag{5.1}$$

THEOREM 5.2. Let $\{r_n\}_{n \geq 1}$ be a sequence of positive real numbers and let $\xi_1 < \xi_2 < \dots$ be a sequence of real numbers such that $\sum r_i \xi_i^k < \infty$, for all $k \geq 0$. Put

$$s_k = \sum_{i=1}^{\infty} r_i \xi_i^k, \quad k = 0, 1, \dots \tag{5.2}$$

Then, the sequence $\{s_k\}_{k \geq 0}$ is positive definite in every interval (a, b) satisfying $-\infty \leq a < \xi_1 < \xi_2 < \dots < b \leq \infty$.

PROOF. Let $\varphi(\lambda)$ be any real nonnegative polynomial in the interval (a, b) , say, $\varphi(\lambda) = \sum_0^n p_k \lambda^k$. We have

$$\sum_{k=0}^n p_k s_k = \sum_{k=0}^n \sum_{i=1}^{\infty} p_k r_i \xi_i^k = \sum_{i=1}^{\infty} r_i \left(\sum_{k=0}^n p_k \xi_i^k \right) = \sum_{i=1}^{\infty} r_i \varphi(\xi_i) > 0. \tag{5.3}$$

Thus the proof is complete. □

COROLLARY 5.3. Let $\{\lambda_i\}_{i \geq 0}$ and $\{\rho_i\}_{i \geq 0}$ be the eigenvalues and the normalization constants, respectively. If $\lambda_i \rho_i > 0$ for $i \geq 0$, then $\{\mu_i\}_{i \geq 1}$ is a positive definite sequence.

PROOF. We define the real numbers

$$r_k = \frac{\lambda_{k-1}}{\rho_{k-1}}, \quad k \geq 1, \quad \xi_k = \lambda_{k-1}, \quad k \geq 1. \tag{5.4}$$

If we set

$$s_k = \sum_1^{\infty} r_i \xi_i^k, \quad k = 0, 1, \dots, \tag{5.5}$$

then $r_k > 0$ for $k \geq 1$. Thus, by [Theorem 5.2](#), the sequence $\{s_k\}_{k \geq 0}$ is positive definite. By formula [\(4.3\)](#), $s_k = \mu_{k+1}$, ($k = 0, 1, \dots$) hence $\{\mu_i\}_{i \geq 1}$ is a positive definite sequence. □

DEFINITION 5.4. An infinite real quadratic form

$$\sum_{i,k=0}^{\infty} a_{ik} \xi_i \xi_k, \quad (a_{ik} = a_{ki}) \tag{5.6}$$

is said to be positive if all its partial sums

$$\sum_{i,k=0}^n a_{ik} \xi_i \xi_k, \quad (n = 0, 1, \dots) \tag{5.7}$$

are positive.

THEOREM 5.5. *The sequence $\{s_n\}_{n \geq 0}$ is positive definite in the interval $(-\infty, \infty)$ if the infinite quadratic form*

$$\sum_{i,k=0}^{\infty} s_{i+k} \xi_i \xi_k \tag{5.8}$$

is positive.

PROOF. See [1, page 3, Theorem 1]. □

THEOREM 5.6. *Let $\{\lambda_i\}_{i \geq 0}$ and $\{\rho_i\}_{i \geq 0}$ be the eigenvalues and the normalization constants of [Problem 4.1](#). Let $\lambda_i \rho_i > 0$, and let $\{\mu_i\}_{i \geq 0}$ be a sequence of moments given by (4.4). Then the quadratic form*

$$\sum_{i,k=0}^{\infty} \mu_{i+k+1} \xi_i \xi_k \tag{5.9}$$

is positive.

PROOF. Since $\lambda_i \rho_i > 0$, by [Corollary 5.3](#), the sequence $\{\mu\}_{i \geq 1}$ is positive definite. By [Theorem 5.5](#), the infinite quadratic form

$$\sum_{i,k=0}^{\infty} \mu_{i+k+1} \xi_i \xi_k \tag{5.10}$$

is positive. □

THEOREM 5.7. *Let $\{\lambda_i\}_{i \geq 0}$ and $\{\rho_i\}_{i \geq 0}$ be the eigenvalues and the normalization constants of [Problem 4.1](#). Let $\lambda_i \rho_i > 0$, and let $\{\mu_i\}_{i \geq 0}$ be a sequence of moments given by (4.4). Then*

$$\det \begin{pmatrix} \mu_1 & \mu_2 & \cdots & \mu_n \\ \mu_2 & \mu_3 & \cdots & \mu_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_n & \mu_{n+1} & \cdots & \mu_{2n-1} \end{pmatrix} > 0, \tag{5.11}$$

for $n = 1, 2, \dots$

PROOF. By [Theorem 5.6](#), the quadratic form

$$\sum_{i,k=0}^{\infty} \mu_{i+k+1} \xi_i \xi_k \tag{5.12}$$

is positive. Then, the determinants of the principal submatrices are positive, that is,

$$\det \begin{pmatrix} \mu_1 & \mu_2 & \cdots & \mu_n \\ \mu_2 & \mu_3 & \cdots & \mu_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_n & \mu_{n+1} & \cdots & \mu_{2n-1} \end{pmatrix} > 0, \tag{5.13}$$

for $n = 1, 2, \dots$ □

6. Construction of a solution for Problem 4.1. In this section, we give an explicit solution for Problem 4.1, if the *limit circle* case holds for the recurrence relation (2.1).

THEOREM 6.1. *Let $\tau(\lambda)$ be the spectral function defined by (3.6), and let $B = \text{diag}(b_0, b_1, \dots)$ be given real matrix, where $\{b_n\}_{n \geq 0}$ is a sequence of nonzero real numbers. If the relation*

$$b_n \Delta_n \Delta_{n+1} > 0 \quad (6.1)$$

holds for all $n \geq 0$, then there exists a countable set of orthogonal polynomials $\{\gamma_n(\lambda)\}_{n \geq 0}$ in the sense of dual orthogonality property (3.13), and the polynomials are determined up to change of sign. Moreover, the polynomials $\{\gamma_n(\lambda)\}_{n \geq 0}$ are dense in L^2_τ .

PROOF. For the first part, we seek polynomials of the form

$$\gamma_n(\lambda) = \beta_n \left(\lambda^n + \sum_{k=0}^{n-1} \alpha_{nk} \lambda^k \right), \quad k = 0, 1, \dots, \quad (6.2)$$

where $\gamma_0(\lambda) = \beta_0$ and $\beta_n \neq 0$. Using (3.13) and (4.3) we get

$$\beta_0^2 = \frac{1}{\mu_0 b_0} = \frac{\Delta_0}{b_0 \Delta_1}, \quad (6.3)$$

that is positive by assumption. It follows from (6.2) that

$$(\gamma_n(\lambda))^2 = \beta_n^2 \left[\lambda^{2n} + \sum_{k=0}^{n-1} \alpha_{nk} \lambda^{n+k} \right] + \beta_n \gamma_n(\lambda) \sum_{k=0}^{n-1} \alpha_{nk} \lambda^k, \quad n = 1, 2, \dots \quad (6.4)$$

Combining (3.13), (4.2), and (6.4), it follows that

$$b_n^{-1} = \beta_n^2 \left(\mu_{2n} + \sum_{k=0}^{n-1} \alpha_{nk} \mu_{n+k} \right). \quad (6.5)$$

This gives β_n in terms of α_{nk} and the moments. To determine α_{nk} , we substitute $\gamma_n(\lambda)$ given by (6.2) in (4.2) and we obtain

$$\mu_{n+k} + \sum_{k=0}^{n-1} \alpha_{nk} \mu_{n+k} = 0, \quad (6.6)$$

where $0 \leq k \leq n-1$ and $n = 1, 2, \dots$. Using matrix notation, we have

$$M_n (\alpha_{n0}, \alpha_{n1}, \dots, \alpha_{n,n-1})^T = (-\mu_n, -\mu_{n+1}, \dots, -\mu_{2n-1})^T. \quad (6.7)$$

Therefore

$$\alpha_{nr} = \frac{\sum_{k=1}^n (-1)^{k+r} \mu_{n-1+k} A_{r+1,k}}{\Delta_n}, \tag{6.8}$$

where $0 \leq r \leq n - 1$ and $n = 1, 2, \dots$. Substituting (4.6) in (6.8), we get

$$\Delta_{n+1} = \Delta_n \left(\mu_{2n} + \sum_{k=0}^{n-1} \alpha_{nk} \mu_{n+k} \right). \tag{6.9}$$

Using (6.5) in (6.9) we obtain

$$\beta_n^2 = \frac{\Delta_n}{\Delta_{n+1} b_n} \tag{6.10}$$

that is positive by the assumption, and this completes the proof. For the second part see [2, page 141]. □

THEOREM 6.2. *Let the assumptions of Theorem 6.1 hold. Then, the Jacobi matrix A of the form*

$$A = \begin{pmatrix} a_0 & c_0 & 0 & \cdots & \cdots & \cdots \\ c_0 & a_1 & c_1 & 0 & \cdots & \cdots \\ 0 & c_1 & a_2 & c_2 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}, \tag{6.11}$$

given by

$$\begin{aligned} a_0 &= b_0 \alpha_{1,0}, & a_n &= b_n (\alpha_{n,n-1} - \alpha_{n+1,n}), & n &= 1, 2, \dots, \\ \alpha_{nr} &= \frac{\sum_{k=1}^n (-1)^{k+r} \mu_{n-1+k} A_{r+1,k}}{\Delta_n}, & 0 \leq r &\leq n - 1, & n &= 1, 2, \dots, \\ c_n &= \frac{b_n \beta_n}{\beta_{n+1}}, & \beta_n &= \frac{\Delta_n}{\Delta_{n+1} b_n}, & n &= 0, 1, \dots, \end{aligned} \tag{6.12}$$

is a solution for Problem 4.1, which assumes $\tau(\lambda)$ as its spectral function, where

$$\mathcal{Y}(\lambda) = (\mathcal{Y}_0(\lambda), \mathcal{Y}_1(\lambda), \dots)^T, \quad \mathcal{Y}_n(\lambda) = \beta_n \left(\lambda^n + \sum_{r=0}^{n-1} \alpha_{nr} \lambda^r \right), \quad n = 0, 1, \dots \tag{6.13}$$

Moreover, the matrix A is positive definite in the sense of Definition 5.4.

PROOF. Substituting $\mathcal{Y}_n(\lambda)$ given by (6.2) in the recurrence relation (2.1) and comparing the corresponding coefficients of the powers of λ , we obtain

$$a_0 = -b_0 \alpha_{1,0}, \quad c_n \beta_{n+1} = b_n \beta_n, \tag{6.14}$$

$$c_n \beta_{n+1} \alpha_{n+1,n} + a_n \beta_n = b_n \beta_n \alpha_{n,n-1}, \quad n \geq 1. \tag{6.15}$$

By substituting (6.14) in (6.15) we get

$$a_n = b_n(\alpha_{n,n-1} - \alpha_{n+1,n}), \quad (6.16)$$

which completes the first part of the proof. In order to prove A is positive definite, by Definition 5.4, it suffices to prove that the determinants of leading principal submatrices of A are all positive. Let D_n be the determinant of the $n \times n$ leading submatrix of A in the upper left corner of A . It is easy to check by induction that

$$D_n = b_0 b_1 \cdots b_{n-1} (-1)^n \alpha_{n,0}. \quad (6.17)$$

This is equivalent to

$$D_n = \frac{b_0 \Delta_0}{\Delta_1} \frac{b_1 \Delta_1}{\Delta_2} \cdots \frac{b_{n-1} \Delta_{n-1}}{\Delta_n} (-1)^n \alpha_{n,0} \Delta_n. \quad (6.18)$$

By using (6.8) we obtain

$$(-1)^n \alpha_{n,0} \Delta_n = \det \begin{pmatrix} \mu_1 & \mu_2 & \cdots & \mu_n \\ \mu_2 & \mu_3 & \cdots & \mu_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_n & \mu_{n+1} & \cdots & \mu_{2n-1} \end{pmatrix}. \quad (6.19)$$

Therefore, combining (6.19) with the assumption of Theorem 6.1 implies that $D_n > 0$ if and only if

$$\det \begin{pmatrix} \mu_1 & \mu_2 & \cdots & \mu_n \\ \mu_2 & \mu_3 & \cdots & \mu_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_n & \mu_{n+1} & \cdots & \mu_{2n-1} \end{pmatrix} > 0, \quad (6.20)$$

which is true by Theorem 5.7, for $n = 1, 2, \dots$. This completes the proof. \square

ACKNOWLEDGEMENTS. The author would like to thank Professor Angelo Mingarelli for his support, and the anonymous referee for valuable comments. The author is also indebted to the Ministry of Culture and Higher Education of Islamic Republic of Iran for the financial support of this work.




REFERENCES

- [1] N. I. Ahiezer and M. Krein, *Some Questions in the Theory of Moments*, Translations of Mathematical Monographs, vol. 2, American Mathematical Society, Rhode Island, 1962. MR 29#5073. Zbl 117.32702.
- [2] F. V. Atkinson, *Discrete and Continuous Boundary Problems*, Mathematics in Science and Engineering, vol. 8, Academic Press, New York, 1964. MR 31#416. Zbl 117.05806.
- [3] K. Ghanbari and A. Mingarelli, *Generalized inverse eigenvalue problem for symmetric matrices*, Int. J. Appl. Math. 4 (2000), no. 2, 199-209.
- [4] O. H. Hald, *Inverse eigenvalue problems for Jacobi matrices*, Linear Algebra and Appl. 14 (1976), no. 1, 63-85. MR 56#5594. Zbl 328.15007.

- [5] J. Janas and S. Naboko, *Jacobi matrices with absolutely continuous spectrum*, Proc. Amer. Math. Soc. **127** (1999), no. 3, 791–800. [MR 99f:47042](#). [Zbl 910.47022](#).

KAZEM GHANBARI: SCHOOL OF MATHEMATICS AND STATISTICS, CARLETON UNIVERSITY,
OTTAWA, CANADA ON K1S 5B6

E-mail address: kghanbar@math.carleton.ca



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

