ON θ -PRECONTINUOUS FUNCTIONS

TAKASHI NOIRI

(Received 16 January 2001)

ABSTRACT. We introduce a new class of functions called θ -precontinuous functions which is contained in the class of weakly precontinuous (or almost weakly continuous) functions and contains the class of almost precontinuous functions. It is shown that the θ -precontinuous image of a p -closed space is quasi H -closed.

2000 Mathematics Subject Classification. 54C08.

1. Introduction. A subset A of a topological space X is said to be preopen [14] or nearly open [26] if $A \subset \text{Int}(\text{Cl}(A))$. A function $f : X \rightarrow Y$ is called precontinuous [14] if the preimage $f^{-1}(V)$ of each open set V of Y is preopen in X . Precontinuity was called near continuity by Pták [26] and also called almost continuity by Frolík [9] and Husain [10]. In 1985, Janković [12] introduced almost weak continuity as a weak form of precontinuity. Popa and Noiri [23] introduced weak precontinuity and showed that almost weak continuity is equivalent to weak precontinuity. Paul and Bhattacharyya [21] called weakly precontinuous functions quasi precontinuous and obtained the further properties of quasi precontinuity. Recently, Nasef and Noiri [16] have introduced and investigated the notion of almost precontinuity. Quite recently, Jafari and Noiri [11] investigated the further properties of almost precontinuous functions.

In this paper, we introduce a new class of functions called θ -precontinuous functions which is contained in the class of weakly precontinuous functions and contains the class of almost precontinuous functions. We obtain basic properties of θ -precontinuous functions. It is shown in the last section that the θ -precontinuous images of p -closed (resp., β -connected) spaces are quasi H -closed (resp., semi-connected).

2. Preliminaries. Throughout, by (X, τ) and (Y, σ) (or simply X and Y) we denote topological spaces. Let S be a subset of X . We denote the interior and the closure of S by $\text{Int}(S)$ and $\text{Cl}(S)$, respectively. A subset S is said to be *preopen* [14] (resp., *semi-open* [13], *α -open* [17]) if $S \subset \text{Int}(\text{Cl}(S))$ (resp., $S \subset \text{Cl}(\text{Int}(S))$, $S \subset \text{Int}(\text{Cl}(\text{Int}(S)))$). The complement of a preopen set is called *preclosed*. The intersection of all preclosed sets containing S is called the *preclosure* [8] of S and is denoted by $\text{pCl}(S)$. The *preinterior* of S is defined by the union of all preopen sets contained in S and is denoted by $\text{pInt}(S)$. The family of all preopen sets of X is denoted by $\text{PO}(X)$. We set $\text{PO}(X, x) = \{U : x \in U \text{ and } U \in \text{PO}(X)\}$. A point x of X is called a θ -cluster point of S if $\text{Cl}(U) \cap S \neq \emptyset$ for every open set U of X containing x . The set of all θ -cluster points of S is called the θ -closure of S and is denoted by $\text{Cl}_\theta(S)$. A subset S is said to be θ -closed [27] if $S = \text{Cl}_\theta(S)$. The complement of a θ -closed set is said to be θ -open. A point x of X

is called a *pre θ -cluster point* of S if $\text{pCl}(U) \cap S \neq \emptyset$ for every preopen set U of X containing x . The set of all pre θ -cluster points of S is called the *pre θ -closure* of S and is denoted by $\text{pCl}_\theta(S)$. A subset S is said to be *pre θ -closed* [20] if $S = \text{pCl}_\theta(S)$. The complement of a pre θ -closed set is said to be *pre θ -open*.

DEFINITION 2.1. A function $f : X \rightarrow Y$ is said to be *precontinuous* [14] (resp., *almost precontinuous* [16], *weakly precontinuous* [23] or *quasi precontinuous* [21]) if for each $x \in X$ and each open set V of Y containing $f(x)$, there exists $U \in \text{PO}(X, x)$ such that $f(U) \subset V$ (resp., $f(U) \subset \text{Int}(\text{Cl}(V))$, $f(U) \subset \text{Cl}(V)$).

DEFINITION 2.2. A function $f : X \rightarrow Y$ is said to be *almost weakly continuous* [12] if $f^{-1}(V) \subset \text{Int}(\text{Cl}(f^{-1}(\text{Cl}(V))))$ for every open set V of Y .

DEFINITION 2.3. A function $f : X \rightarrow Y$ is said to be *strongly θ -precontinuous* [19] if for each $x \in X$ and each open set V of Y containing $f(x)$, there exists $U \in \text{PO}(X, x)$ such that $f(\text{pCl}(U)) \subset V$.

DEFINITION 2.4. A function $f : X \rightarrow Y$ is said to be *θ -precontinuous* if for each $x \in X$ and each open set V of Y containing $f(x)$, there exists $U \in \text{PO}(X, x)$ such that $f(\text{pCl}(U)) \subset \text{Cl}(V)$.

REMARK 2.5. By the above definitions and Theorem 3.3 below, we have the following implications and none of these implications is reversible by [19, Example 2.2], [11, Example 2.9], and Examples 2.6 and 5.11 below.

$$\begin{aligned} \text{strongly } \theta\text{-precontinuous} &\Rightarrow \text{precontinuous} \Rightarrow \text{almost precontinuous} \\ &\Rightarrow \theta\text{-precontinuous} \Rightarrow \text{weakly precontinuous.} \end{aligned} \quad (2.1)$$

EXAMPLE 2.6. This example is due to Arya and Deb [4]. Let X be the set of all real numbers. The topology τ on X is the cocountable topology. Let $Y = \{a, b, c\}$, $\sigma = \{\emptyset, Y, \{a\}, \{c\}, \{a, c\}\}$. We define a function $f : (X, \tau) \rightarrow (Y, \sigma)$ by $f(x) = a$ if x is rational; $f(x) = b$ if x is irrational. Then f is a θ -precontinuous function which is not almost precontinuous

3. Characterizations

THEOREM 3.1. For a function $f : X \rightarrow Y$ the following properties are equivalent:

- (1) f is θ -precontinuous;
- (2) $\text{pCl}_\theta(f^{-1}(B)) \subset f^{-1}(\text{Cl}_\theta(B))$ for every subset B of Y ;
- (3) $f(\text{pCl}_\theta(A)) \subset \text{Cl}_\theta(f(A))$ for every subset A of X .

PROOF. (1) $\Rightarrow$ (2). Let B be any subset of Y . Suppose that $x \notin f^{-1}(\text{Cl}_\theta(B))$. Then $f(x) \notin \text{Cl}_\theta(B)$ and there exists an open set V containing $f(x)$ such that $\text{Cl}(V) \cap B = \emptyset$. Since f is θ -p.c., there exists $U \in \text{PO}(X, x)$ such that $f(\text{pCl}(U)) \subset \text{Cl}(V)$. Therefore, we have $f(\text{pCl}(U)) \cap B = \emptyset$ and $\text{pCl}(U) \cap f^{-1}(B) = \emptyset$. This shows that $x \notin \text{pCl}_\theta(f^{-1}(B))$. Thus, we obtain $\text{pCl}_\theta(f^{-1}(B)) \subset f^{-1}(\text{Cl}_\theta(B))$.

(2) $\Rightarrow$ (3). Let A be any subset of X . Then we have $\text{pCl}_\theta(A) \subset \text{pCl}_\theta(f^{-1}(f(A))) \subset f^{-1}(\text{Cl}_\theta(f(A)))$ and hence $f(\text{pCl}_\theta(A)) \subset \text{Cl}_\theta(f(A))$.

(3) $\Rightarrow$ (2). Let B be a subset of Y . We have $f(\text{pCl}_\theta(f^{-1}(B))) \subset \text{Cl}_\theta(f(f^{-1}(B))) \subset \text{Cl}_\theta(B)$ and hence $\text{pCl}_\theta(f^{-1}(B)) \subset f^{-1}(\text{Cl}_\theta(B))$.

(2) $\Rightarrow$ (1). Let $x \in X$ and V be an open set of Y containing $f(x)$. Then we have $\text{Cl}(V) \cap (Y - \text{Cl}(V)) = \emptyset$ and $f(x) \notin \text{Cl}_\theta(Y - \text{Cl}(V))$. Hence, $x \notin f^{-1}(\text{Cl}_\theta(Y - \text{Cl}(V)))$ and $x \notin \text{pCl}_\theta(f^{-1}(Y - \text{Cl}(V)))$. There exists $U \in \text{PO}(X, x)$ such that $\text{pCl}(U) \cap f^{-1}(Y - \text{Cl}(V)) = \emptyset$; hence $f(\text{pCl}(U)) \subset \text{Cl}(V)$. Therefore, f is $\theta.p.c.$ $\square$

THEOREM 3.2. For a function $f : X \rightarrow Y$ the following properties are equivalent:

- (1) f is θ -precontinuous;
- (2) $f^{-1}(V) \subset \text{pInt}_\theta(f^{-1}(\text{Cl}(V)))$ for every open set V of Y ;
- (3) $\text{pCl}_\theta(f^{-1}(V)) \subset f^{-1}(\text{Cl}(V))$ for every open set V of Y .

PROOF. (1) $\Rightarrow$ (2). Suppose that V is any open set of Y and $x \in f^{-1}(V)$. Then $f(x) \in V$ and there exists $U \in \text{PO}(X, x)$ such that $f(\text{pCl}(U)) \subset \text{Cl}(V)$. Therefore, $x \in U \subset \text{pCl}(U) \subset f^{-1}(\text{Cl}(V))$. This shows that $x \in \text{pInt}_\theta(f^{-1}(\text{Cl}(V)))$. Therefore, we obtain $f^{-1}(V) \subset \text{pInt}_\theta(f^{-1}(\text{Cl}(V)))$.

(2) $\Rightarrow$ (3). Suppose that V is any open set of Y and $x \notin f^{-1}(\text{Cl}(V))$. Then $f(x) \notin \text{Cl}(V)$ and there exists an open set W containing $f(x)$ such that $W \cap V = \emptyset$; hence $\text{Cl}(W) \cap V = \emptyset$. Therefore, we have $f^{-1}(\text{Cl}(W)) \cap f^{-1}(V) = \emptyset$. Since $x \in f^{-1}(W)$, by (2) $x \in \text{pInt}_\theta(f^{-1}(\text{Cl}(W)))$. There exists $U \in \text{PO}(X, x)$ such that $\text{pCl}(U) \subset f^{-1}(\text{Cl}(W))$. Thus we have $\text{pCl}(U) \cap f^{-1}(V) = \emptyset$ and hence $x \notin \text{pCl}_\theta(f^{-1}(V))$. This shows that $\text{pCl}_\theta(f^{-1}(V)) \subset f^{-1}(\text{Cl}(V))$.

(3) $\Rightarrow$ (1). Suppose that $x \in X$ and V is any open set of Y containing $f(x)$. Then $V \cap (Y - \text{Cl}(V)) = \emptyset$ and $f(x) \notin \text{Cl}(Y - \text{Cl}(V))$. Therefore, $x \notin f^{-1}(\text{Cl}(Y - \text{Cl}(V)))$ and by (3) $x \notin \text{pCl}_\theta(f^{-1}(Y - \text{Cl}(V)))$. There exists $U \in \text{PO}(X, x)$ such that $\text{pCl}(U) \cap f^{-1}(Y - \text{Cl}(V)) = \emptyset$. Therefore, we obtain $f(\text{pCl}(U)) \subset \text{Cl}(V)$. This shows that f is $\theta.p.c.$ $\square$

THEOREM 3.3. For a function $f : X \rightarrow Y$ the following properties hold:

- (1) if f is almost precontinuous, then it is θ -precontinuous;
- (2) if f is θ -precontinuous, then it is weakly precontinuous.

PROOF. Statement (2) is obvious. We will show statement (1). Suppose that $x \in X$ and V is any open set of Y containing $f(x)$. Since f is almost precontinuous, $f^{-1}(\text{Int}(\text{Cl}(V)))$ is preopen and $f^{-1}(\text{Cl}(V))$ is preclosed in X by [16, Theorem 3.1]. Now, set $U = f^{-1}(\text{Int}(\text{Cl}(V)))$. Then we have $U \in \text{PO}(X, x)$ and $\text{pCl}(U) \subset f^{-1}(\text{Cl}(V))$. Therefore, we obtain $f(\text{pCl}(U)) \subset \text{Cl}(V)$. This shows that f is $\theta.p.c.$ $\square$

COROLLARY 3.4. Let Y be a regular space. Then, for a function $f : X \rightarrow Y$ the following properties are equivalent:

- (1) f is strongly θ -precontinuous;
- (2) f is precontinuous;
- (3) f is almost precontinuous;
- (4) f is θ -precontinuous;
- (5) f is weakly precontinuous.

PROOF. This is an immediate consequence of [19, Theorem 3.2]. $\square$

DEFINITION 3.5. A topological space X is said to be *pre-regular* [20] if for each preclosed set F and each point $x \in X - F$, there exist disjoint preopen sets U and V such that $x \in U$ and $F \subset V$.

LEMMA 3.6 (see [20]). *A topological space X is pre-regular if and only if for each $U \in \text{PO}(X)$ and each point $x \in U$, there exists $V \in \text{PO}(X, x)$ such that $x \in V \subset \text{pCl}(V) \subset U$.*

THEOREM 3.7. *Let X be a pre-regular space. Then $f : X \rightarrow Y$ is $\theta.p.c.$ if and only if it is weakly precontinuous.*

PROOF. Suppose that f is weakly precontinuous. Let $x \in X$ and V is any open set of Y containing $f(x)$. Then, there exists $U \in \text{PO}(X, x)$ such that $f(U) \subset \text{Cl}(V)$. Since X is pre-regular, there exists $U_* \in \text{PO}(X, x)$ such that $x \in U_* \subset \text{pCl}(U_*) \subset U$. Therefore, we obtain $f(\text{pCl}(U_*)) \subset \text{Cl}(V)$. This shows that f is $\theta.p.c.$ $\square$

THEOREM 3.8. *Let $f : X \rightarrow Y$ be a function and $g : X \rightarrow X \times Y$ the graph function of f defined by $g(x) = (x, f(x))$ for each $x \in X$. Then g is $\theta.p.c.$ if and only if f is $\theta.p.c.$*

PROOF

NECESSITY. Suppose that g is $\theta.p.c.$ Let $x \in X$ and V be an open set of Y containing $f(x)$. Then $X \times V$ is an open set of $X \times Y$ containing $g(x)$. Since g is $\theta.p.c.$, there exists $U \in \text{PO}(X, x)$ such that $g(\text{pCl}(U)) \subset \text{Cl}(X \times V)$. It follows that $\text{Cl}(X \times V) = X \times \text{Cl}(V)$. Therefore, we obtain $f(\text{pCl}(U)) \subset \text{Cl}(V)$. This shows that f is $\theta.p.c.$

SUFFICIENCY. Let $x \in X$ and W be any open set of $X \times Y$ containing $g(x)$. There exist open sets $U_1 \subset X$ and $V \subset Y$ such that $g(x) = (x, f(x)) \in U_1 \times V \subset W$. Since f is $\theta.p.c.$, there exists $U_2 \in \text{PO}(X, x)$ such that $f(\text{pCl}(U_2)) \subset \text{Cl}(V)$. Let $U = U_1 \cap U_2$, then $U \in \text{PO}(X, x)$. Therefore, we obtain $g(\text{pCl}(U)) \subset \text{Cl}(U_1) \times f(\text{pCl}(U_2)) \subset \text{Cl}(U_1) \times \text{Cl}(V) \subset \text{Cl}(W)$. This shows that g is $\theta.p.c.$ $\square$

4. Some properties

LEMMA 4.1 (see [15]). *Let A and X_0 be subsets of a space X .*

- (1) *If $A \in \text{PO}(X)$ and X_0 is semi-open in X , then $A \cap X_0 \in \text{PO}(X_0)$.*
- (2) *If $A \in \text{PO}(X_0)$ and $X_0 \in \text{PO}(X)$, then $A \in \text{PO}(X)$.*

LEMMA 4.2 (see [7]). *Let A and X_0 be subsets of a space X such that $A \subset X_0 \subset X$. Let $\text{pCl}_{X_0}(A)$ denote the preclosure of A in the subspace X_0 .*

- (1) *If X_0 is semi-open in X , then $\text{pCl}_{X_0}(A) \subset \text{pCl}(A)$.*
- (2) *If $A \in \text{PO}(X_0)$ and $X_0 \in \text{PO}(X)$, then $\text{pCl}(A) \subset \text{pCl}_{X_0}(A)$.*

THEOREM 4.3. *If $f : X \rightarrow Y$ is $\theta.p.c.$ and X_0 is a semi-open subset of X , then the restriction $f/X_0 : X_0 \rightarrow Y$ is $\theta.p.c.$*

PROOF. For any $x \in X_0$ and any open neighborhood V of $f(x)$, there exists $U \in \text{PO}(X, x)$ such that $f(\text{pCl}(U)) \subset \text{Cl}(V)$ since f is $\theta.p.c.$ Put $U_0 = U \cap X_0$, then by Lemmas 4.1 and 4.2 $U_0 \in \text{PO}(X_0, x)$ and $\text{pCl}_{X_0}(U_0) \subset \text{pCl}(U_0)$. Therefore, we obtain

$$(f/X_0)(\text{pCl}_{X_0}(U_0)) = f(\text{pCl}_{X_0}(U_0)) \subset f(\text{pCl}(U_0)) \subset f(\text{pCl}(U)) \subset \text{Cl}(V). \quad (4.1)$$

This shows that f/X_0 is $\theta.p.c.$ $\square$

THEOREM 4.4. *A function $f : X \rightarrow Y$ is $\theta.p.c.$ if for each $x \in X$ there exists $X_0 \in \text{PO}(X, x)$ such that the restriction $f|_{X_0} : X_0 \rightarrow Y$ is $\theta.p.c.$*

PROOF. Let $x \in X$ and V be any open neighborhood of $f(x)$. There exists $X_0 \in \text{PO}(X, x)$ such that $f|_{X_0} : X_0 \rightarrow Y$ is $\theta.p.c.$ Thus, there exists $U \in \text{PO}(X_0, x)$ such that $(f|_{X_0})(\text{pCl}_{X_0}(U)) \subset \text{Cl}(V)$. By Lemmas 4.1 and 4.2, $U \in \text{PO}(X, x)$ and $\text{pCl}(U) \subset \text{pCl}_{X_0}(U)$. Hence, we have $f(\text{pCl}(U)) = (f|_{X_0})(\text{pCl}(U)) \subset (f|_{X_0})(\text{pCl}_{X_0}(U)) \subset \text{Cl}(V)$. This shows that f is $\theta.p.c.$ $\square$

COROLLARY 4.5. *Let $\{U_\lambda : \lambda \in \Lambda\}$ be an α -open cover of a topological space X . A function $f : X \rightarrow Y$ is $\theta.p.c.$ if and only if the restriction $f|_{U_\lambda} : U_\lambda \rightarrow Y$ is $\theta.p.c.$ for each $\lambda \in \Lambda$.*

PROOF. This is an immediate consequence of Theorems 4.3 and 4.4. $\square$

Let $\{X_\alpha : \alpha \in \mathcal{A}\}$ be a family of topological spaces, A_α a nonempty subset of X_α for each $\alpha \in \mathcal{A}$ and $X = \Pi\{X_\alpha : \alpha \in \mathcal{A}\}$ denote the product space, where $\mathcal{A}$ is nonempty.

LEMMA 4.6 (see [8]). *Let n be a positive integer and $A = \Pi_{j=1}^n A_{\alpha_j} \times \Pi_{\alpha \neq \alpha_j} X_\alpha$.*

(1) *$A \in \text{PO}(X)$ if and only if $A_{\alpha_j} \in \text{PO}(X_{\alpha_j})$ for each $j = 1, 2, \dots, n$.*

(2) *$\text{pCl}(\Pi_{\alpha \in \mathcal{A}} A_\alpha) \subset \Pi_{\alpha \in \mathcal{A}} \text{pCl}(A_\alpha)$.*

THEOREM 4.7. *If a function $f_\alpha : X_\alpha \rightarrow Y_\alpha$ is $\theta.p.c.$ for each $\alpha \in \mathcal{A}$. Then the product function $f : \Pi X_\alpha \rightarrow \Pi Y_\alpha$, defined by $f(\{x_\alpha\}) = \{f_\alpha(x_\alpha)\}$ for each $x = \{x_\alpha\}$, is $\theta.p.c.$*

PROOF. Let $x = \{x_\alpha\} \in \Pi X_\alpha$ and W be any open set of ΠY_α containing $f(x)$. Then, there exists an open set V_{α_j} of Y_{α_j} such that

$$f(x) = \{f_\alpha(x_\alpha)\} \in \Pi_{j=1}^n V_{\alpha_j} \times \Pi_{\alpha \neq \alpha_j} Y_\alpha \subset W. \quad (4.2)$$

Since f_α is $\theta.p.c.$ for each α , there exists $U_{\alpha_j} \in \text{PO}(X_{\alpha_j}, x_{\alpha_j})$ such that $f_{\alpha_j}(\text{pCl}(U_{\alpha_j})) \subset \text{Cl}(V_{\alpha_j})$ for $j = 1, 2, \dots, n$. Now, put $U = \Pi_{j=1}^n U_{\alpha_j} \times \Pi_{\alpha \neq \alpha_j} X_\alpha$. Then, it follows from Lemma 4.6 that $U \in \text{PO}(\Pi X_\alpha, x)$. Moreover, we have

$$\begin{aligned} f(\text{pCl}(U)) &\subset f(\Pi_{j=1}^n \text{pCl}(U_{\alpha_j}) \times \Pi_{\alpha \neq \alpha_j} X_\alpha) \\ &\subset \Pi_{j=1}^n f_{\alpha_j}(\text{pCl}(U_{\alpha_j})) \times \Pi_{\alpha \neq \alpha_j} Y_\alpha \\ &\subset \Pi_{j=1}^n \text{Cl}(V_{\alpha_j}) \times \Pi_{\alpha \neq \alpha_j} Y_\alpha \subset \text{Cl}(W). \end{aligned} \quad (4.3)$$

This shows that f is $\theta.p.c.$ $\square$

5. Preservation property

DEFINITION 5.1. A topological space X is said to be

- (1) *p -closed* [7] (resp., *p -Lindelöf*) if every cover of X by preopen sets has a finite (resp., countable) subfamily whose preclosures cover X ,
- (2) *countably p -closed* if every countable cover of X by preopen sets has a finite subfamily whose preclosures cover X ;
- (3) *quasi H -closed* [25] (resp., *almost Lindelöf* [6]) if every cover of X by open sets has a finite (resp., countable) subfamily whose closures cover X ,
- (4) *lightly compact* [5] if every countable cover of X by open sets has a finite subfamily whose closures cover X .

DEFINITION 5.2. A subset K of a space X is said to be

- (1) *p-closed relative to X* [7] if for every cover $\{V_\alpha : \alpha \in \mathcal{A}\}$ of K by preopen sets of X , there exists a finite subset $\mathcal{A}_*$ of $\mathcal{A}$ such that $K \subset \cup \{pCl(V_\alpha) : \alpha \in \mathcal{A}_*\}$,
- (2) *quasi H-closed relative to X* [25] if for every cover $\{V_\alpha : \alpha \in \mathcal{A}\}$ of K by open sets of X , there exists a finite subset $\mathcal{A}_*$ of $\mathcal{A}$ such that $K \subset \cup \{Cl(V_\alpha) : \alpha \in \mathcal{A}_*\}$.

THEOREM 5.3. If $f : X \rightarrow Y$ is a $\theta.p.c.$ function and K is *p-closed relative to X* , then $f(K)$ is *quasi H-closed relative to Y* .

PROOF. Suppose that $f : X \rightarrow Y$ is $\theta.p.c.$ and K is *p-closed relative to X* . Let $\{V_\alpha : \alpha \in \mathcal{A}\}$ be a cover of $f(K)$ by open sets of Y . For each point $x \in K$, there exists $\alpha(x) \in \mathcal{A}$ such that $f(x) \in V_{\alpha(x)}$. Since f is $\theta.p.c.$, there exists $U_x \in PO(X, x)$ such that $f(pCl(U_x)) \subset Cl(V_{\alpha(x)})$. The family $\{U_x : x \in K\}$ is a cover of K by preopen sets of X and hence there exists a finite subset K_* of K such that $K \subset \cup_{x \in K_*} pCl(U_x)$. Therefore, we obtain $f(K) \subset \cup_{x \in K_*} Cl(V_{\alpha(x)})$. This shows that $f(K)$ is *quasi H-closed relative to Y* . $\square$

COROLLARY 5.4. Let $f : X \rightarrow Y$ be a $\theta.p.c.$ surjection. Then, the following properties hold:

- (1) If X is *p-closed*, then Y is *quasi H-closed*.
- (2) If X is *p-Lindelöf*, then Y is *almost Lindelöf*.
- (3) If X is *countably p-closed*, then Y is *lightly compact*.

A subset S of a topological space X is said to be *β -open* [1] or *semipreopen* [3] if $S \subset Cl(Int(Cl(S)))$. It is well known that α -openness implies both preopenness and semi-openness which imply β -openness. The complement of a semipreopen set is said to be *semipreclosed* [3]. The intersection of all semipreclosed sets of X containing a subset S is the *semipreclosure* of S and is denoted by $spCl(S)$ [3].

DEFINITION 5.5. A topological space X is said to be

- (1) *β -connected* [24] or *semipreconnected* [2] if X cannot be expressed as the union of two nonempty disjoint β -open sets,
- (2) *semi-connected* [22] if X cannot be expressed as the union of two nonempty disjoint semi-open sets.

REMARK 5.6. We have the following implications:

$$\beta\text{-connected} \Rightarrow \text{semi-connected} \Rightarrow \text{connected}. \quad (5.1)$$

But, the converses need not be true as the following simple examples show.

EXAMPLE 5.7. (1) Let $X = \{a, b, c\}$ and $\tau = \{X, \emptyset, \{a\}, \{b\}, \{a, b\}\}$. Then (X, τ) is connected but not semi-connected.

(2) Let $X = \{a, b, c\}$ and $\tau = \{X, \emptyset, \{b, c\}\}$. Then (X, τ) is semi-connected but not β -connected.

LEMMA 5.8. For a topological space X , the following properties are equivalent:

- (1) X is *β -connected* or *semipreconnected*.
- (2) The intersection of two nonempty semipreopen subsets of X is always nonempty.
- (3) The intersection of two nonempty preopen subsets of X is always nonempty.

(4) $\text{pCl}(V) = X$ for every nonempty preopen subset V of X .

(5) $\text{spCl}(V) = X$ for every nonempty semipreopen subset V of X .

PROOF. The proofs of equivalences of (1), (2), and (3) are given in [2, Theorem 6.4]. The other properties (4) and (5), which are stated in [18], are easily equivalent to (3) and (2), respectively. $\square$

THEOREM 5.9. *If $f : X \rightarrow Y$ is a θ .p.c. surjection and X is β -connected, then Y is semi-connected.*

PROOF. Let V be any nonempty open set of Y . Let $y \in V$. Since f is surjective, there exists $x \in X$ such that $f(x) = y$. Since f is θ .p.c., there exists $U \in \text{PO}(X, x)$ such that $f(\text{pCl}(U)) \subset \text{Cl}(V)$. Since X is β -connected, by Lemma 5.8 $\text{pCl}(U) = X$ and hence $\text{Cl}(V) = Y$ since f is surjective. Therefore, it follows from [22, Theorem 4.3] that Y is semi-connected. $\square$

REMARK 5.10. The following example shows that the image of β -connectedness under weakly precontinuous surjections is not necessarily semi-connected.

EXAMPLE 5.11. Let X be the set of real numbers, $\tau = \{\emptyset\} \cup \{V \subset X : 0 \in V\}$, $Y = \{a, b, c\}$, and $\sigma = \{Y, \emptyset, \{a\}, \{b\}, \{a, b\}\}$. Define a function $f : (X, \tau) \rightarrow (Y, \sigma)$ as follows: $f(x) = a$ if $x < 0$; $f(x) = c$ if $x = 0$; $f(x) = b$ if $x > 0$. Then f is a weakly precontinuous surjection which is not θ .p.c. The topological space (X, τ) is β -connected by Lemma 5.8. By Example 5.7(1), (Y, σ) is connected but not semi-connected.

REFERENCES

- [1] M. E. Abd El-Monsef, S. N. El-Deeb, and R. A. Mahmoud, β -open sets and β -continuous mapping, *Bull. Fac. Sci. Assiut Univ. A* **12** (1983), no. 1, 77–90. [MR 87b:54002](#). [Zbl 577.54008](#).
- [2] T. Aho and T. Nieminen, *Spaces in which preopen subsets are semiopen*, *Ricerche Mat.* **43** (1994), no. 1, 45–59. [MR 96b:54058](#). [Zbl 912.54030](#).
- [3] D. Andrijević, *Semipreopen sets*, *Mat. Vesnik* **38** (1986), no. 1, 24–32. [MR 87j:54002](#). [Zbl 604.54002](#).
- [4] S. P. Arya and M. Deb, *On θ -continuous mappings*, *Math. Student* **42** (1974), 81–89. [MR 53#3982](#).
- [5] R. W. Bagley, E. H. Connell, and J. D. McKnight Jr., *On properties characterizing pseudo-compact spaces*, *Proc. Amer. Math. Soc.* **9** (1958), 500–506. [MR 20#3523](#). [Zbl 089.17601](#).
- [6] K. Dłaska and M. Ganster, *Almost rc -Lindelöf spaces*, *Bull. Malaysian Math. Soc.* (2) **17** (1997), 51–56.
- [7] J. Dontchev, M. Ganster, and T. Noiri, *On p -closed spaces*, *Int. J. Math. Math. Sci.* **24** (2000), no. 3, 203–212. [MR 2001c:54017](#). [Zbl 0965.54024](#).
- [8] N. El-Deeb, I. A. Hasanein, A. S. Mashhour, and T. Noiri, *On p -regular spaces*, *Bull. Math. Soc. Sci. Math. R. S. Roumanie (N.S.)* **27(75)** (1983), no. 4, 311–315. [MR 85d:54018](#). [Zbl 524.54016](#).
- [9] Z. Frolík, *Remarks concerning the invariance of Baire spaces under mappings*, *Czechoslovak Math. J.* **11(86)** (1961), 381–385. [MR 24#A2932](#). [Zbl 104.17204](#).
- [10] T. Husain, *Almost continuous mappings*, *Prace Mat.* **10** (1966), 1–7. [MR 36#3322](#). [Zbl 138.17601](#).
- [11] S. Jafari and T. Noiri, *On almost precontinuous functions*, *Int. J. Math. Math. Sci.* **24** (2000), no. 3, 193–201. [CMP 1 776 323](#). [Zbl 958.54011](#).

- [12] D. S. Janković, *θ -regular spaces*, Int. J. Math. Math. Sci. **8** (1985), no. 3, 615–619. [MR 87h:54030](#). [Zbl 577.54012](#).
- [13] N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly **70** (1963), 36–41. [MR 29#4025](#). [Zbl 113.16304](#).
- [14] A. S. Mashhour, M. E. Abd El-Monsef, and S. N. El-Deep, *On precontinuous and weak precontinuous mappings*, Proc. Math. Phys. Soc. Egypt **53** (1982), 47–53. [Zbl 571.54011](#).
- [15] A. S. Mashhour, I. A. Hasanein, and S. N. El-Deeb, *A note on semicontinuity and precontinuity*, Indian J. Pure Appl. Math. **13** (1982), no. 10, 1119–1123. [MR 84a:54021](#). [Zbl 499.54009](#).
- [16] A. A. Nasef and T. Noiri, *Some weak forms of almost continuity*, Acta Math. Hungar. **74** (1997), no. 3, 211–219. [CMP 1 440 246](#). [Zbl 924.54017](#).
- [17] O. Njåstad, *On some classes of nearly open sets*, Pacific J. Math. **15** (1965), 961–970. [MR 33#3245](#). [Zbl 137.41903](#).
- [18] T. Noiri, *Properties of hyperconnected spaces*, Acta Math. Hungar. **66** (1995), no. 1-2, 147–154. [MR 95k:54037](#). [Zbl 818.54020](#).
- [19] ———, *Strongly θ -precontinuous functions*, Acta Math. Hungar. **90** (2001), 315–324.
- [20] M. C. Pal and P. Bhattacharyya, *Feeble and strong forms of preirresolute functions*, Bull. Malaysian Math. Soc. (2) **19** (1996), no. 2, 63–75. [CMP 1 464 562](#). [Zbl 885.54010](#).
- [21] R. Paul and P. Bhattacharyya, *Quasi-pre-continuous functions*, J. Indian Acad. Math. **14** (1992), no. 2, 115–126. [CMP 1 261 772](#). [Zbl 871.54018](#).
- [22] V. Pipitone and G. Russo, *Spazi semiconnessi e spazi semiaperti*, Rend. Circ. Mat. Palermo (2) **24** (1975), no. 3, 273–285 (Italian). [MR 57#4084](#).
- [23] V. Popa and T. Noiri, *Almost weakly continuous functions*, Demonstratio Math. **25** (1992), no. 1-2, 241–251. [MR 93f:54020](#). [Zbl 789.54014](#).
- [24] ———, *Weakly β -continuous functions*, An. Univ. Timișoara Ser. Mat.-Inform. **32** (1994), no. 2, 83–92. [MR 97g:54019](#). [Zbl 864.54009](#).
- [25] J. Porter and J. Thomas, *On H -closed and minimal Hausdorff spaces*, Trans. Amer. Math. Soc. **138** (1969), 159–170. [MR 38#6544](#). [Zbl 175.49501](#).
- [26] V. Pták, *Completeness and the open mapping theorem*, Bull. Soc. Math. France **86** (1958), 41–74. [MR 21#4345](#). [Zbl 082.32502](#).
- [27] N. V. Veličko, *H -closed topological spaces*, Amer. Math. Soc. Transl. (2) **78** (1968), 103–118.

TAKASHI NOIRI: DEPARTMENT OF MATHEMATICS, YATSUSHIRO COLLEGE OF TECHNOLOGY,
YATSUSHIRO, KUMAMOTO, 866-8501, JAPAN

E-mail address: noiri@as.yatsushiro-nct.ac.jp