

## ON SOME CLASSES OF BCH-ALGEBRAS

MUHAMMAD ANWAR CHAUDHRY and HAFIZ FAKHAR-UD-DIN

(Received 22 September 1999)

**ABSTRACT.** The concept of a BCH-algebra is a generalization of the concept of a BCI-algebra. It is shown that weakly commutative BCH-algebras are weakly commutative BCI-algebras. Moreover, the concepts of weakly positive implicative and weakly implicative BCH-algebras are defined and it is shown that every weakly implicative BCH-algebra is a weakly positive implicative BCH-algebra. The weakly positive implicative BCH-algebras are characterized with the help of their self maps. Two open problems are posed.

2000 Mathematics Subject Classification. Primary 06F35, 03G25.

**1. Introduction.** In 1966, Imai and Iséki introduced two classes of abstract algebras, BCK-algebras and BCI-algebras [6, 7]. BCI-algebras are a generalization of BCK-algebras. These algebras have been extensively studied since their introduction. In 1983, Hu and Li [4, 5] introduced the notion of a BCH-algebra, which is a generalization of the notions of BCK- and BCI-algebras. They have studied a few properties of these algebras. Certain other properties have been studied by Chaudhry [2] and Dudek and Thomys [3]. It has been shown [3, 4, 5] that there are no proper associative and medial BCH-algebras, that is, associative and medial BCH-algebras are associative and medial BCI-algebras, respectively.

The purpose of this paper is to investigate the existence of certain classes of proper BCH-algebras and study their properties. It is shown that proper weakly commutative BCH-algebras do not exist. However, proper weakly positive implicative and proper weakly implicative BCH-algebras exist and every weakly implicative BCH-algebra is a weakly positive implicative BCH-algebra but not conversely. Weakly positive implicative BCH-algebras have been characterized in terms of their self maps. The results proved in this paper are general in the sense that corresponding results for BCK-algebras and BCI-algebras become special cases.

**2. Preliminaries.** In this section, we describe certain definitions, known results, and examples that will be used in the sequel.

**DEFINITION 2.1** (see [9]). A BCI-algebra is an algebra  $(X, *, 0)$  of type  $(2, 0)$  satisfying the following conditions:

- (1)  $(x * y) * (x * z) \leq z * y$ ,
- (2)  $x * (x * y) \leq y$ ,
- (3)  $x \leq x$ ,
- (4)  $x \leq y$  and  $y \leq x$  imply  $x = y$ ,
- (5)  $x \leq 0$  implies  $x = 0$ , where  $x \leq y$  is defined by  $x * y = 0$ .

If (5) is replaced by  $0 \leq x$ , then the algebra is called a BCK-algebra. It is known that every BCK-algebra is a BCI-algebra but not conversely. Further, in a BCI-algebra the identity  $(x * y) * z = (x * z) * y$  holds [9].

**DEFINITION 2.2** (see [4]). A BCH-algebra is an algebra  $(X, *, 0)$  of type  $(2, 0)$  satisfying the following conditions:

- (3)  $x \leq x$ ,
- (4)  $x \leq y, y \leq x$  imply  $x = y$ ,
- (6)  $(x * y) * z = (x * z) * y$ , where  $x \leq y$  if and only if  $x * y = 0$ .

In any BCH-algebra, the following hold:

- (2)  $x * (x * y) \leq y$  [4],
- (5)  $x * 0 = 0$  implies  $x = 0$  [4],
- (7)  $0 * (x * y) = (0 * x) * (0 * y)$  [3],
- (8)  $x * 0 = x$  [3],
- (9)  $(x * y) * x = 0 * y$  [4],
- (10)  $x \leq y$  implies  $0 * x = 0 * y$  [2].

It is known that every BCI-algebra is a BCH-algebra but the following example shows that the converse is not true.

**EXAMPLE 2.3** (see [4]). Let  $X = \{0, 1, 2, 3\}$  in which  $*$  is defined by:

$*$	0	1	2	3
0	0	0	0	0
1	1	0	3	3
2	2	0	0	2
3	3	0	0	0

Then  $(X, *, 0)$  is a BCH-algebra but it is not a BCI-algebra because

$$(2 * 3) * (2 * 1) = 2 * 0 = 2 \not\leq 1 * 3 = 3. \tag{2.1}$$

**EXAMPLE 2.4** (see [2]). Let  $X = \{0, 1, 2, 3, 4\}$  in which  $*$  is defined by:

$*$	0	1	2	3	4
0	0	0	0	0	4
1	1	0	0	1	4
2	2	2	0	0	4
3	3	3	3	0	4
4	4	4	4	4	0

Routine calculations give that  $(X, *, 0)$  is a BCH-algebra but it is not a BCI-algebra because

$$(1 * 3) * (1 * 2) = 1 * 0 = 1 \not\leq 2 * 3 = 0. \tag{2.2}$$

In the sequel a BCH-algebra will be simply denoted by  $X$ .

**DEFINITION 2.5** (see [5]). A BCH-algebra  $X$  is called proper if it is not a BCI-algebra.

We note that BCH-algebras of Examples 2.3 and 2.4 are proper BCH-algebras.

**DEFINITION 2.6** (see [4]). A BCH/BCI-algebra  $X$  is called associative if  $(x * y) * z = x * (y * z)$ .

**DEFINITION 2.7** (see [3]). A BCH/BCI-algebra  $X$  is called medial if  $(x * y) * (z * \mu) = (x * z) * (y * \mu)$ .

In the sequel, we shall need the following result.

(11) A BCH-algebra  $X$  is proper if and only if it does not satisfy (1) (see [4]).

**3. Classification of BCH-algebras.** It is known that associative BCH-algebras are associative BCI-algebras and medial BCH-algebras are medial BCI-algebras [3, 4]. Thus a natural question arises whether there exist some interesting classes of proper BCH-algebras or not? We show that there exist proper weakly positive implicative BCH-algebras as well as weakly implicative BCH-algebras. Moreover, the class of weakly implicative BCH-algebras is a proper subclass of the class of weakly positive implicative BCH-algebras. However, weakly commutative BCH-algebras are weakly commutative BCI-algebras.

**DEFINITION 3.1** (see [8]). A BCK-algebra  $X$  is called positive implicative if  $(x * y) * z = (x * z) * (y * z)$ . It is called implicative if  $x * (y * x) = x$ . It is commutative if  $x * (x * y) = y * (y * x)$ .

It is well known that positive implicative BCI-algebras, implicative BCI-algebras and commutative BCI-algebras are positive implicative BCK-algebras, implicative BCK-algebras and commutative BCK-algebras, respectively [9].

In [1], Chaudhry defined three classes of proper BCI-algebras, namely, weakly positive implicative BCI-algebras, weakly implicative BCI-algebras, and weakly commutative BCI-algebras. He also investigated a few properties of these algebras. We recall these definitions and the following result.

**DEFINITION 3.2** (see [1]). A BCI-algebra  $X$  is called weakly positive implicative if

$$(12) \quad (x * y) * z = ((x * z) * z) * (y * z).$$

It is called weakly implicative if

$$(13) \quad (x * (y * x)) * (0 * (y * x)) = x.$$

It is called weakly commutative if

$$(14) \quad (x * (x * y)) * (0 * (x * y)) = y * (y * x).$$

**THEOREM 3.3** (see [1]). A BCI-algebra  $X$  is weakly positive implicative if and only if

$$(15) \quad x * y = ((x * y) * y) * (0 * y).$$

We note that Theorem 3.3 tells us that (12) and (15) are equivalent in a BCI-algebra. However, they are not equivalent in a BCH-algebra. We consider the BCH-algebra  $X$  of Example 2.4. Then easy calculations give that (15) is satisfied but (12) is not satisfied because  $(1 * 2) * 3 = 0 \neq 1 = ((1 * 3) * 3) * (2 * 3)$ . Further the following theorem tells us that BCH-algebras satisfying (12) are BCI-algebras.

**THEOREM 3.4.** A BCH-algebra satisfying  $(x * y) * z = ((x * z) * z) * (y * z)$  is a BCI-algebra.

**PROOF.** In view of (11) it is sufficient to prove that (1) holds.

Consider

$$\begin{aligned}
 ((x * y) * (x * z)) * (z * y) &= ((x * (x * z)) * y) * (z * y) \\
 &= (((x * y) * y) * ((x * z) * y)) * (z * y) \quad (\text{by (12)}) \\
 &= (((x * y) * y) * (z * y)) * ((x * z) * y) \\
 &= ((x * z) * y) * ((x * z) * y) \quad (\text{by (12)}) \\
 &= 0.
 \end{aligned} \tag{3.1}$$

This completes the proof.  $\square$

In view of Theorems 3.3 and 3.4 and the comments made between them, we adopt the following definitions for BCH-algebras.

**DEFINITION 3.5.** A BCH-algebra  $X$  is weakly positive implicative if

$$x * y = ((x * y) * y) * (0 * y) \quad \forall x, y \in X. \tag{3.2}$$

We note that the BCH-algebra of Example 2.4 satisfies (3.2). Thus there exist proper weakly positive implicative BCH-algebras.

**DEFINITION 3.6.** A BCH-algebra  $X$  is weakly implicative if

$$(x * (y * x)) * (0 * (y * x)) = x \quad \forall x, y \in X. \tag{3.3}$$

**DEFINITION 3.7.** A BCH-algebra  $X$  is weakly commutative if

$$(x * (x * y)) * (0 * (x * y)) = y * (y * x). \tag{3.4}$$

**THEOREM 3.8.** Every weakly implicative BCH-algebra  $X$  is a weakly positive implicative BCH-algebra.

**PROOF.** Let  $X$  be weakly implicative. Then

$$(x * (z * x)) * (0 * (z * x)) = x. \tag{3.5}$$

Putting  $x = z * x$  in (3.5), we get

$$((z * x) * (z * (z * x))) * (0 * (z * (z * x))) = z * x. \tag{3.6}$$

Since  $z * (z * x) \leq x$ , therefore (10) gives  $0 * (z * (z * x)) = 0 * x$ . Thus

$$z * x = ((z * x) * (z * (z * x))) * (0 * x) = ((z * (z * (z * x))) * x) * (0 * x). \tag{3.7}$$

Now

$$\begin{aligned}
 &(z * x) * (z * (z * (z * x))) \\
 &= ((z * (z * (z * x))) * x) * (0 * x) * (z * (z * (z * x))) \\
 &= ((z * (z * (z * x))) * x) * (z * (z * (z * x))) * (0 * x) \\
 &= (0 * x) * (0 * x) = 0.
 \end{aligned} \tag{3.8}$$

Hence  $z * x \leq z * (z * (z * x)) \leq z * x$ . Thus

$$z * (z * (z * x)) = z * x \tag{3.9}$$

holds in a weakly implicative BCH-algebra. Putting (3.9) in (3.7) we get  $z * x = ((z * x) * x) * (0 * x)$ . Hence,  $X$  is weakly positive implicative. This completes the proof.  $\square$

**REMARK 3.9.** It is known that  $0 * x = 0 * (0 * (0 * x))$  holds in a BCH-algebra [3], but it is still not known that in a BCH-algebra the identity  $x * y = x * (x * (x * y))$  holds or not, although it holds in BCI-algebras and weakly implicative BCH-algebras (as shown in (3.9)).

**REMARK 3.10.** Since every BCI-algebra is a BCH-algebra and weak positive implicativeness and weak implicativeness coincide with positive implicativeness and implicativeness, respectively, in BCK-algebras [1], therefore the following results of Chaudhry and Iséki follow as corollaries of Theorem 3.4.

**COROLLARY 3.11** (see [1]). *Every weakly implicative BCI-algebra is a weakly positive implicative BCI-algebra.*

**COROLLARY 3.12** (see [8]). *Every implicative BCK-algebra is a positive implicative BCK-algebra.*

**THEOREM 3.13.** *A BCH-algebra  $X$  satisfying  $(x * (x * y)) * (0 * (x * y)) = y * (y * x)$  is a BCI-algebra.*

**PROOF.** It is sufficient to show that (1) holds. We consider

$$\begin{aligned} & ((x * y) * (x * z)) * (z * y) \\ &= ((x * (x * z)) * y) * (z * y) \\ &= (((z * (z * x)) * (0 * (z * x))) * y) * (z * y) \text{ (by given condition)} \\ &= (((z * (z * x)) * y) * (0 * (z * x))) * (z * y) \tag{3.10} \\ &= (((z * y) * (z * x)) * (0 * (z * x))) * (z * y) \\ &= (((z * y) * (z * y)) * (z * x)) * (0 * (z * x)) \\ &= (0 * (z * x)) * (0 * (z * x)) = 0. \end{aligned}$$

This completes the proof.  $\square$

We now pose the following open problem.

**OPEN PROBLEM 1.** Do there exist classes of proper BCH-algebras other than the classes of weakly positive implicative and weakly implicative BCH-algebras, which are generalizations of the known classes of BCI- as well as BCK-algebras.

**4. Characterization of weakly positive implicative BCH-algebras.** In this section, we characterize weakly positive implicative BCH-algebras by their self maps.

**DEFINITION 4.1.** Let  $X$  be a BCH-algebra. For a fixed  $x$  in  $X$ , the map  $R_x : X \rightarrow X$  given by  $R_x(t) = t * x$  for all  $t \in X$  is called a right self map.

**DEFINITION 4.2.** Let  $X$  be a BCH-algebra. For a fixed  $x$  in  $X$ , the map  $R'_x : X \rightarrow X$  given by  $R'_x(t) = (t * x) * (0 * x)$  for all  $t \in X$  is called a weak right self map.

The following theorem gives us a characterization of a weakly positive implicative BCH-algebra with the help of its right and weak right self maps.

**THEOREM 4.3.** A BCH-algebra  $X$  is weakly positive implicative if and only if  $R_z = R'_z \circ R_z$  for all  $z \in X$ , where "o" is composition of functions.

**PROOF.** Let  $X$  be a BCH-algebra and  $R_z = R'_z \circ R_z$ . Then  $R_z(y) = R'_z \circ R_z(y)$  for all  $y \in X$ . Thus  $y * z = R'_z(y * z) = ((y * z) * z) * (0 * z)$  for all  $y, z \in X$ . Hence  $X$  is a weakly positive implicative BCH-algebra. Conversely, if  $X$  is a weakly positive implicative BCH-algebra, then  $y * z = ((y * z) * z) * (0 * z)$ . Thus  $R_z(y) = (R_z(y) * z) * (0 * z) = R'_z(R_z(y)) = R'_z \circ R_z(y)$  for all  $y, z \in X$ . Hence  $R_z = R'_z \circ R_z$ . This completes the proof. □

**THEOREM 4.4.** Let  $X$  be a weakly positive implicative BCH-algebra. Then  $R'_y = R'_y \circ R'_y = (R'_y)^2$ .

**PROOF.** Since  $X$  is weakly positive implicative, therefore  $x * y = ((x * y) * y) * (0 * y)$ . Thus

$$\begin{aligned} (x * y) * (0 * y) &= (((x * y) * y) * (0 * y)) * (0 * y) \\ &= (((x * y) * (0 * y)) * y) * (0 * y). \end{aligned} \tag{4.1}$$

Hence

$$R'_y(x) = R'_y((x * y) * (0 * y)) = R'_y(R'_y(x)) = R'_y \circ R'_y(x) = (R'_y)^2(x) \tag{4.2}$$

for all  $x, y \in X$ . This completes the proof. □

The following example shows that the converse of the above theorem is not true.

**EXAMPLE 4.5.** Let  $X = \{0, a, b, c\}$  in which  $*$  is defined by:

$*$	0	a	b	c
0	0	0	b	b
a	a	0	b	b
b	b	b	0	0
c	c	b	a	0

Then  $X$  is a BCI-algebra. Further  $X$  is not weakly positive implicative because  $a = c * b \neq ((c * b) * b) * (0 * b) = (a * b) * (0 * b) = b * b = 0$ . Moreover, easy calculations give that

$$R'_0 = (R'_0)^2, \quad R'_a = (R'_a)^2, \quad R'_b = (R'_b)^2, \quad R'_c = (R'_c)^2. \tag{4.3}$$

This shows that the converse of Theorem 4.4 does not hold for the class of BCH-algebras, because it does not hold for BCI-algebras.

We now pose another open problem.

**OPEN PROBLEM 2.** What are the characterizations of weakly positive implicative BCH-algebras and weakly implicative BCH-algebras in terms of their ideals.

**ACKNOWLEDGEMENT.** The first author gratefully acknowledges the support provided by the King Fahd University of Petroleum and Minerals.

#### REFERENCES

- [1] M. A. Chaudhry, *Weakly positive implicative and weakly implicative BCI-algebras*, Math. Japon. **35** (1990), no. 1, 141-151. MR 90k:06019. Zbl 702.06014.
- [2] ———, *On BCH-algebras*, Math. Japon. **36** (1991), no. 4, 665-676. MR 92e:06042. Zbl 733.06008.
- [3] W. A. Dudek and J. Thomys, *On decompositions of BCH-algebras*, Math. Japon. **35** (1990), no. 6, 1131-1138. MR 91j:06040. Zbl 723.06014.
- [4] Q. P. Hu and X. Li, *On BCH-algebras*, Math. Sem. Notes Kobe Univ. **11** (1983), no. 2, part 2, 313-320. MR 86a:06016. Zbl 579.03047.
- [5] ———, *On proper BCH-algebras*, Math. Japon. **30** (1985), no. 4, 659-661. MR 87d:06042. Zbl 583.03050.
- [6] Y. Imai and K. Iséki, *On axiom systems of propositional calculi. XIV*, Proc. Japan Acad. **42** (1966), 19-22. MR 33#3902. Zbl 156.24812.
- [7] K. Iséki, *An algebra related with a propositional calculus*, Proc. Japan Acad. **42** (1966), 26-29. MR 34#2433. Zbl 207.29304.
- [8] K. Iséki and S. Tanaka, *An introduction to the theory of BCK-algebras*, Math. Japon. **23** (1978/79), no. 1, 1-26. MR 80a:03081. Zbl 385.03051.
- [9] K. Iséki and A. B. Thaheem, *Note on BCI-algebras*, Math. Japon. **29** (1984), no. 2, 255-258. MR 85i:06020a. Zbl 541.03038.

MUHAMMAD ANWAR CHAUDHRY: DEPARTMENT OF MATHEMATICAL SCIENCES, KING FAHD UNIVERSITY OF PETROLEUM AND MINERALS, DHAHRAN 31261, SAUDI ARABIA  
*E-mail address:* chaudhry@kfupm.edu.sa

HAFIZ FAKHAR-UD-DIN: DEPARTMENT OF MATHEMATICS, ISLAMIA UNIVERSITY, BAHAWALPUR, PAKISTAN



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

