

THE MONAD INDUCED BY THE HOM-FUNCTOR IN THE CATEGORY OF TOPOLOGICAL SPACES AND ITS ASSOCIATED EILENBERG-MOORE ALGEBRAS

KOENA R. NAILANA

Received 15 September 2001

We discuss the monad associated with the topology of pointwise convergence. We also study examples of the Eilenberg-Moore algebras for this monad.

2000 Mathematics Subject Classification: 18C15, 54B30, 54F05.

1. Introduction. Let Top denote the category of topological spaces and continuous functions. Let \mathbb{R} denote the real line with the usual topology, and for each topological space X , let $C(X, \mathbb{R})$ be the set of continuous real-valued functions from X to \mathbb{R} . Consider the contravariant hom-functor $C_p : \text{Top} \rightarrow \text{Top}^{\text{op}}$ defined by assigning to each space X the space of continuous real-valued functions with the topology of pointwise convergence. We denote this space by $C_p X$. The space $C_p X$ has been extensively studied. A fundamental reference on $C_p X$ is Arkhangel'skii [2]. We recall that the sub-basic open sets of $C_p X$ are sets of the form $[f, V]$, where $[x, V] = \{f \in C_p X : f(x) \in V, V \text{ open in } \mathbb{R}\}$.

2. The monad induced by the hom-functor in Top and the associated M -algebras. We now consider the composite functor $C_p^{\text{op}} C_p : \text{Top} \rightarrow \text{Top}^{\text{op}} \rightarrow \text{Top}$ where C_p^{op} is the dual functor. Let $M = C_p^{\text{op}} C_p$. If $x \in X$, then the function $\hat{x} : C_p X \rightarrow \mathbb{R}$ defined by $\hat{x}(f) = f(x)$ is called the evaluation map at x . The following propositions are important since they ensure that our morphisms are continuous. The proofs are straightforward and will be omitted.

PROPOSITION 2.1. (i) For all $x \in X$, $\hat{x} : C_p X \rightarrow \mathbb{R}$ is continuous.
(ii) For all $g \in C_p X$, $\hat{g} : M C_p X \rightarrow \mathbb{R}$ is continuous.

PROPOSITION 2.2. Let X be any topological space. Then

- (i) $\eta_X : X \rightarrow M X$, where $\eta_X(x) = \hat{x}$ is continuous.
- (ii) $\mu_X : M M X \rightarrow M X$, where $\mu_X(y)[g] = y(\hat{g})$ is continuous.

We recall from [1] that a monad on a category \mathbb{A} is a triplet $\mathbb{M} = (M, \eta, \mu)$ consisting of a functor $M : \mathbb{A} \rightarrow \mathbb{A}$ and natural transformations $\eta : \text{id}_{\mathbb{A}} \rightarrow M$ and $\mu : M M \rightarrow M$ such that $\mu \circ M \mu = \mu \circ \mu M$, $\mu \circ M \eta = \text{id}$, and $\mu \circ \eta M = \text{id}$.

PROPOSITION 2.3. The triplet (M, η, μ) , where $\eta : \text{id}_{\text{Top}} \rightarrow M$ and $\mu : M M \rightarrow M$ are defined by $\eta_X(x) = \hat{x}$ and $\mu_X(y)[g] = y(\hat{g})$, respectively, where $x \in X$, $g \in C_p X$, is a monad.

PROOF. We first check that $\eta : \text{id}_{\text{Top}} \rightarrow M$ and $\mu : MM \rightarrow M$ are natural transformations. Let $f : X \rightarrow Y$ be a continuous function. We show that $M(f) \circ \eta_X = \eta_Y \circ f$. We define η_X and $M(f)$ by $\eta_X(x) = \hat{x}$ and $M(f)(y)[g] = y(g \circ f)$ where $g : Y \rightarrow \mathbb{R}$ is continuous, $y \in M(X)$, and $\hat{}$ denotes evaluation, for example, $\hat{x}(g) = g(x)$. Then $M(f) \circ \eta_X(x) = M(f)(\hat{x})$. Let $g \in C_p X$. Then $M(f)(\hat{x})[g] = \hat{x}(g \circ f) = g \circ f(x) = g(f(x)) = \widehat{f(x)}[g]$. Hence $M(f)(\hat{x}) = \widehat{f(x)}$. Now $\eta_Y \circ f(x) = \eta_Y(f(x)) = \widehat{f(x)}$. Let $g \in C_p X$. Then $\widehat{f(x)}[g] = g(f(x))$. Hence $M(f) \circ \eta_X = \eta_Y \circ f$. We define μ_X by $\mu_X(y)[g] = y(\hat{g})$ where $y \in MM(X)$, $g \in C_p X$, and \hat{g} denotes the evaluation function at g , that is, $\hat{g} : M(X) \rightarrow \mathbb{R}$. We now show that $\mu : MM \rightarrow M$ is a natural transformation, that is, $M(f) \circ \mu_X = \mu_Y \circ M^2(f)$. Let $h \in C_p Y$. Then

$$(M(f) \circ \mu_X)(y)[h] = M(f)(\mu_X(y))[h] = \mu_X(y)(h \circ f) = y(\widehat{h \circ f}). \quad (2.1)$$

On the other hand,

$$\begin{aligned} \mu_Y \circ M^2(f)(y)[h] &= \mu_Y(M^2(f)(y))[h] \\ &= M^2(f)(y)(\hat{h}) \\ &= M(M(f))(y)(\hat{h}) \\ &= y(\hat{h} \circ M(f)). \end{aligned} \quad (2.2)$$

Let $\lambda : C_p X \rightarrow \mathbb{R}$. Then $(\hat{h} \circ M(f))(\lambda) = \hat{h}(M(f)(\lambda)) = M(f)(\lambda)[h] = \lambda(h \circ f) = \widehat{h \circ f}(\lambda)$. Therefore, $\hat{h} \circ M(f) = \widehat{h \circ f}$. From the equations

$$\begin{aligned} M(f) \circ \mu_X(y)[h] &= y(\widehat{h \circ f}), \\ \mu_Y \circ M^2(f)(y)[h] &= y(\hat{h} \circ M(f)), \\ (\hat{h} \circ M(f))(\lambda) &= \widehat{h \circ f}(\lambda), \end{aligned} \quad (2.3)$$

we get $M(f) \circ \mu_X = \mu_Y \circ M^2(f)$. Therefore $\mu : MM \rightarrow M$ is a natural transformation. We now show that the other monad conditions are satisfied. First, we show that $\mu_X \circ M\eta = \text{id}$. We prove that $\mu_X \circ M\eta = \text{id}$ and $\mu_X \circ \eta_M = \text{id}$. Let $y \in M(X)$ and $f \in C_p X$. Then $\hat{f} : M(X) \rightarrow \mathbb{R}$ and $(\mu_X \circ M\eta)(y) \in M(X)$. Then $(\mu_X \circ M\eta)(y)[f] = \mu_X(M\eta(y))[f] = M\eta(y)(\hat{f}) = y(\hat{f} \circ \eta) = y[f]$. Therefore $\mu_X \circ M\eta = \text{id}$. On the other hand $(\mu_X \circ \eta_M)(y)[f] = \mu_X(\eta_M(y))[f] = \eta_M(y)(\hat{f}) = \hat{y}(\hat{f}) = \hat{f}(y) = y(f)$. Therefore $\mu_X \circ \eta_M = \text{id}$. Second, we show that $\mu_X \circ \mu_M = \mu_X \circ M\mu_X$. We prove that $\mu_X \circ \mu_M = \mu_X \circ M\mu_X$. Let $y \in MMM(X)$. Then $\mu_X \circ M\mu_X(y) \in M(X)$. Let $f \in C_p X$. Then $(\mu_X \circ M\mu_X)(y)[f] = \mu_X(M\mu_X(y))[f] = M\mu_X(y)(\hat{f}) = y(\hat{f} \circ \mu_X) = y(\hat{\hat{f}})$. On the other hand, $(\mu_X \circ \mu_M)(y)[f] = \mu_X(\mu_M(y))[f] = \mu_M(y)(\hat{f}) = y(\hat{f})$. Therefore $\mu_X \circ \mu_M = \mu_X \circ M\mu_X$. Therefore (M, η, μ) is a monad. \square

If $\mathbb{M} = (M, \eta, \mu)$ is a monad on \mathbb{A} , then (A, h_A) is called an Eilenberg-Moore algebra or simply an M -algebra if the algebra map $h_A : MA \rightarrow A$ satisfies $h_A \circ \eta_A = \text{id}_A$ and $h_A \circ Mh_A = h_A \circ \mu_A$.

We now look at examples of the M -algebras of the monad (M, η, μ) .

PROPOSITION 2.4. *The real line \mathbb{R} is an M -algebra.*

PROOF. We define $h_{\mathbb{R}} : M\mathbb{R} \rightarrow \mathbb{R}$ as $\hat{\mathbf{1}}_{\mathbb{R}}$, that is, the identity map with respect to \mathbb{R} , and show that the M -algebra conditions are satisfied. It is obvious that the map $h_{\mathbb{R}}$ is continuous. Let $x \in \mathbb{R}$. Then $h_{\mathbb{R}} \circ \eta_{\mathbb{R}}(x) = h_{\mathbb{R}}(\hat{x}) = \hat{x}(\mathbf{1}_{\mathbb{R}}) = \mathbf{1}_{\mathbb{R}}(x)$. Therefore $h_{\mathbb{R}} \circ \eta_{\mathbb{R}} = \mathbf{1}_{\mathbb{R}}$. Now let $y \in MM(\mathbb{R})$. Then $h_{\mathbb{R}} \circ \mu_{\mathbb{R}}(y) = \hat{\mathbf{1}}_{\mathbb{R}}(\mu_{\mathbb{R}}(y)) = \mu_{\mathbb{R}}(y)(\mathbf{1}_{\mathbb{R}}) = y(\hat{\mathbf{1}}_{\mathbb{R}})$. On the other hand, $h_{\mathbb{R}} \circ Mh_{\mathbb{R}}(y) = \hat{\mathbf{1}}_{\mathbb{R}}(M\hat{\mathbf{1}}_{\mathbb{R}}(y)) = \hat{\mathbf{1}}_{\mathbb{R}}(y \circ C_p(\hat{\mathbf{1}}_{\mathbb{R}})) = y \circ C_p(\hat{\mathbf{1}}_{\mathbb{R}})(\mathbf{1}_{\mathbb{R}}) = y(C_p(\hat{\mathbf{1}}_{\mathbb{R}})(\mathbf{1}_{\mathbb{R}})) = y(\mathbf{1}_{\mathbb{R}} \circ \hat{\mathbf{1}}_{\mathbb{R}}) = y(\hat{\mathbf{1}}_{\mathbb{R}})$. Therefore $h_{\mathbb{R}} \circ \mu_{\mathbb{R}} = h_{\mathbb{R}} \circ Mh_{\mathbb{R}}$. \square

PROPOSITION 2.5. *For each $X \in \text{Top}$, $C_p X$ is an M -algebra with $h_{C_p X} = C_p(\eta_X)$.*

PROOF. We first define $h_{C_p X} : MC_p X \rightarrow C_p X$. Let $\varphi \in MC_p X$. We define $h_{C_p X}$ by $h_{C_p X}(\varphi) = \varphi \circ \eta_X = C_p \eta_X(\varphi)$. Then the map $h_{C_p X}$ is continuous, since it is the composite of continuous functions φ and η_X . We now show that the conditions for an M -algebra are satisfied. Thus, we must show that $h_{C_p X} \circ \eta_{C_p X} = \text{id}_{C_p X}$. Let $f \in C_p X$. Then $h_{C_p X} \circ \eta_{C_p X}(f) = h_{C_p X}(\eta_{C_p X}(f)) = C_p \eta_X(\hat{f}) = \hat{f} \circ \eta_X = f = \text{id}_{C_p X}(f)$, since $\hat{f} \circ \eta_X(x) = \hat{f}(\eta_X(x)) = \hat{x}(f) = f(x)$. Therefore $h_{C_p X} \circ \eta_{C_p X} = \text{id}_{C_p X}$.

We must now show that $h_{C_p X} \circ \mu_{C_p X} = h_{C_p X} \circ Mh_{C_p X}$. Let $y \in MMC_p X$. Then $h_{C_p X} \circ \mu_{C_p X}(y) = h_{C_p X}(\mu_{C_p X}(y)) = C_p \eta_X(\mu_{C_p X}(y)) = \mu_{C_p X}(y) \circ \eta_X$. Now let $x \in X$. Then $\mu_{C_p X}(y) \circ \eta_X(x) = \mu_{C_p X}(y)(\hat{x}) = y(\hat{\hat{x}})$. On the other hand, $h_{C_p X} \circ Mh_{C_p X}(y) = C_p \eta_X \circ MC_p \eta_X(y) = C_p(M\eta_X \circ \eta_X)(y) = y \circ M\eta_X \circ \eta_X$. Let $x \in X$. Then $M\eta_X \circ \eta_X(x) = M\eta_X(\hat{x}) = \hat{\hat{x}} \circ C_p \eta_X = \hat{\hat{x}}$. Therefore $h_{C_p X} \circ \mu_{C_p X} = h_{C_p X} \circ Mh_{C_p X}$. Hence $C_p X$ is an M -algebra. \square

PROPOSITION 2.6. *Retracts of $C_p X$ are M -algebras.*

PROOF. Let $g : C_p Y \rightarrow X$ be a retraction. Then there is a continuous function $f : X \rightarrow C_p Y$ such that $g \circ f = \text{id}_X$. The following diagram will help us define the algebra map $h_X : MX \rightarrow X$:

$$\begin{array}{ccccc}
 X & \xrightarrow{\eta_X} & MX & \xrightarrow{Mf} & MC_p Y & \xrightarrow{C_p \eta_Y} & C_p Y \\
 & & \searrow \text{id}_{MX} & & \downarrow Mg & & \downarrow g \\
 & & & & MX & & \\
 & & \searrow \text{id}_X & & \swarrow h_X & & \\
 & & & & & & X
 \end{array}
 \tag{2.4}$$

Define

$$h_X = g \circ C_p \eta \circ Mf = g \circ C_p(C_p f \circ \eta_Y).
 \tag{2.5}$$

Since h_X is the composite of continuous functions, then it is continuous.

Now,

$$\begin{aligned} h_X \circ \eta_X(x) &= h_X(\hat{x}) = g(C(Cf \circ \eta_Y)(\hat{x})) = g(\hat{x} \circ Cf \circ \eta_Y) \\ &= g(\widehat{f(x)} \circ \eta_Y) = g(f(x)) = \text{id}_X(x), \end{aligned} \quad (2.6)$$

since g is a retraction.

We now show that $h_X \circ \mu_X = h_X \circ Mh_X$. Let $y \in MMX$. Then

$$\begin{aligned} h_X \circ \mu_X(y) &= h_X(\mu_X(y)) = g \circ C_p(C_p f \circ \eta_Y)(\mu_X(y)) \\ &= g(C_p(C_p f \circ \eta_Y)(\mu_X(y))) = g(\mu_X(y) \circ C_p f \circ \eta_Y). \end{aligned} \quad (2.7)$$

If $k \in C_p X$, then

$$\mu_X(y)(k) = y(\hat{k}). \quad (2.8)$$

On the other hand,

$$\begin{aligned} h_X \circ Mh_X &= g \circ C_p(C_p f \circ \eta_Y) \circ M(g \circ C_p(C_p f \circ \eta_Y)) \\ &= g \circ C_p \eta_Y \circ Mf \circ Mg \circ MC_p(C_p f \circ \eta_Y) \\ &= g \circ C_p \eta_Y \circ M(f \circ g) \circ MC_p(C_p f \circ \eta_Y) \\ &= g \circ C_p \eta_Y \circ M(\text{id}_X) \circ MC_p(C_p f \circ \eta_Y) \\ &= g \circ C_p \eta_Y \circ MC_p(C_p f \circ \eta_Y) \\ &= g \circ C_p(M(C_p f \circ \eta_Y) \circ \eta_Y) \\ &= g \circ C_p(\eta_{C_p X} \circ C_p f \circ \eta_Y). \end{aligned} \quad (2.9)$$

Now,

$$\begin{aligned} h_X \circ Mh_X(y) &= g(C_p(\eta_{C_p X} \circ C_p f \circ \eta_Y)(y)) \\ &= g(y \circ \eta_{C_p X} \circ C_p f \circ \eta_Y). \end{aligned} \quad (2.10)$$

We only need to show that $y \circ \eta_{C_p X} = \mu_X$. Let $k \in C_p X$. Then $y \circ \eta_{C_p X}(k) = y(\eta_{C_p X}(k)) = y(\hat{k})$. From (2.8), we have $y \circ \eta_{C_p X} = \mu_X$ and therefore $h_X \circ \mu_X = h_X \circ Mh_X$. Hence retracts of $C_p X$ are M -algebras. \square

3. The algebra morphisms and the transfer of ring structure from MX to X for an M -algebra (X, h_X) . For an M -algebra (X, h_X) the ring structure on MX can be transferred to X , via h_X , in such a way that X becomes a ring with respect to the induced operations.

DEFINITION 3.1. On an M -algebra (X, h_X) define

- (i) $x_1 + x_2$ to be $h_X(\eta_X(x_1) + \eta_X(x_2))$,
- (ii) $x_1 \cdot x_2$ to be $h_X(\eta_X(x_1) \cdot \eta_X(x_2))$.

In addition to the ring structure defined above we also define the scalar multiplication in the following way: define tx to be $h_X(t\eta_X(x))$, where t is a scalar.

According to [Definition 3.1](#), $C_p X$ (being an M -algebra, [Proposition 2.5](#)) has now two concepts of the operations “+” and “ \cdot ”, the natural one defined pointwise

$$(h_X(x + y) = h_X(x) + h_X(y), h_X(xy) = h_X(x)h_X(y)) \quad (3.1)$$

and Definition 3.1. The same applies to MX . We omit the straightforward proof of the following proposition.

PROPOSITION 3.2. *The natural operations on MX defined pointwise coincide with the corresponding ones defined above.*

LEMMA 3.3. *The topology on X is initial with respect to η_X , that is, X has the weak topology induced by η_X into $C_p C_p X = MX$.*

PROOF. Basic neighborhoods of $\eta_X(x)$ have inverse images of η_X of the form $\cap_{i=1}^n f_i^{-1}[W_i]$. □

LEMMA 3.4 [2]. *Let $\varphi \in C_p C_p X$ such that $\varphi : (C_p X, h_{C_p X}) \rightarrow (\mathbb{R}, h_{\mathbb{R}})$ is a linear functional. Then there are $x_1, \dots, x_n \in X, \lambda_1, \dots, \lambda_n \in \mathbb{R}$ such that $\varphi = \sum_{i=1}^n \lambda_i \hat{x}_i$.*

PROPOSITION 3.5. *If $\varphi : (C_p X, h_{C_p X}) \rightarrow (\mathbb{R}, h_{\mathbb{R}})$ is a nontrivial continuous multiplicative linear functional, then there is $x \in X$ such that $\varphi = \hat{x}$, that is, φ is a point evaluation.*

PROOF. By Lemma 3.4, there are points $x_1, \dots, x_n \in X$, and scalars $\lambda_1, \dots, \lambda_n \in \mathbb{R}$ such that $\varphi = \sum_{i=1}^n \lambda_i \hat{x}_i$ where $\lambda_i = \varphi(g_i), g_i \in C_p X$ being such that $g_i(x_i) = 1, g_i(x_j) = 0$ for $i \neq j, 0 \leq g_i \leq 1$. Now $\varphi(g_k^2) = \varphi(g_k)^2 = \lambda_k$. Also $\varphi(g_k^2) = \sum_{i=1}^n \lambda_i \hat{x}_i(g_k^2) = \sum_{i=1}^n \lambda_i g_k^2(x_i) = \lambda_k$.

Thus $\lambda_k = \lambda_k^2$, so that $\lambda_k = 0$ or $\lambda_k = 1$ for $k = 1, 2, \dots, n$. Moreover, $\lambda_k = g_k(x_k) \geq 0$. Furthermore, $\varphi(\mathbf{1}) = 1$ gives $1 = \varphi(\mathbf{1}) = \sum_{i=1}^n \lambda_i \hat{x}_i(\mathbf{1}) = \sum_{i=1}^n \lambda_i$. Consequently, all λ_i 's except one are zero, the exceptional one being one 1. Let $x = x_l$, where $\lambda_l = 1$. Then $\lambda_i = 0$ for $i \neq l$, so that $\varphi = \lambda_l \hat{x}_l = \hat{x}_l$. □

PROPOSITION 3.6. *Let $\varphi : (C_p X, h_{C_p X}) \rightarrow (\mathbb{R}, h_{\mathbb{R}})$ be an algebra map. Then φ is a continuous ring homomorphism.*

PROOF. Given $f, g \in C_p X$, consider $\eta_{C_p X}(f) + \eta_{C_p X}(g)$ in $MC_p X$. We have

$$h_{\mathbb{R}} \circ C^2 \varphi(\eta_{C_p X}(f) + \eta_{C_p X}(g)) = h_{\mathbb{R}} \circ M\varphi(\eta_{C_p X}(f)) + h_{\mathbb{R}} \circ M\varphi(\eta_{C_p X}(g)) \tag{3.2}$$

by Lemma 3.4.

Hence $\varphi \circ h_X(\eta_{C_p X}(f) + \eta_{C_p X}(g)) = \varphi \circ h_X(\eta_{C_p X}(f)) + \varphi \circ h_X(\eta_{C_p X}(g))$, so that $\varphi(f + g) = \varphi(f) + \varphi(g)$, since h_X preserves the ring structure. Similarly, $\varphi(f \cdot g) = \varphi(f) \cdot \varphi(g)$. We also have $\varphi(tf) = t\varphi(f), t \in \mathbb{R}$. Moreover $\varphi(\mathbf{1}) = 1$, where $\mathbf{1}$ denotes the constant function with value equal to 1. □

PROPOSITION 3.7. *Every algebra map $\varphi : (C_p X, h_{C_p X}) \rightarrow (\mathbb{R}, h_{\mathbb{R}})$ is a point evaluation map.*

PROOF. By the above proposition, φ is a continuous ring homomorphism, that is, a continuous multiplicative linear functional. Thus, there is some $x \in X$ such that $\varphi(f) = f(x)$ for all f in $C_p X$, by the above proposition. □

THEOREM 3.8. *The algebra morphisms $\varphi : (C_p X, h_{C_p X}) \rightarrow (\mathbb{R}, h_{\mathbb{R}})$ are precisely the morphisms \hat{x} , where $x \in X$, that is, the point evaluation map.*

PROOF. Suppose $\varphi = \hat{x}$, for some $x \in X$. Let $y \in MC_p X$. Take $h_X = C_p \eta_{C_p X}$ and $h_{\mathbb{R}} = \hat{1}_{\mathbb{R}}$. Then $\varphi \circ h_X(y) = \hat{x} \circ C_p \eta_{C_p X}(y) = \hat{x}(C_p \eta_{C_p X}(y)) = y \circ \eta_X(x) = y(\hat{x})$. On the other hand, $h_{\mathbb{R}} \circ M(\varphi)(y) = h_{\mathbb{R}}(M(\varphi)(y)) = \hat{1}_{\mathbb{R}}(M(\varphi)(y)) = M(\varphi)(y)(1_{\mathbb{R}}) = y(1_{\mathbb{R}} \circ \varphi) = y(\varphi) = y(\hat{x})$. Therefore, $\varphi \circ h_X = h_{\mathbb{R}} \circ M(\varphi)$ and thus $\varphi = \hat{x}$ is an algebra morphism. The converse follows from [Proposition 3.7](#). \square

PROPOSITION 3.9. *The algebra morphisms $\varphi : (C_p X, h_{C_p X}) \rightarrow (C_p Y, h_{C_p Y})$ are the maps $C_p(f)$, where $f : Y \rightarrow X$ is continuous.*

PROOF. Suppose that $\varphi : (C_p X, h_{C_p X}) \rightarrow (C_p Y, h_{C_p Y})$ is an algebra map. Given $y \in Y$, $\hat{y} \circ \varphi : (C_p X, h_{C_p X}) \rightarrow (\mathbb{R}, h_{\mathbb{R}})$ is an algebra map, since the composition of two algebra maps is an algebra map. Thus the following diagram is commutative:

$$\begin{array}{ccccc}
 MC_p X & \xrightarrow{M\varphi} & MC_p(Y) & \xrightarrow{M\hat{y}} & M\mathbb{R} \\
 \downarrow h_{C_p X} & & \downarrow h_{C_p Y} & & \downarrow h_{\mathbb{R}} \\
 C_p X & \xrightarrow{\varphi} & C_p Y & \xrightarrow{\hat{y}} & \mathbb{R}
 \end{array} \tag{3.3}$$

By [Theorem 3.8](#), $\hat{y} \circ \varphi = \hat{x}$ for some $x \in X$. Put $x = f(y)$. Thus f maps Y into X . Since X has the initial topology induced by η_X , f will be continuous if $\eta_X \circ f$ is continuous. Now $\eta_X \circ f(y) = \hat{x} = \hat{y} \circ \varphi = C\varphi(\eta_Y(y))$. Thus $\eta_X \circ f = C\varphi \circ \eta_Y$, so that $\eta_X \circ f$ is continuous, hence f is continuous, as required. It remains to prove that $\varphi = Cf$. Since the functions \hat{y} distinguish the points of $C_p Y$, it suffices to prove that $\hat{y} \circ \varphi = \hat{y} \circ Cf$ for every $y \in Y$. Now $\hat{y}(Cf(g)) = \hat{y}(g \circ f) = g \circ f(y) = g(f(y)) = g(x)$. Also $\hat{y}(\varphi(g)) = \hat{y} \circ \varphi(g) = \hat{x}(g) = g(x)$. Hence $\hat{y} \circ \varphi = \hat{y} \circ Cf$ for all $y \in Y$, so that $\varphi = Cf$.

Conversely suppose the morphism $\varphi : (C_p X, h_{C_p X}) \rightarrow (C_p Y, h_{C_p Y})$ is such that $\varphi = Cf$. Then by [Proposition 3.9](#), φ is an algebra morphism. \square

PROPOSITION 3.10. *The map $h_{C_p X} : MC_p X \rightarrow C_p X$ preserves the ring structure of the function spaces, operations being defined pointwise.*

PROOF. Let $\varphi, \psi \in MC_p X$, so that $\varphi, \psi : MX \rightarrow \mathbb{R}$. The maps $\varphi + \psi$, $\varphi \cdot \psi$, and $t\varphi$ (where $t \in \mathbb{R}$) are both defined pointwise, so that $(\varphi + \psi)(\lambda) = \varphi(\lambda) + \psi(\lambda)$, $\varphi \cdot \psi(\lambda) = \varphi(\lambda) \cdot \psi(\lambda)$, and $(t\varphi)(\lambda) = t\varphi(\lambda)$, for all $\lambda \in C_p X$. Now $h_{C_p X}(\varphi) = C\eta_X(\varphi) = \varphi \circ \eta_X$, hence $h_{C_p X}(\varphi + \psi) = (\varphi + \psi) \circ \eta_X$. Thus

$$\begin{aligned}
 (\varphi + \psi) \circ \eta_X(x) &= (\varphi + \psi)(\eta_X(x)) = \varphi(\eta_X(x)) + \psi(\eta_X(x)) \\
 &= h_{C_p X}(\varphi)(x) + h_{C_p X}(\psi)(x) = (h_{C_p X}(\varphi) + h_{C_p X}(\psi))(x).
 \end{aligned} \tag{3.4}$$

Since this holds for every $x \in X$, we have $h_{C_p X}(\varphi + \psi) = h_{C_p X}(\varphi) + h_{C_p X}(\psi)$.

The proof that $h_{C_p X}(\varphi \cdot \psi) = h_{C_p X}(\varphi) \cdot h_{C_p X}(\psi)$ is similar. We also have $h_{C_p X}(t\varphi) = th_{C_p X}(\varphi)$, where t is a scalar. \square

PROPOSITION 3.11. *For any $f : Y \rightarrow X$, the map $C_p f : C_p X \rightarrow C_p Y$ preserves the ring structure.*

PROOF. Since $C_p f$ acts by composition on the right, the result is clear. We will verify one case only: $C_p f(\varphi + \psi) = C_p f(\varphi) + C_p f(\psi)$. Then

$$\begin{aligned} C_p f(\varphi + \psi)(\gamma) &= (\varphi + \psi)(f(\gamma)) = \varphi(f(\gamma)) + \psi(f(\gamma)) \\ &= C_p f(\varphi)(\gamma) + C_p f(\psi)(\gamma) = (C_p f(\varphi) + C_p f(\psi))(\gamma). \end{aligned} \quad (3.5)$$

Since the equality holds for every $\gamma \in Y$, $C_p f(\varphi + \psi) = C_p f(\varphi) + C_p f(\psi)$. \square

PROBLEM 3.12. Characterize fully the Eilenberg-Moore category of M -algebras.

REFERENCES

- [1] J. Adámek, H. Herrlich, and G. E. Strecker, *Abstract and Concrete Categories*, Pure and Applied Mathematics, John Wiley & Sons, New York, 1990.
- [2] A. V. Arkhangel'skiĭ, *Topological Function Spaces*, Mathematics and Its Applications (Soviet Series), vol. 78, Kluwer Academic Publishers, Dordrecht, 1992.

KOENA R. NAILANA: DEPARTMENT OF MATHEMATICS, APPLIED MATHEMATICS AND ASTRONOMY, UNIVERSITY OF SOUTH AFRICA, P.O. BOX 392, PRETORIA 0003, SOUTH AFRICA
E-mail address: nailkr@unisa.ac.za



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

