COMMON FIXED POINT THEOREMS FOR A WEAK DISTANCE IN COMPLETE METRIC SPACES

JEONG SHEOK UME and SUCHEOL YI

Received 17 October 2001

Using the concept of a w -distance, we obtain common fixed point theorems on complete metric spaces. Our results generalize the corresponding theorems of Jungck, Fisher, Dien, and Liu.

2000 Mathematics Subject Classification: 47H10.

1. Introduction. In 1976, Caristi [1] proved a fixed point theorem in a complete metric space which generalizes the Banach contraction principle. This theorem is very useful and has many applications. Later, Dien [3] showed that a pair of mappings satisfying both the Banach contraction principle and Caristi's condition in a complete metric space has a common fixed point. That is to say, let (X, d) be a complete metric space and let S and T be two orbitally continuous mappings of X into itself. Suppose that there exists a finite number of functions $\{\varphi_i\}_{1 \leq i \leq N_0}$ of X into $\mathbb{R}_+$ such that

$$d(Sx, Ty) \leq q \cdot d(x, y) + \sum_{i=1}^{N_0} [\varphi_i(x) - \varphi_i(Sx) + \varphi_i(y) - \varphi_i(Ty)] \quad (1.1)$$

for all $x, y \in X$ and some $q \in [0, 1)$. Then S and T have a unique common fixed point z in X . Further, if $x \in X$ then $S^n x \rightarrow z$ and $T^n x \rightarrow z$ as $n \rightarrow \infty$. In particular, if S is an identity mapping, $q = 0$, and $N_0 = 1$, then this means a Caristi's fixed point theorem.

Recently, Liu [7] obtained necessary and sufficient conditions for the existence of fixed point of continuous self-mapping by using the ideas of Jungck [5] and Dien [3]: let f be a continuous self-mapping of a metric space (X, d) , then f has a fixed point in X if and only if there exist $z \in X$, a mapping $g : X \rightarrow X$, and a function Φ from X into $[0, \infty)$ such that f and g are compatible, $g(X) \subset f(X)$, g is continuous, and

$$d(gx, z) \leq r d(fx, z) + [\Phi(fx) - \Phi(gx)] \quad (1.2)$$

for all $x \in X$ and some $r \in [0, 1)$.

In 1996, Kada et al. [6] introduced the concept of w -distance on a metric space as follows: let X be a metric space with metric d , then a function $p : X \times X \rightarrow [0, \infty)$ is called a w -distance on X if the following are satisfied:

- (1) $p(x, z) \leq p(x, y) + p(y, z)$ for any $x, y, z \in X$;
- (2) for any $x \in X$, $p(x, \cdot) : X \rightarrow [0, \infty)$ is lower semicontinuous;

- (3) for any $\epsilon > 0$, there exists $\delta > 0$ such that $p(z, x) \leq \delta$ and $p(z, y) \leq \delta$ imply $d(x, y) \leq \epsilon$.

In this paper, using the concept of a w -distance, we obtain common fixed point theorems on complete metric spaces. Our results generalize the corresponding theorems of Jungck [5], Fisher [4], Dien [3], and Liu [7].

2. Definitions and preliminaries. Throughout, we denote by $\mathbb{N}$ the set of positive integers and by $\mathbb{R}_+$ the set of nonnegative real numbers, that is, $\mathbb{R}_+ := [0, \infty)$.

DEFINITION 2.1 (see [3]). A mapping T of a space X into itself is said to be orbitally continuous if $x_0 \in X$ such that $x_0 = \lim_{i \rightarrow \infty} T^{n_i}x$ for some $x \in X$, then $Tx_0 = \lim_{i \rightarrow \infty} T(T^{n_i}x)$.

DEFINITION 2.2 (see [2]). Let T be a mapping of a metric space X into itself. For each $x \in X$, let

$$\begin{aligned} O(T, x, n) &= \{x, Tx, \dots, T^n x\}, \quad n = 1, 2, \dots, \\ O(T, x, \infty) &= \{x, Tx, \dots\}. \end{aligned} \tag{2.1}$$

A space X is said to be T -orbitally complete if and only if every Cauchy sequence, which is contained in $O(T, x, \infty)$ for some $x \in X$, converges in X .

DEFINITION 2.3 (see [6]). Let X be a metric space with metric d . Then a function $p : X \times X \rightarrow \mathbb{R}_+$ is called a w -distance on X if the following properties are satisfied:

- (1) $p(x, z) \leq p(x, y) + p(y, z)$ for any $x, y, z \in X$;
- (2) for any $x \in X$, $p(x, \cdot) : X \rightarrow \mathbb{R}_+$ is lower semicontinuous;
- (3) for any $\epsilon > 0$, there exists $\delta > 0$ such that $p(z, x) \leq \delta$ and $p(z, y) \leq \delta$ imply $d(x, y) \leq \epsilon$.

The metric d is a w -distance on X . Other examples of w -distance are stated in [6].

DEFINITION 2.4 (see [5]). Let (X, d) be a metric space and $f, g : X \rightarrow X$. The mappings f and g are called compatible if and only if for every sequence $\{x_n\}_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow \infty} f x_n = \lim_{n \rightarrow \infty} g x_n = t$ for some $t \in X$, it implies

$$\lim_{n \rightarrow \infty} d(f g x_n, g f x_n) = 0. \tag{2.2}$$

LEMMA 2.5 (see [6]). Let X be a metric space with metric d , and p a w -distance on X . Let $\{x_n\}$ and $\{y_n\}$ be sequences in X , let $\{\alpha_n\}$ and $\{\beta_n\}$ be sequences in $\mathbb{R}_+$ converging to 0, and let $x, y, z \in X$. Then the following properties hold:

- (i) if $p(x_n, y) \leq \alpha_n$ and $p(x_n, z) \leq \beta_n$ for any $n \in \mathbb{N}$, then $y = z$. In particular, if $p(x, y) = 0$ and $p(x, z) = 0$, then $y = z$;
- (ii) if $p(x_n, y_n) \leq \alpha_n$ and $p(x_n, z) \leq \beta_n$ for any $n \in \mathbb{N}$, then $\{y_n\}$ converges to z ;
- (iii) if $p(x_n, x_m) \leq \alpha_n$ for any $n, m \in \mathbb{N}$ with $m > n$, then $\{x_n\}$ is a Cauchy sequence;
- (iv) if $p(y, x_n) \leq \alpha_n$ for any $n \in \mathbb{N}$, then $\{x_n\}$ is a Cauchy sequence.

3. Main results

THEOREM 3.1. Let (X, d) be a complete metric space with a w -distance p . Suppose that two mappings $f, g : X \rightarrow X$ and a function φ from X into $\mathbb{R}_+$ are satisfying the following conditions:

- (i) $g(X) \subseteq f(X)$,
- (ii) there exists $t \in X$ such that $p(t, gx) \leq r \cdot p(t, fx) + [\varphi(fx) - \varphi(gx)]$ for all $x \in X$ and some $r \in [0, 1)$,
- (iii) for every sequence $\{x_n\}_{n \in \mathbb{N}}$ in X satisfying

$$\lim_{n \rightarrow \infty} p(t, fx_n) = \lim_{n \rightarrow \infty} p(t, gx_n) = 0, \quad (3.1)$$

it implies that

$$\lim_{n \rightarrow \infty} \max \{p(t, fx_n), p(t, gx_n), p(fgx_n, gfx_n)\} = 0, \quad (3.2)$$

- (iv) for each $u \in X$ with $u \neq fu$ or $u \neq gu$,

$$\inf \{p(u, fx) + p(u, gx) + p(fgx, gfx) : x \in X\} > 0. \quad (3.3)$$

Then f and g have a unique common fixed point in X .

PROOF. Let x_0 be a given point of X . By (i), there exists $x_n \in X$ such that $gx_{n-1} = fx_n$ for $n \geq 1$. From [Theorem 3.1\(ii\)](#), we have

$$p(t, fx_{j+1}) = p(t, gx_j) \leq r \cdot p(t, fx_j) + [\varphi(fx_j) - \varphi(gx_j)], \quad (3.4)$$

which implies that

$$\sum_{j=0}^{n-1} p(t, fx_{j+1}) \leq r \cdot \sum_{j=0}^{n-1} p(t, fx_j) + \sum_{j=0}^{n-1} [\varphi(fx_j) - \varphi(gx_j)], \quad (3.5)$$

that is,

$$\begin{aligned} \sum_{j=1}^n p(t, fx_j) &\leq \frac{r}{1-r} p(t, fx_0) + \frac{1}{1-r} [\varphi(fx_0) - \varphi(fx_n)], \\ &\leq \frac{r}{1-r} p(t, fx_0) + \frac{1}{1-r} \varphi(fx_0), \end{aligned} \quad (3.6)$$

which means that the series $\sum_{n=1}^{\infty} p(t, fx_n)$ is convergent, so

$$\lim_{n \rightarrow \infty} p(t, fx_n) = \lim_{n \rightarrow \infty} p(t, gx_n) = 0. \quad (3.7)$$

Suppose that $t \neq ft$ or $t \neq gt$. Then, from [Theorem 3.1](#)(iii) and (iv) we obtain that

$$\begin{aligned} 0 &< \inf \{p(t, fx) + p(t, gx) + p(fgx, gfx) : x \in X\} \\ &\leq \inf \{p(t, fx_n) + p(t, gx_n) + p(fgx_n, gfx_n) : n \in \mathbb{N}\} \\ &= 0. \end{aligned} \quad (3.8)$$

This is a contradiction. Hence t is a common fixed point of f and g .

We prove that t is a unique common fixed point of f and g . Let u be a common fixed point of f and g . Then, by [Theorem 3.1](#)(ii),

$$\begin{aligned} p(t, t) &= p(t, gt) \leq r \cdot p(t, ft) + [\varphi(ft) - \varphi(gt)] = r \cdot p(t, t), \\ p(t, u) &= p(t, gu) \leq r \cdot p(t, fu) + [\varphi(fu) - \varphi(gu)] = r \cdot p(t, u). \end{aligned} \quad (3.9)$$

Thus $p(t, t) = p(t, u) = 0$. From [Lemma 2.5](#), we obtain $t = u$. Therefore t is a unique common fixed point of f and g . $\square$

REMARK 3.2. [Theorem 3.1](#) generalizes and improves Dien [3, Theorem 2.2] and Liu [7, Theorem 3.2].

THEOREM 3.3. *Let f be a continuous self-mapping of metric space (X, d) . Assume that f has a fixed point in X . Then there exists a w -distance $p, t \in X$, a continuous mapping $g : X \rightarrow X$, and a function φ from X into $\mathbb{R}_+$ satisfying [Theorem 3.1](#)(i), (ii), (iii), and (iv).*

PROOF. Let z be a fixed point of f , $r = 1/2$, $gx = t = z$, and $\varphi(x) = 1$ for all $x \in X$. Define $p : X \times X \rightarrow \mathbb{R}_+$ by

$$p(x, y) = \max \{d(fx, x), d(fx, y), d(fx, fy)\} \quad \forall x, y \in X. \quad (3.10)$$

Suppose that

$$\lim_{n \rightarrow \infty} p(t, fx_n) = \lim_{n \rightarrow \infty} p(t, gx_n) = 0. \quad (3.11)$$

Then it is easy to verify that the results of [Theorem 3.3](#) follow. $\square$

THEOREM 3.4. *Let f and g be a continuous compatible self-mappings of the metric space (X, d) . There exists $t \in X$ satisfying*

$$d(t, gx) \leq r \cdot d(t, fx) + [\varphi(fx) - \varphi(gx)] \quad (3.12)$$

for all $x \in X$ and some $r \in [0, 1)$. Then

(i) for every sequence $\{x_n\}_{n \in \mathbb{N}}$ in X such that

$$\lim_{n \rightarrow \infty} d(t, fx_n) = \lim_{n \rightarrow \infty} d(t, gx_n) = 0 \quad (3.13)$$

for some $t \in X$, it implies that

$$\lim_{n \rightarrow \infty} \max \{d(t, fx_n), d(t, gx_n), d(fgx_n, gfx_n)\} = 0; \quad (3.14)$$

(ii) for each $u \in X$ with $u \neq fu$ or $u \neq gu$,

$$\inf \{d(u, fx) + d(u, gx) + d(fgx, gfx) : x \in X\} > 0. \quad (3.15)$$

PROOF. The results follow by elementary calculation. $\square$

REMARK 3.5. Since the metric d is w -distance, from Theorems 3.1, 3.3, and 3.4, we obtain Liu [7, Theorem 3.1].

THEOREM 3.6. Let (X, d) be a complete metric space with a w -distance p , two mappings $f, g : X \rightarrow X$, and two functions φ, ψ from X into $\mathbb{R}_+$ such that Theorem 3.1(i), (iv) are satisfied,

(i) for every sequence $\{x_n\}_{n \in \mathbb{N}}$ in X such that

$$\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = t \quad (3.16)$$

for some $t \in X$, it implies that

$$\lim_{n \rightarrow \infty} \max \{p(t, fx_n), p(t, gx_n), p(fgx_n, gfx_n)\} = 0, \quad (3.17)$$

(ii)

$$\begin{aligned} p(gx, gy) &\leq a_1 p(fx, fy) + a_2 p(fx, gx) + a_3 p(fy, gy) \\ &\quad + a_4 p(fx, gy) + a_5 [p(gx, fy) d(fy, gx)]^{1/2} \\ &\quad + [\varphi(fx) - \varphi(gx)] + [\psi(fy) - \psi(gy)] \end{aligned} \quad (3.18)$$

for all $x, y \in X$, where a_1, a_2, a_3, a_4 , and a_5 are in $[0, 1)$ with $a_1 + a_4 + a_5 < 1$ and $a_1 + a_2 + a_3 + 2a_4 < 1$.

Then f and g have a unique common fixed point in X .

PROOF. Let x_0 be an arbitrary point of X . By Theorem 3.1(i), we obtain a sequence $\{x_n\}$ in X such that $gx_{n-1} = fx_n$ for $n \geq 1$. Let $y_n = p(fx_n, fx_{n+1})$ for $n \geq 0$. It follows from Theorem 3.6(ii) that

$$\begin{aligned} y_{j+1} &= p(gx_j, gx_{j+1}) \\ &\leq a_1 p(fx_j, fx_{j+1}) + a_2 p(fx_j, gx_j) + a_3 p(fx_{j+1}, gx_{j+1}) \\ &\quad + a_4 p(fx_j, gx_{j+1}) + a_5 [p(gx_j, fx_{j+1}) d(fx_{j+1}, gx_j)]^{1/2} \\ &\quad + [\varphi(fx_j) - \varphi(gx_j)] + [\psi(fx_{j+1}) - \psi(gx_{j+1})] \\ &\leq (a_1 + a_2 + a_4) y_j + (a_3 + a_4) y_{j+1} \\ &\quad + [\varphi(fx_j) - \varphi(fx_{j+1})] + [\psi(fx_{j+1}) - \psi(fx_{j+2})], \end{aligned} \quad (3.19)$$

which implies that

$$y_{j+1} \leq L_1 y_j + L_2 [\varphi(fx_j) - \varphi(fx_{j+1}) + \psi(fx_{j+1}) - \psi(fx_{j+2})], \quad (3.20)$$

where

$$L_1 = \frac{a_1 + a_2 + a_4}{1 - a_3 - a_4}, \quad L_2 = \frac{1}{1 - a_3 - a_4}. \quad (3.21)$$

Thus

$$\sum_{j=1}^n y_j \leq \frac{L_1}{1-L_1} y_0 + \frac{L_2}{1-L_1} [\varphi(fx_0) + \psi(fx_1)] \quad (3.22)$$

for all $n \geq 1$. Hence, the series $\sum_{n=1}^{\infty} y_n$ is convergent. For any $n, r \geq 1$, we have

$$p(fx_n, fx_{n+r}) \leq \sum_{i=n}^{n+r-1} y_i. \quad (3.23)$$

By Lemma 2.5, this implies that $\{fx_n\}_{n=1}^{\infty}$ is a Cauchy sequence in X . Since X is a complete metric space, there exists $t \in X$ such that $fx_n \rightarrow t$ as $n \rightarrow \infty$. From Theorem 3.6(i), we have

$$\lim_{n \rightarrow \infty} \max \{p(t, fx_n), p(t, gx_n), p(fgx_n, gfx_n)\} = 0. \quad (3.24)$$

Suppose that $t \neq ft$ or $t \neq gt$, then from Theorem 3.1(iv) we obtain that

$$\begin{aligned} 0 &< \inf \{p(t, fx) + p(t, gx) + p(fgx, gfx) : x \in X\} \\ &\leq \inf \{p(t, fx_n) + p(t, gx_n) + p(fgx_n, gfx_n) : n \in \mathbb{N}\} \\ &= 0. \end{aligned} \quad (3.25)$$

which is a contradiction. Therefore t is a common fixed point of f and g . It follows from Lemma 2.5 and Theorem 3.6(ii) that t is a unique common fixed point of f and g . $\square$

THEOREM 3.7. *Let f be a continuous self-mapping of a metric space (X, d) . Assume that f has a fixed point in X . Then there exist a w -distance $p, t \in X$, a continuous mapping $g : X \rightarrow X$, and functions φ, ψ from X into $\mathbb{R}_+$ satisfying Theorem 3.1(i), (iv) and Theorem 3.6(i), (ii).*

PROOF. By a method similar to that in the proof of Theorem 3.3, the results follow. $\square$

REMARK 3.8. Since the metric d is w -distance, from Theorems 3.4, 3.6, and 3.7, we obtain Jungck [5, Theorem], Fisher [4, Theorem 2], and Liu [7, Theorem 3.3].

ACKNOWLEDGMENT. This work was supported by Korea Research Foundation Grant (KRF-2001-015-DP0025).

REFERENCES

- [1] J. Caristi, *Fixed point theorems for mappings satisfying inwardness conditions*, Trans. Amer. Math. Soc. **215** (1976), 241–251.
- [2] L. B. Ćirić, *A generalization of Banach's contraction principle*, Proc. Amer. Math. Soc. **45** (1974), 267–273.
- [3] N. H. Dien, *Some remarks on common fixed point theorems*, J. Math. Anal. Appl. **187** (1994), no. 1, 76–90.
- [4] B. Fisher, *Mappings with a common fixed point*, Math. Sem. Notes Kobe Univ. **7** (1979), no. 1, 81–84.

- [5] G. Jungck, *Commuting mappings and fixed points*, Amer. Math. Monthly **83** (1976), no. 4, 261-263.
- [6] O. Kada, T. Suzuki, and W. Takahashi, *Nonconvex minimization theorems and fixed point theorems in complete metric spaces*, Math. Japon. **44** (1996), no. 2, 381-391.
- [7] Z. Liu, Y. Xu, and Y. J. Cho, *On characterizations of fixed and common fixed points*, J. Math. Anal. Appl. **222** (1998), no. 2, 494-504.

JEONG SHEOK UME: DEPARTMENT OF APPLIED MATHEMATICS, CHANGWON NATIONAL UNIVERSITY, CHANGWON 641-773, KOREA

E-mail address: jsume@sarim.changwon.ac.kr

SUCHEOL YI: DEPARTMENT OF APPLIED MATHEMATICS, CHANGWON NATIONAL UNIVERSITY, CHANGWON 641-773, KOREA

E-mail address: scyi@sarim.changwon.ac.kr