ON HOPF DEMEYER-KANZAKI GALOIS EXTENSIONS

GEORGE SZETO and LIANYONG XUE

Received 15 October 2002

Let H be a finite-dimensional Hopf algebra over a field k , B a left H -module algebra, and H^* the dual Hopf algebra of H . For an H^* -Azumaya Galois extension B with center C , it is shown that B is an H^* -DeMeyer-Kanzaki Galois extension if and only if C is a maximal commutative separable subalgebra of the smash product $B\#H$. Moreover, the characterization of a commutative Galois algebra as given by S. Ikehata (1981) is generalized.

2000 Mathematics Subject Classification: 16W30, 16H05.

1. Introduction. Let H be a finite-dimensional Hopf algebra over a field k , B a left H -module algebra, and H^* the dual Hopf algebra of H . In [7], the class of Azumaya Galois extensions of a ring as studied in [1, 2] was generalized to H^* -Azumaya Galois extensions. An H^* -Azumaya Galois extension B was characterized in terms of the smash product $B\#H$ see [7, Theorem 3.4]. Observing that the commutator $V_B(B^H)$ of B^H in B is also an H^* -Azumaya Galois extension (see [7, Lemma 4.1]), in the present paper, we will give a characterization of an H^* -Azumaya Galois extension B in terms of $V_B(B^H)$. Moreover, we will investigate the class of H^* -Azumaya Galois extensions B such that $V_B(B^H) = C$, where C is the center of B . We note that when $H = kG$, where G is a finite automorphism group of B , such a B is precisely a DeMeyer-Kanzaki Galois extension with Galois group G [3, 6, 8, 9]. Several equivalent conditions are then given for an H^* -Azumaya Galois extension being an H^* -DeMeyer-Kanzaki Galois extension, and the characterization of a commutative Galois algebra as given by Ikehata [5, Theorem 2] is generalized to an H^* -DeMeyer-Kanzaki Galois extension.

2. Basic definitions and notation. Throughout, H denotes a finite-dimensional Hopf algebra over a field k with comultiplication Δ and counit ε , H^* the dual Hopf algebra of H , B a left H -module algebra, C the center of B , $B^H = \{b \in B \mid hb = \varepsilon(h)b \text{ for all } h \in H\}$, and $B\#H$ the smash product of B with H , where $B\#H = B \otimes_k H$ such that, for all $b\#h$ and $b'\#h'$ in $B\#H$, $(b\#h)(b'\#h') = \sum b(h_1b')\#h_2h'$, where $\Delta(h) = \sum h_1 \otimes h_2$.

For a subring A of B with the same identity 1, we denote the commutator subring of A in B by $V_B(A)$. We call B a separable extension of A if there

exist $\{a_i, b_i \text{ in } B, i = 1, 2, \dots, m \text{ for some integer } m\}$ such that $\sum a_i b_i = 1$ and $\sum b a_i \otimes b_i = \sum a_i \otimes b_i b$ for all b in B where $\otimes$ is over A . An Azumaya algebra is a separable extension of its center. A ring B is called a Hirata separable extension of A if $B \otimes_A B$ is isomorphic to a direct summand of a finite direct sum of B as a B -bimodule. A ring B is called an H^* -Galois extension of B^H if B is a right H^* -comodule algebra with structure map $\rho : B \rightarrow B \otimes_k H^*$ such that $\beta : B \otimes_{B^H} B \rightarrow B \otimes_k H^*$ is a bijection where $\beta(a \otimes b) = (a \otimes 1)\rho(b)$. An H^* -Galois extension B is called an H^* -Azumaya Galois extension if B is separable over B^G which is an Azumaya algebra over C^G , and an H^* -DeMeyer-Kanzaki Galois extension if B is an H^* -Azumaya Galois extension and $V_B(B^H) = C$.

Let P be a finitely generated and projective module over a commutative ring R . Then for a prime ideal p of R , $P_p (= P \otimes_R R_p)$ is a free module over $R_p (= \text{the local ring of } R \text{ at } p)$, and the rank of P_p over R_p is the number of copies of R_p in P_p , that is, $\text{rank}_{R_p}(P_p) = m$ for some integer m . It is known that the $\text{rank}_R(P)$ is a continuous function ($\text{rank}_R(P)(p) = \text{rank}_{R_p}(P_p) = m$) from $\text{Spec}(R)$ to the set of nonnegative integers with the discrete topology (see [4, Corollary 4.11, page 31]). We will use the $\text{rank}_R(P)$ -function for a finitely generated and projective module P over a commutative ring R .

3. H^* -Azumaya Galois extensions. In this section, keeping all notations as given in Section 2, we will characterize an H^* -Azumaya Galois extension B in terms of the commutator $V_B(B^H)$ of B^H in B .

THEOREM 3.1. *If $B = B^H \cdot V_B(B^H)$, then $(V_B(B^H))^H = C^H$.*

PROOF. Since $C \subset V_B(B^H)$, $C^H \subset (V_B(B^H))^H$. Conversely, since $V_B(B^H) \subset B$, $(V_B(B^H))^H \subset B^H$. Hence $(V_B(B^H))^H \subset B^H \cap V_B(B^H) \subset \text{the center of } V_B(B^H)$. But $B = B^H \cdot V_B(B^H)$, so the center of $V_B(B^H)$ is C . Thus, $(V_B(B^H))^H \subset C^H$. $\square$

THEOREM 3.2. *A ring B is an H^* -Azumaya Galois extension of B^H if and only if $B = B^H \cdot V_B(B^H)$ such that $V_B(B^H)$ is an H^* -Azumaya Galois extension of C^H and B^H is an Azumaya C^H -algebra.*

PROOF. ($\Rightarrow$) Since B is an H^* -Azumaya Galois extension of B^H , then $V_B(B^H)$ is an H^* -Azumaya Galois extension of $(V_B(B^H))^H$ (see [7, Lemma 4.1]) and B^H is an Azumaya C^H -algebra (see [7, Theorem 3.4]). Moreover, by the proof of [7, Lemma 4.1], $B \# H$ is an Azumaya C^H -algebra such that $B \# H \cong B^H \otimes_{C^H} (V_B(B^H) \# H) \cong B^H (V_B(B^H) \# H)$, where B^H and $V_B(B^H) \# H$ are Azumaya C^H -algebras. But H is a finite-dimensional Hopf algebra over a field k , so $B \cong B^H \otimes_{C^H} V_B(B^H)$ from the isomorphism $B \# H \cong B^H \otimes_{C^H} (V_B(B^H) \# H)$, and so $B = B^H \cdot V_B(B^H)$. Hence $(V_B(B^H))^H = C^H$ by Theorem 3.1. Thus $V_B(B^H)$ is an H^* -Azumaya Galois C^H -algebra.

($\Leftarrow$) Since $V_B(B^H)$ is an H^* -Azumaya Galois algebra over C^H , $V_B(B^H) \# H$ is an Azumaya C^H -algebra [7, Theorem 3.4]. By hypothesis, B^H is an Azumaya C^H -algebra, so $B^H \otimes_{C^H} (V_B(B^H) \# H) \cong B^H V_B(B^H) \# H = B \# H$ which is an Azumaya

C^H -algebra. Thus $B\#H$ is a Hirata separable extension of B (see [5, Theorem 1]). Moreover, $V_B(B^H)$ is a separable C^H -algebra (see [7, Theorem 3.4]) and B^H is an Azumaya C^H -algebra by hypothesis, so $B^H \cdot V_B(B^H) (= B)$ is also a separable C^H -algebra. Thus B is an H^* -Azumaya Galois extension of B^H [7, Theorem 3.4]. $\square$

Next we generalize the characterization of a commutative Galois algebra as given by Ikehata (see [5, Theorem 2]) to a commutative H^* -Galois algebra.

LEMMA 3.3. *If C is a commutative H^* -Galois algebra over C^H , then C is a maximal commutative subalgebra of $C\#H$.*

PROOF. Since C is a commutative H^* -Galois algebra over C^H , $C\#H \cong \text{Hom}_{C^H}(C, C)$ [6, Theorem 1.7]. Hence it suffices to show that $V_{\text{Hom}_{C^H}(C, C)}(C_L) = C_L$ where $C_L = \{c_L, \text{ the left multiplication map induced by } c \in C\}$. In fact, $C_L \subset V_{\text{Hom}_{C^H}(C, C)}(C_L)$ is clear. Conversely, let $f \in V_{\text{Hom}_{C^H}(C, C)}(C_L)$. Then, for each $c \in C$, $(cf)(x) = (fc)(x)$ for all $x \in C$. Hence $cf(x) = f(cx)$, and so $cf(1) = f(c)$ for all $c \in C$. Thus $f(c) = d_f(c)$ for all $c \in C$, where $d_f = f(1) \in C$, that is, $f = (d_f)_L \in C_L$. $\square$

THEOREM 3.4. *Let C be a commutative separable C^H -algebra containing C^H as a direct summand as a C^H -module. Then, C is a commutative H^* -Galois algebra over C^H if and only if $C \otimes_{C^H} (C\#H) \cong M_n(C)$, the matrix algebra over C of order n where n is the dimension of H over k .*

PROOF. $(\Rightarrow)$ Since C is an H^* -Galois algebra over C^H , $C\#H \cong \text{Hom}_{C^H}(C, C)$ such that C is finitely generated and projective over C^H [6, Theorem 1.7]. Hence $C\#H$ is an Azumaya C^H -algebra and C is a maximal commutative subalgebra of the Azumaya C^H -algebra $C\#H$ by Lemma 3.3. By hypothesis, C is also a separable C^H -algebra, so C is a splitting ring for the Azumaya C^H -algebra $C\#H$ such that $C \otimes_{C^H} (C\#H) \cong \text{Hom}_C(C\#H, C\#H)$ (see the proof of [4, Theorem 5.5, page 64]). Noting that $C\#H = C \otimes_k H$ which is a free C -module of rank n where $n = \dim_k(H)$, we have that $C \otimes_{C^H} (C\#H) \cong M_n(C)$.

$(\Leftarrow)$ Since $C \otimes_{C^H} (C\#H) \cong M_n(C)$, $C \otimes_{C^H} (C\#H)$ is an Azumaya C -algebra. By hypothesis, C^H is a direct summand of C as a C^H -module, so $C\#H$ is an Azumaya C^H -algebra [4, Corollary 1.10, page 45]. Hence $C\#H$ is a Hirata separable extension of C . But C is a separable C^H -algebra by hypothesis, so C is an H^* -Galois algebra over C^H [7, Theorem 3.4].

We remark that the necessity does not need the hypothesis that C^H is a direct summand of C . $\square$

4. H^* -DeMeyer-Kanzaki Galois extensions. We recall that B is an H^* -DeMeyer-Kanzaki Galois extension of B^H if B is an H^* -Azumaya Galois extension of B^H and $V_B(B^H) = C$. In this section, we characterize an H^* -DeMeyer-Kanzaki Galois extension in terms of the smash product $V_B(B^H)\#H$ and prove that C is a splitting ring for the Azumaya C^H -algebras $V_B(B^H)\#H$ and $B\#H$.

THEOREM 4.1. *Let B be an H^* -Azumaya Galois extension of B^H . Then the following statements are equivalent:*

- (1) B is an H^* -DeMeyer-Kanzaki Galois extension of B^H ;
- (2) $\text{rank}_{C^H}(V_B(B^H)) = \text{rank}_{C^H}(C)$;
- (3) C is a maximal commutative separable subalgebra of $V_B(B^H) \# H$.

PROOF. (1) $\Rightarrow$ (2). It is clear.

(2) $\Rightarrow$ (1). Since B is an H^* -Azumaya Galois extension of B^H , $V_B(B^H)$ is an H^* -Azumaya Galois algebra over C^H by Theorem 3.2 such that $V_B(B^H)$ is a separable and finitely generated projective module over C^H (see [7, Theorem 3.4]). Hence the rank function $\text{rank}_{C^H}(V_B(B^H))$ is defined and $V_B(B^H)$ is an Azumaya algebra over its center [4, Theorem 3.8, page 55]. But $B = B^H \cdot V_B(B^H)$ by Theorem 3.2, so the center of $V_B(B^H)$ is C . Thus $V_B(B^H)$ is an Azumaya C -algebra; and so C is a direct summand $V_B(B^H)$ as a C -module. This implies that C is a direct summand $V_B(B^H)$ as a C^H -module. Therefore the rank function $\text{rank}_{C^H}(C)$ is also defined. Now by hypothesis, $\text{rank}_{C^H}(V_B(B^H)) = \text{rank}_{C^H}(C)$, so $V_B(B^H) = C$, that is, B is an H^* -DeMeyer-Kanzaki Galois extension of B^H .

(1) $\Rightarrow$ (3). Since B is an H^* -DeMeyer-Kanzaki Galois extension of B^H , B is an H^* -Azumaya Galois extension such that $V_B(B^H) = C$. Hence $B = B^H \cdot V_B(B^H) \cong B^H \otimes_{C^H} C$ such that C is an H^* -Galois algebra over C^H by Theorem 3.2, and so C is a separable C^H -algebra containing C^H as a direct summand as a C^H -module [7, Theorem 3.4]. Hence C is a maximal commutative separable subalgebra of $C \# H$ where $C = V_B(B^H)$ by Lemma 3.3.

(3) $\Rightarrow$ (2). Since B is an H^* -Azumaya Galois extension of B^H , $B = B^H \cdot V_B(B^H) \cong B^H \otimes_{C^H} V_B(B^H)$ such that $V_B(B^H)$ is an H^* -Azumaya Galois algebra over C^H by Theorem 3.2. Hence $V_B(B^H) \# H$ is an Azumaya C^H -algebra and $V_B(B^H)$ is an Azumaya C -algebra [7, Theorem 3.4]. By hypothesis, C is a maximal commutative separable subalgebra of $V_B(B^H) \# H$, so

$$C \otimes_{C^H} (V_B(B^H) \# H) \cong \text{Hom}_C(V_B(B^H) \# H, V_B(B^H) \# H) \quad (4.1)$$

(see [4, Theorem 5.5, page 64]). On the other hand, $V_B(B^H) \# H \cong \text{Hom}_{C^H}(V_B(B^H), V_B(B^H))$ (see [7, Theorem 3.4]). Thus

$$\begin{aligned} C \otimes_{C^H} (V_B(B^H) \# H) &\cong C \otimes_{C^H} \text{Hom}_{C^H}(V_B(B^H), V_B(B^H)) \\ &\cong \text{Hom}_C(C \otimes_{C^H} V_B(B^H), C \otimes_{C^H} V_B(B^H)); \end{aligned} \quad (4.2)$$

and so $\text{Hom}_C(V_B(B^H) \# H, V_B(B^H) \# H) \cong \text{Hom}_C(C \otimes_{C^H} V_B(B^H), C \otimes_{C^H} V_B(B^H))$. This implies that $V_B(B^H) \# H \cong P \otimes_C (C \otimes_{C^H} V_B(B^H))$ for some finitely generated projective C -module P of rank 1, that is, $V_B(B^H) \# H \cong P \otimes_{C^H} V_B(B^H)$. Taking $\text{rank}_{C^H}(\cdot)$ both sides, we have that $n \cdot \text{rank}_{C^H}(V_B(B^H)) = (\text{rank}_{C^H}(P)) \cdot (\text{rank}_{C^H}(V_B(B^H)))$ where $n = \dim_k(H)$. But $\text{rank}_{C^H}(V_B(B^H))$ is also n , so $\text{rank}_{C^H}(C) = \text{rank}_{C^H}(P) = n = \text{rank}_{C^H}(V_B(B^H))$. $\square$

Theorem 4.1 implies that the Azumaya C^H -algebras $V_B(B^H)\#H$ and $B\#H$ have a nice splitting ring C which is an H^* -Galois algebra over C^H and separable over C^H such that $C \otimes_{C^H} (V_B(B^H)\#H)$ and $C \otimes_{C^H} (B\#H)$ are matrix algebras.

COROLLARY 4.2. *If B is an H^* -DeMeyer-Kanzaki Galois extension of B^H , then $C \otimes_{C^H} (V_B(B^H)\#H) \cong M_n(C)$, the matrix algebra over C of order n where $n = \dim_k(H)$.*

PROOF. By hypothesis, B is an H^* -DeMeyer-Kanzaki Galois extension of B^H , so $C (= V_B(B^H))$ is an H^* -Galois algebra over C^H by **Theorem 3.2**. Hence C is a separable C^H -algebra and $C\#H$ is an Azumaya C^H -algebra [7, Theorem 3.4]. Thus C^H is a direct summand of C as a C^H -module. Therefore, $C \otimes_{C^H} (C\#H) \cong M_n(C)$ by **Theorem 3.4**. $\square$

COROLLARY 4.3. *If B is an H^* -DeMeyer-Kanzaki Galois extension of B^H , then $C \otimes_{C^H} (B\#H) \cong M_n(B)$, the matrix algebra over B of order n where $n = \dim_k(H)$.*

PROOF. By **Corollary 4.2**, $C \otimes_{C^H} (C\#H) \cong M_n(C)$, so

$$B^H \otimes_{C^H} C \otimes_{C^H} (C\#H) \cong B^H \otimes_{C^H} M_n(C). \quad (4.3)$$

Since $B = B^H \cdot V_B(B^H) \cong B^H \otimes_{C^H} V_B(B^H) = B^H \otimes_{C^H} C$, we have that

$$\begin{aligned} C \otimes_{C^H} (B\#H) &\cong C \otimes_{C^H} ((B^H \otimes_{C^H} C)\#H) \\ &\cong C \otimes_{C^H} B^H \otimes_{C^H} (C\#H) \\ &\cong B^H \otimes_{C^H} C \otimes_{C^H} (C\#H) \\ &\cong B^H \otimes_{C^H} M_n(C) \cong M_n(B^H \otimes_{C^H} C) \\ &\cong M_n(B). \end{aligned} \quad (4.4) \quad \square$$

ACKNOWLEDGMENTS. This paper was written under the support of a Caterpillar Fellowship at Bradley University. The authors would like to thank Caterpillar Inc. for the support.

REFERENCES

- [1] R. Alfaro and G. Szeto, *Skew group rings which are Azumaya*, Comm. Algebra **23** (1995), no. 6, 2255–2261.
- [2] ———, *On Galois extensions of an Azumaya algebra*, Comm. Algebra **25** (1997), no. 6, 1873–1882.
- [3] F. R. DeMeyer, *Some notes on the general Galois theory of rings*, Osaka J. Math. **2** (1965), 117–127.
- [4] F. R. DeMeyer and E. Ingraham, *Separable Algebras over Commutative Rings*, Lecture Notes in Mathematics, vol. 181, Springer-Verlag, Berlin, 1971.
- [5] S. Ikehata, *Note on Azumaya algebras and H -separable extensions*, Math. J. Okayama Univ. **23** (1981), no. 1, 17–18.
- [6] H. F. Kreimer and M. Takeuchi, *Hopf algebras and Galois extensions of an algebra*, Indiana Univ. Math. J. **30** (1981), no. 5, 675–692.
- [7] M. Ouyang, *Azumaya extensions and Galois correspondence*, Algebra Colloq. **7** (2000), no. 1, 43–57.

- [8] G. Szeto and L. Xue, *Some correspondences for a center Galois extension*, Math. Japon. **52** (2000), no. 3, 463–468.
- [9] L. Xue, *The invariant subrings of DeMeyer-Kanzaki Galois extensions*, Port. Math. (N.S.) **59** (2002), no. 1, 37–45.

George Szeto: Department of Mathematics, Bradley University, Peoria, IL 61625, USA
E-mail address: szeto@hilltop.bradley.edu

Lianyong Xue: Department of Mathematics, Bradley University, Peoria, IL 61625, USA
E-mail address: lxue@hilltop.bradley.edu