A NEW CHARACTERIZATION OF SOME ALTERNATING AND SYMMETRIC GROUPS

AMIR KHOSRAVI and BEHROOZ KHOSRAVI

Received 5 February 2002

We suppose that $p = 2^\alpha 3^\beta + 1$, where $\alpha \geq 1$, $\beta \geq 0$, and $p \geq 7$ is a prime number. Then we prove that the simple groups A_n , where $n = p, p+1$, or $p+2$, and finite groups S_n , where $n = p, p+1$, are also uniquely determined by their order components. As corollaries of these results, the validity of a conjecture of J. G. Thompson and a conjecture of Shi and Bi (1990) both on A_n , where $n = p, p+1$, or $p+2$, is obtained. Also we generalize these conjectures for the groups S_n , where $n = p, p+1$.

2000 Mathematics Subject Classification: 20D05, 20D60, 20D08.

1. Introduction. Let G be a finite group. We denote by $\pi(G)$ the set of all prime divisors of $|G|$. We construct the prime graph of G as follows. *The prime graph* $\Gamma(G)$ of a group G is the graph whose vertex set is $\pi(G)$, and two distinct primes p and q are joined by an edge (we write $p \sim q$) if and only if G contains an element of order pq . Let $t(G)$ be the number of connected components of $\Gamma(G)$ and let $\pi_1, \pi_2, \dots, \pi_{t(G)}$ be the connected components of $\Gamma(G)$. If $2 \in \pi(G)$, then we always suppose that $2 \in \pi_1$.

Now $|G|$ can be expressed as a product of coprime positive integers m_i , $i = 1, 2, \dots, t(G)$, where $\pi(m_i) = \pi_i$. These integers are called *the order components* of G . The set of order components of G will be denoted by $OC(G)$. Also we call $m_2, \dots, m_{t(G)}$ *the odd-order components* of G . The order components of non-abelian simple groups having at least three prime graph components are obtained by Chen [7, Tables 1, 2, 3]. Similarly, the order components of non-abelian simple groups with two-order components can be obtained by using the tables in [18, 28].

The following groups are uniquely determined by their order components: Suzuki-Ree groups [6], Sporadic simple groups [4], $PSL_2(q)$ [7], $E_8(q)$ [2], $G_2(q)$, where $q \equiv 0 \pmod{3}$ [3], $F_4(q)$, where q is even [15], $PSL_3(q)$, where q is an odd prime power [14], $PSL_3(q)$, where $q = 2^n$ [13], $PSU_3(q)$, where $q > 5$ [16], and A_p , where p and $p-2$ are primes [12].

It was proved by Oyama [20] that a finite group which has the same table of characters as an alternating group A_n is isomorphic to A_n . It was also proved by Koike [17] that a finite group which has the isomorphic subgroup-lattice as an alternating group A_n is isomorphic to A_n .

Let $\pi_e(G)$ denote the set of orders of elements in G . Shi and Bi [27] proved that if $\pi_e(G) = \pi_e(A_n)$ and $|G| = |A_n|$, then $G \cong A_n$. Iranmanesh and Alavi [12] proved that if p and $p-2$ are primes and $\text{OC}(G) = \text{OC}(A_p)$, then $G \cong A_p$. Praeger and Shi [21] and Shi and Bi [26] proved that A_8 , A_9 , A_{11} , A_{13} , S_7 , and S_8 are characterizable by their element orders. Also recently, Kondrat'ev and Mazurov [19] and Zavarnitsin [29] proved that if $\pi_e(G) = \pi_e(A_n)$, where $n = s, s+1, s+2$ and s is a prime number, then $G \cong A_n$.

Now we prove the following theorems.

THEOREM 1.1. *Let $p = 2^\alpha 3^\beta + 1$, where $\alpha \geq 1$, $\beta \geq 0$, and $p \geq 7$ is a prime number. Let $M = A_n$, where $n = p, p+1, p+2$. Then $\text{OC}(G) = \text{OC}(M)$ if and only if $G \cong M$.*

THEOREM 1.2. *Let $p = 2^\alpha 3^\beta + 1$, where $\alpha \geq 1$, $\beta \geq 0$, and $p \geq 7$ is a prime number. Let $M = S_n$, where $n = p, p+1$. Then $\text{OC}(G) = \text{OC}(M)$ if and only if $G \cong M$.*

In this paper, all groups are finite and by simple groups we mean non-abelian simple groups. All further unexplained notations are standard and we refer, for example, to [10]. Also frequently we use the results of Williams [28] and Kondrat'ev [18] about the prime graph of simple groups.

2. Preliminary results

REMARK 2.1. Let N be a normal subgroup of G and $p \sim q$ in $\Gamma(G/N)$. Then $p \sim q$ in $\Gamma(G)$. In fact if $xN \in G/N$ has order pq , then there is a power of x which has order pq .

DEFINITION 2.2 (see [11]). A finite group G is called a 2-Frobenius group if it has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$, where K and G/H are Frobenius groups with kernels H and K/H , respectively.

LEMMA 2.3 (see [28, Theorem A]). *If G is a finite group with its prime graph having more than one component, then G is one of the following groups:*

- (a) *a Frobenius or 2-Frobenius group;*
- (b) *a simple group;*
- (c) *an extension of a π_1 -group by a simple group;*
- (d) *an extension of a simple group by a π_1 -solvable group;*
- (e) *an extension of a π_1 -group by a simple group by a π_1 -group.*

LEMMA 2.4 (see [28, Lemma 3]). *If G is a finite group with more than one prime graph component and has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that H and G/K are π_1 -groups and K/H is a simple group, then H is a nilpotent group.*

The next lemma follows from [1, Theorem 2].

LEMMA 2.5. *Let G be a Frobenius group of even order and let H, K be Frobenius complement and Frobenius kernel of G , respectively. Then $t(\Gamma(G)) = 2$,*

and the prime graph components of G are $\pi(H)$, $\pi(K)$ and G has one of the following structures:

- (a) $2 \in \pi(K)$ and all Sylow subgroups of H are cyclic;
- (b) $2 \in \pi(H)$, K is an abelian group, H is a solvable group, the Sylow subgroups of odd order of H are cyclic groups, and the 2-Sylow subgroups of H are cyclic or generalized quaternion groups;
- (c) $2 \in \pi(H)$, K is an abelian group, and there exists $H_0 \leq H$ such that $|H : H_0| \leq 2$, $H_0 = Z \times \text{SL}(2, 5)$, $(|Z|, 2.3.5) = 1$, and the Sylow subgroups of Z are cyclic.

The next lemma follows from [1, Theorem 2] and Lemma 2.4.

LEMMA 2.6. *Let G be a 2-Frobenius group of even order. Then $t(\Gamma(G)) = 2$ and G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that*

- (a) $\pi_1 = \pi(G/K) \cup \pi(H)$ and $\pi(K/H) = \pi_2$;
- (b) G/K and K/H are cyclic, $|G/K|$ divides $|\text{Aut}(K/H)|$, $(|G/K|, |K/H|) = 1$, and $|G/K| < |K/H|$;
- (c) H is nilpotent and G is a solvable group.

LEMMA 2.7 (see [8, Lemma 8]). *Let G be a finite group with $t(\Gamma(G)) \geq 2$ and let N be a normal subgroup of G . If N is a π_i -group for some prime graph component of G and $m_1, m_2, \dots, m_r$ are some order components of G but not a π_i -number, then $m_1 m_2 \cdots m_r$ is a divisor of $|N| - 1$.*

The next lemma follows from [5, Lemma 1.4].

LEMMA 2.8. *Suppose that G and M are two finite groups satisfying $t(\Gamma(M)) \geq 2$, $N(G) = N(M)$, where $N(G) = \{n \mid G \text{ has a conjugacy class of size } n\}$, and $Z(G) = 1$. Then $|G| = |M|$.*

LEMMA 2.9 (see [5, Lemma 1.5]). *Let G_1 and G_2 be finite groups satisfying $|G_1| = |G_2|$ and $N(G_1) = N(G_2)$. Then $t(\Gamma(G_1)) = t(\Gamma(G_2))$ and $\text{OC}(G_1) = \text{OC}(G_2)$.*

LEMMA 2.10. *Let G be a finite group and let M be a non-abelian finite group with $t(M) = 2$ satisfying $\text{OC}(G) = \text{OC}(M)$.*

- (1) *Let $|M| = m_1 m_2$, $\text{OC}(M) = \{m_1, m_2\}$, and $\pi(m_i) = \pi_i$ for $i = 1, 2$. Then $|G| = m_1 m_2$ and one of the following holds:*
 - (a) *G is a Frobenius or 2-Frobenius group;*
 - (b) *G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that G/K is a π_1 -group, H is a nilpotent π_1 -group, and K/H is a non-abelian simple group. Moreover, $\text{OC}(K/H) = \{m'_1, m'_2, \dots, m'_s, m_2\}$, $|K/H| = m'_1 m'_2 \cdots m'_s m_2$, and $m'_1 m'_2 \cdots m'_s \mid m_1$, where $\pi(m'_j) = \pi'_j$, $1 \leq j \leq s$.*
- (2) *In case (b), $|G/K| \mid |\text{Out}(K/H)|$.*

PROOF. The proof of (1) follows from the above lemmas. Since $t(G) \geq 2$, we have $t(G/H) \geq 2$. Otherwise $t(G/H) = 1$, so that $t(G) = 1$. Since H is a π_i -group,

TABLE 2.1

p	Finite simple C_{pp} groups
2	A_5, A_6 ; $L_2(q)$, where q is a Fermat prime, a Mersenne prime, or $q = 2^n$, $n \geq 3$, $L_3(2^2)$, $Sz(2^{2n+1})$, $n \geq 1$
3	A_5, A_6 ; $L_2(q)$, where $q = 2^3$, 3^{n+1} , or $2 \cdot 3^n \pm 1$ which is a prime, $n \geq 1$, $L_3(2^2)$
5	A_5, A_6, A_7 ; M_{11}, M_{22} ; $L_2(q)$, where $q = 7^2$, 5^n , or $2 \cdot 5^n \pm 1$ which is a prime, $n \geq 1$, $L_3(2^2)$, $S_4(q)$, $q = 3, 7$, $U_4(3)$; $Sz(q)$, $q = 2^3, 2^5$
7	A_7, A_8, A_9 ; M_{22}, J_1, J_2, HS ; $L_2(q)$, $q = 2^3, 7^n$, or $2 \cdot 7^n - 1$ which is a prime, $n \geq 1$, $L_3(2^2)$, $S_6(2)$, $O_8^+(2)$, $G_2(q)$, $q = 3, 19$; $U_3(q)$, $q = 3, 5, 19$; $U_4(3)$, $U_6(2)$, $Sz(2^3)$
13	A_{13}, A_{14}, A_{15} ; Suz, Fi_{22} ; $L_2(q)$, $q = 3^3, 5^2, 13^n$, or $2 \cdot 13^n - 1$ which is a prime, $n \geq 1$, $L_3(3)$, $L_4(3)$, $O_7(3)$, $S_4(5)$, $S_6(3)$, $O_8^+(3)$, $G_2(q)$, $q = 2^3, 3$; $F_4(2)$, $U_3(q)$, $q = 2^2, 23$, $Sz(2^3)$, ${}^3D_4(2)$, ${}^2E_6(2)$, ${}^2F_4(2)'$
17	A_{17}, A_{18}, A_{19} ; $J_3, He, Fi_{23}, Fi'_{24}$; $L_2(q)$, $q = 2^4, 17^n$, $2 \cdot 17^n \pm 1$ which is a prime, $n \geq 1$, $S_4(4)$, $S_8(2)$, $F_4(2)$, $O_8^-(2)$, $O_{10}^-(2)$, ${}^2E_6(2)$
19	A_{19}, A_{20}, A_{21} ; $J_1, J_3, O'N, Th, HN$; $L_2(q)$, $q = 19^n$, $2 \cdot 19^n - 1$ which is a prime, $n \geq 1$, $L_3(7)$, $U_3(2^3)$, $R(3^3)$, ${}^2E_6(2)$
37	A_{37}, A_{38}, A_{39} ; J_4, Ly ; $L_2(q)$, $q = 37^n$, $2 \cdot 37^n - 1$ which is a prime, $n \geq 1$, $U_3(11)$, $R(3^3)$, ${}^2F_4(2^3)$
73	A_{73}, A_{74}, A_{75} ; $L_2(q)$, $q = 73^n$, $2 \cdot 73^n - 1$ which is a prime, $n \geq 1$, $L_3(2^3)$, $S_6(2^3)$, $G_2(q)$, $q = 2^3, 3^2$; $F_4(3)$, $E_6(2)$, $E_7(2)$, $U_3(3^2)$, ${}^3D_4(3)$
109	$A_{109}, A_{110}, A_{111}$; $L_2(q)$, $q = 109^n$, $2 \cdot 109^n - 1$ which is a prime, $n \geq 1$, ${}^2F_4(2^3)$
$p = 2^m + 1$, $m = 2^s$	A_p, A_{p+1}, A_{p+2} ; $L_2(q)$, $q = 2^m, p^k$, $2 \cdot p^k \pm 1$ which is a prime, $s \geq k \geq 1$, $S_a(2^b)$, $a = 2^{c+1}$ and $b = 2^d$, $c \geq 1$, $c + d = s$, $F_4(2^e)$, $e \geq 1$, $4e = 2^s$, $O_{2(m+1)}^-(2)$, $s \geq 2$, $O_a^-(2^b)$, $c \geq 2$, $c + d = s$
Other	A_p, A_{p+1}, A_{p+2} ; $L_2(q)$, $q = p^k$, $2 \cdot p^k - 1$ which is a prime, $k \geq 1$

we arrive at a contradiction. Moreover, we have $Z(G/H) = 1$. For any $xH \in G/H$ and $xH \notin K/H$, xH induces an automorphism of K/H and this automorphism

is trivial if and only if $xH \in Z(G/H)$. Therefore, $G/K \leq \text{Out}(K/H)$ and since $Z(G/H) = 1$, (2) follows. $\square$

DEFINITION 2.11. A group G is called a C_{pp} group if the centralizers of its elements of order p in G are p -groups.

LEMMA 2.12 (see [9]). *Let p be a prime and $p = 2^\alpha 3^\beta + 1$, $\alpha \geq 0$ and $\beta \geq 0$. Then any finite simple C_{pp} group is given by Table 2.1.*

3. Characterization of some alternating and symmetric groups. In the sequel, we suppose that $p = 2^\alpha 3^\beta + 1$, where $\alpha \geq 1$, $\beta \geq 0$, and $p \geq 7$ is a prime number.

LEMMA 3.1. *Let G be a finite group and let M be A_n , where $n = p, p+1$, or $p+2$, or S_n , where $n = p, p+1$. If $\text{OC}(G) = \text{OC}(M)$, then G is neither a Frobenius group nor a 2-Frobenius group.*

PROOF. If G is a Frobenius group, then by Lemma 2.5, $\text{OC}(G) = \{|H|, |K|\}$, where K and H are Frobenius kernel and Frobenius complement of G , respectively. Since $|H| \mid |K| - 1$, we have $|H| < |K|$. Therefore, $2 \nmid |H|$, and hence $2 \mid |K|$. So, $|H| = p, |K| = |G|/p$. We claim that there exists a prime p' such that $3n/4 < p'$. Note that $p \leq n$, and hence $p'^2 \nmid |A_n|$. Let $\beta(n)$ be the number of prime numbers less than or equal to n . In fact, by [22, Theorem 2] we have

$$\frac{n}{\log n - 1/2} < \beta(n) < \frac{n}{\log n - 3/2}, \quad (3.1)$$

where $n \geq 67$. Thus

$$\beta(n) - \beta\left(\frac{3n}{4}\right) > \frac{n}{\log n - 1/2} - \frac{3n/4}{\log(3n/4) - 3/2}. \quad (3.2)$$

When $n \geq 405$, we get $\beta(n) - \beta(3n/4) > 1$, and for $n < 405$, we can immediately obtain the result by checking the table of prime numbers. Now let P' be the p' -Sylow subgroup of K . Since K is nilpotent, $P' \triangleleft G$. Then $p \mid p' - 1$, by Lemma 2.7, which is a contradiction since $p' < p$. Therefore, G is not a Frobenius group.

Now let G be a 2-Frobenius group. By Lemma 2.6, there is a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that $|K/H| = p$ and $|G/K| < p$. So, $|H| \neq 1$ since $|G| = |G/K| \cdot |K/H| \cdot |H|$. Since $2 \mid |H|$, let p' be as above and let P' be the p' -Sylow subgroup of H . Now, $p \mid p' - 1$, which is impossible. Hence, G is not a 2-Frobenius group. $\square$

LEMMA 3.2. *Let G be a finite group and $M = A_n$, where $n = p, p+1$, or $p+2$, or S_n , where $n = p, p+1$. If $\text{OC}(G) = \text{OC}(M)$, then G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that H and G/K are π_1 -groups and K/H is a simple group. Moreover, the odd-order component of M is equal to an odd-order component of K/H . In particular, $t(\Gamma(K/H)) \geq 2$. Also $|G/H|$ divides $|\text{Aut}(K/H)|$, and in fact $G/H \leq \text{Aut}(K/H)$.*

PROOF. The first part of the lemma follows from the above lemmas since the prime graph of M has two prime graph components. For primes p and q , if K/H has an element of order pq , then G has one. Hence, by the definition of prime graph component, the odd-order component of G must be an odd-order component of K/H . Since $K/H \triangleleft G/H$ and $C_{G/H}(K/H) = 1$, we have

$$G/H = \frac{N_{G/H}(K/H)}{C_{G/H}(K/H)} \cong T, \quad T \leq \text{Aut}(K/H). \quad (3.3)$$

□

THEOREM 3.3. *Let $p = 2^{\alpha}3^{\beta} + 1$, where $\alpha \geq 1$, $\beta \geq 0$, and $p \geq 7$ is a prime number. Let $M = A_n$, where $n = p, p+1, p+2$. Then $\text{OC}(G) = \text{OC}(M)$ if and only if $G \cong M$.*

PROOF. By Lemma 3.2, G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that $\pi(H) \cup \pi(G/K) \subset \pi_1$, K/H is a non-abelian simple group, $t(\Gamma(K/H)) \geq 2$, and the odd-order component of M is an odd-order component of K/H . Therefore, K/H is a finite simple C_{pp} group. Now using Table 2.1, we consider each possibility of K/H separately.

In the sequel, we frequently use the results of [28, Table I] and [18, Tables 2, 3].

STEP 1. Let $p = 7, 13, 17, 19, 37, 73$, or 109 .

Since the proofs of these cases are similar, we state only one of them, say $p = 13$. Using Table 2.1, we have

- (1) $K/H \cong \text{Suz}$ or Fi_{22} . It is a contradiction since $3^7 \mid |\text{Suz}|$ and $3^9 \mid |\text{Fi}_{22}|$ but $3^7 \nmid |A_n|$, where $n = 13, 14, 15$;
- (2) $K/H \cong L_2(27)$, $L_2(25)$, $L_3(3)$, $L_4(3)$, $\text{Sz}(8)$, ${}^2F_4(2)'$, or $U_3(4)$. If $K/H \cong L_2(27)$, then $|G|/|K/H| = |H| \cdot |G/K| \neq 1$. By Lemma 2.6, $|G/K| \mid |\text{Out}(K/H)| = 6$. So, $|H| \neq 1$. Let P be the 5-Sylow subgroup of H . But since H is nilpotent, $P \triangleleft G$. Hence, $13 \mid (|P| - 1)$, which is a contradiction. Other cases are similar;
- (3) $K/H \cong L_2(13^r)$ or $L_2(2 \cdot 13^r - 1)$, where $2 \cdot 13^r - 1$ is a prime, $r \geq 1$. Note that $13^2 \nmid |G|$, hence $r = 1$. So, $K/H \cong L_2(13)$ or $L_2(25)$, and we can proceed similar to (2);
- (4) $K/H \cong O_7(3)$. It is a contradiction since $3^9 \mid |O_7(3)|$ but $3^9 \nmid |A_n|$;
- (5) $K/H \cong S_4(5)$ or $S_6(3)$. It is a contradiction since $5^4 \mid |S_4(5)|$ but $5^4 \nmid |A_n|$. Also $3^9 \mid |S_6(3)|$ but $3^9 \nmid |A_n|$;
- (6) $K/H \cong O_8^+(3)$. It is a contradiction since $3^{12} \mid |O_8^+(3)|$ but $3^{12} \nmid |A_n|$;
- (7) $K/H \cong G_2(3)$ or $G_2(8)$. If $K/H \cong G_2(3)$, then we get a contradiction since for $n = 13, 14$ we have $3^6 \mid |G_2(3)|$ but $3^6 \nmid |A_n|$. For $n = 15$, since $|\text{Out}(G_2(3))| = 2$, we have $|H| \neq 1$. Now we proceed similar to (2). If $K/H \cong G_2(8)$, then we get a contradiction since $2^{18} \mid |G_2(8)|$ but $2^{18} \nmid |A_n|$;
- (8) $K/H \cong F_4(2)$. It is a contradiction since $17 \mid |F_4(2)|$ but $17 \nmid |A_n|$;
- (9) $K/H \cong U_3(23)$. It is a contradiction since $23 \mid |U_3(23)|$ but $23 \nmid |A_n|$;

- (10) $K/H \cong {}^3D_4(2)$ or ${}^2E_6(2)$. It is a contradiction since $2^{12} \nmid |A_n|$. Also $19 \nmid |A_n|$;
- (11) K/H is an alternating group, namely A_{13} , A_{14} , or A_{15} .

First suppose that $n = 13$. Since $|K/H| \leq |A_{13}|$, $K/H \cong A_{13}$. But $|G| = |A_{13}|$, and hence $H = 1$ and $K = G \cong A_{13}$. If $n = 14$, then $K/H \cong A_{13}$ or A_{14} . But if $r \neq 6$, then $\text{Aut}(A_r) = S_r$, and hence $|\text{Out}(A_r)| = 2$. If $K/H \cong A_{13}$, then $|G/K| \mid 2$, and hence $|H| \neq 1$. Now we get a contradiction similar to (2). Therefore, $K/H \cong A_{14}$, and hence $G \cong A_{14}$. If $n = 15$, we do similarly.

STEP 2. Let $p = 2^m + 1$, where $m = 2^s$.

Using Table 2.1, we have

- (i) $K/H \cong L_2(2^m)$. Note that for every m we have $|L_2(2^m)| \mid |G|$. Using Lemma 2.6, $|G/K| \mid |\text{Out}(K/H)|$. Also $|\text{Out}(L_2(2^m))| = m$. Hence, $|H| \neq 1$. Now let p' be a prime number less than p such that

$$p' \parallel \frac{|A_n|}{m|K/H|}. \quad (3.4)$$

Let P' be the p' -Sylow subgroup of H . Since H is nilpotent, $P' \triangleleft G$. Hence, $p \mid (|P'| - 1)$, which is a contradiction;

- (ii) $K/H \cong L_2(p^k)$ or $L_2(2p^k \pm 1)$, where $2p^k \pm 1$ is a prime and $1 \leq k \leq s$. We know that $p \parallel |A_n|$, hence $k = 1$. Now we proceed similar to (i);
- (iii) $K/H \cong S_a(2^b)$, where $a = 2^{c+1}$ and $b = 2^d$, $c \geq 1$, $c + d = s$. Let $q = 2^b$ and $f = 2^c$. Then $p = q^f + 1$ and we have

$$|S_a(2^b)| = q^{f^2}(q^f - 1)(q^f + 1)\prod_{i=1}^{f-1}(q^i - 1)(q^i + 1). \quad (3.5)$$

Each factor of the form $(q^j \pm 1)$ is less than or equal to p and therefore divides $|A_n|$. Also $q^{f^2} = (2^m)^f \leq 2^{m^2} \leq 2^{2^m}$. Hence, $|S_a(2^b)| \mid |A_n|$. But $|\text{Out}(S_a(2^b))| = b$. Then $|H| \neq 1$ and we can proceed similar to (i);

- (iv) $K/H \cong F_4(2^e)$, where $e \geq 1$, $4e = 2^s$, or $O_{2(m+1)}^-(2)$, where $s \geq 2$, or $O_a^-(2^b)$, where $c \geq 2$, $c + d = s$. Again this part is similar to (iii);
- (v) $K/H \cong A_p, A_{p+1}, A_{p+2}$.

First suppose that $n = p$. Since $|K/H| \leq |A_p|$, $K/H \cong A_p$. But $|G| = |A_p|$, and hence $H = 1$ and $K = G \cong A_p$. If $n = p + 1$, then $K/H \cong A_p$ or A_{p+1} . But if $r \neq 6$, then $\text{Aut}(A_r) = S_r$, and hence $|\text{Out}(A_r)| = 2$. If $K/H \cong A_p$, then $|G/K| \mid 2$, and hence $|H| \neq 1$. Now we get a contradiction similar to (i). Therefore, $K/H \cong A_{p+1}$, and hence $G \cong A_{p+1}$. If $n = p + 2$, we do similarly.

STEP 3. For other primes p , we have $K/H \cong A_p, A_{p+1}, A_{p+2}; L_2(q)$, where $q = p^k, 2p^k - 1$ which is a prime, $k \geq 1$.

In fact the proof of this step is exactly similar to that of Step 2 and we omit it for convenience. $\square$

THEOREM 3.4. *If G is a non-abelian finite group with connected prime graph, then G is not characterizable with its order component.*

PROOF. Clearly, $\text{OC}(G) = \text{OC}(\mathbb{Z}_{|G|})$, but $G \not\cong \mathbb{Z}_{|G|}$. □

COROLLARY 3.5. *Every simple group with one component (see [28, Table I]) is not characterizable with this method.*

THEOREM 3.6. *Let n be a positive integer. If there exist at least two non-isomorphic abelian groups of order n , then abelian groups of order n are not characterizable with their order component.*

PROOF. The proof is obvious. □

REMARK 3.7. It was a conjecture that every finite simple group M , where $\Gamma(M)$ is not connected, is characterizable with its order components. But the following example is a counterexample.

EXAMPLE 3.8. If q is an odd-prime power and $n = 2^k \geq 4$, then $\text{OC}(S_{2n}(q)) = \text{OC}(O_{2n+1}(q))$, but obviously $S_{2n}(q) \not\cong O_{2n+1}(q)$.

THEOREM 3.9. *Let $p = 2^\alpha 3^\beta + 1$, where $\alpha \geq 1$, $\beta \geq 0$, and $p \geq 7$ is a prime number. Let $M = S_n$, where $n = p, p + 1$. Then $\text{OC}(G) = \text{OC}(M)$ if and only if $G \cong M$.*

PROOF. Similar to the proof of [Theorem 3.3](#), since G is a C_{pp} group, we have $K/H \cong A_n$. Now using [Lemma 3.2](#), we have

$$A_n \leq \frac{G}{H} \leq \text{Aut}(A_n) = S_n. \quad (3.6)$$

Therefore, $G/H \cong A_n$ or $\text{Aut}(A_n) = S_n$. If $G/H \cong A_n$, then $|H| = 2$ and $H \triangleleft G$. Hence, $H \subseteq Z(G) = 1$, which is a contradiction. Therefore, $G/H \cong S_n$, and since $|G| = |S_n|$, we have $G \cong S_n$. □

4. Some related results

REMARK 4.1. It is a well known conjecture of J. G. Thompson that if G is a finite group with $Z(G) = 1$ and M is a non-abelian simple group satisfying $N(G) = N(M)$, then $G \cong M$.

We can generalize this conjecture for the groups under discussion by our characterization of these groups.

COROLLARY 4.2. *Let G be a finite group with $Z(G) = 1$ and let M be A_p , A_{p+1} , A_{p+2} , S_p , or S_{p+1} . If $N(G) = N(M)$, then $G \cong M$.*

PROOF. By [Lemmas 2.8](#) and [2.9](#), if G and M are two finite groups satisfying the conditions of [Corollary 4.2](#), then $\text{OC}(G) = \text{OC}(M)$. So, [Theorems 3.3](#) and [3.9](#) imply this corollary. □

REMARK 4.3. Shi and Bi in [26] put forward the following conjecture.

SHI'S CONJECTURE. *Let G be a group and M a finite simple group. Then $G \cong M$ if and only if*

- (i) $|G| = |M|$,
- (ii) $\pi_e(G) = \pi_e(M)$, where $\pi_e(G)$ denotes the set of orders of elements in G .

This conjecture is valid for sporadic simple groups [24], groups of alternating type [27], and some simple groups of Lie type [23, 25, 26]. As a consequence of Theorems 3.3 and 3.9, we prove a generalization of this conjecture for the groups under discussion.

COROLLARY 4.4. *Let G be a finite group and let M be A_p , A_{p+1} , A_{p+2} , S_p , or S_{p+1} . If $|G| = |M|$ and $\pi_e(G) = \pi_e(M)$, then $G \cong M$.*

PROOF. By assumption, we must have $\text{OC}(G) = \text{OC}(M)$. Thus the corollary follows by Theorems 3.3 and 3.9. $\square$

REFERENCES

- [1] G. Y. Chen, *On structure of Frobenius groups and 2-Frobenius groups*, J. Southwest China Normal Univ. **20** (1995), no. 5, 485–487.
- [2] ———, *A new characterization of $E_8(q)$* , J. Southwest China Normal Univ. **21** (1996), no. 3, 215–217.
- [3] ———, *A new characterization of $G_2(q)$, [$q \equiv 0 \pmod{3}$]*, J. Southwest China Normal Univ. (1996), 47–51.
- [4] ———, *A new characterization of sporadic simple groups*, Algebra Colloq. **3** (1996), no. 1, 49–58.
- [5] ———, *On Thompson's conjecture*, J. Algebra **185** (1996), no. 1, 184–193.
- [6] ———, *A new characterization of Suzuki-Ree group*, Sci. China Ser. A **40** (1997), no. 8, 807–812.
- [7] ———, *A new characterization of $\text{PSL}_2(q)$* , Southeast Asian Bull. Math. **22** (1998), no. 3, 257–263.
- [8] ———, *Further reflections on Thompson's conjecture*, J. Algebra **218** (1999), no. 1, 276–285.
- [9] Z. M. Chen and W. J. Shi, *On simple C_{pp} groups*, J. Southwest China Teachers Univ. (Nat. Sci) **18** (1993), no. 3, 249–256.
- [10] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, *Atlas of Finite Groups*, Oxford University Press, Oxford, 1985.
- [11] K. W. Gruenberg and K. W. Roggenkamp, *Decomposition of the augmentation ideal and of the relation modules of a finite group*, Proc. London Math. Soc. (3) **31** (1975), no. 2, 149–166.
- [12] A. Iranmanesh and S. H. Alavi, *A new characterization of A_p where p and $p-2$ are primes*, Korean J. Comput. Appl. Math. **8** (2001), no. 3, 665–673.
- [13] A. Iranmanesh, S. H. Alavi, and B. Khosravi, *A characterization of $\text{PSL}(3, q)$ for $q = 2^m$* , Acta Math. Sin. (Engl. Ser.) **18** (2002), no. 3, 463–472.
- [14] ———, *A characterization of $\text{PSL}(3, q)$ where q is an odd prime power*, J. Pure Appl. Algebra **170** (2002), no. 2-3, 243–254.
- [15] A. Iranmanesh and B. Khosravi, *A characterization of $F_4(q)$ where $q = 2^n$ ($n > 1$)*, Far East J. Math. Sci. **2** (2000), no. 6, 853–859.

- [16] A. Iranmanesh, B. Khosravi, and S. H. Alavi, *A Characterization of $\text{PSU}_3(q)$ for $q > 5$* , S. E. Bull. Math. **26** (2002), no. 1, 33–44.
- [17] K. Koike, *On some groups which are determined by their subgroup-lattices*, Tokyo J. Math. **6** (1983), no. 2, 413–421.
- [18] A. S. Kondrat'ev, *Prime graph components of finite simple groups*, Math. USSR-Sb. **67** (1990), no. 1, 235–247.
- [19] A. S. Kondrat'ev and V. D. Mazurov, *Recognition of alternating groups of prime degree from their element orders*, Siberian Math. J. **41** (2000), no. 2, 294–302.
- [20] T. Oyama, *On the groups with the same table of characters as alternating groups*, Osaka J. Math. **1** (1964), no. 1, 91–101.
- [21] C. E. Praeger and W. J. Shi, *A characterization of some alternating and symmetric groups*, Comm. Algebra **22** (1994), no. 5, 1507–1530.
- [22] J. B. Rosser and L. Schoenfeld, *Approximate formulas for some functions of prime numbers*, Illinois J. Math. **6** (1962), 64–94.
- [23] W. J. Shi, *A new characterization of some simple groups of Lie type*, Classical Groups and Related Topics (Beijing, 1987), Contemp. Math., vol. 82, American Mathematical Society, Rhode Island, 1989, pp. 171–180.
- [24] ———, *A new characterization of the sporadic simple groups*, Group Theory (Singapore, 1987), de Gruyter, Berlin, 1989, pp. 531–540.
- [25] ———, *Pure quantitative characterization of finite simple groups. I*, Progr. Natur. Sci. (English Ed.) **4** (1994), no. 3, 316–326.
- [26] W. J. Shi and J. X. Bi, *A characteristic property for each finite projective special linear group*, Groups—Canberra 1989, Lecture Notes in Math., vol. 1456, Springer, Berlin, 1990, pp. 171–180.
- [27] ———, *A new characterization of the alternating groups*, Southeast Asian Bull. Math. **16** (1992), no. 1, 81–90.
- [28] J. S. Williams, *Prime graph components of finite groups*, J. Algebra **69** (1981), no. 2, 487–513.
- [29] A. V. Zavarnitsin, *Recognition of alternating groups of degrees $r + 1$ and $r + 2$ for prime r and of a group of degree 16 by the set of their element orders*, Algebra i Logika **39** (2000), no. 6, 648–661 (Russian), translated in Algebra and Logic **39** (2000), no. 6, 370–377.

Amir Khosravi: Faculty of Mathematical Sciences and Computer Engineering, University for Teacher Education, 599 Taleghani Ave., Tehran 15614, Iran

Behrooz Khosravi: 241 Golnaz Street, Velenjak, Tehran 19847, Iran

E-mail address: khosravibbb@yahoo.com