

BERGMAN SPACES OF TEMPERATURE FUNCTIONS ON A CYLINDER

MARCOS LÓPEZ-GARCÍA

Received 19 April 2002

We define the weighted Bergman space $b_{\beta}^p(S_T)$ consisting of temperature functions on the cylinder $S_T = S^1 \times (0, T)$ and belonging to $L^p(\Omega_T, t^{\beta} dx dt)$, where $\Omega_T = (0, 2) \times (0, T)$. For $\beta > -1$ we construct a family of bounded projections of $L^p(\Omega_T, t^{\beta} dx dt)$ onto $b_{\beta}^p(S_T)$. We use this to get, for $1 < p < \infty$ and $1/p + 1/p' = 1$, a duality $b_{\beta}^p(S_T)^* = b_{\beta'}^p(S_T)$, where β' depends on p and β .

2000 Mathematics Subject Classification: 46E22, 35K05.

1. Introduction. For \mathbb{D} , the open unit disk in the complex plane \mathbb{C} , the classic Bergman space L_a^p is the subspace of holomorphic functions $f: \mathbb{D} \rightarrow \mathbb{C}$ such that $f \in L^p(\mathbb{D})$. It can be verified by the mean value theorem and Hölder inequality that L_a^p is a closed subspace of $L^p(\mathbb{D})$. This implies the existence of an orthogonal projection P from $L^2(\mathbb{D})$ onto L_a^2 , which is called the Bergman projection. The projection P can be written as an integral operator

$$Pf(z) = \int_{\mathbb{D}} K(z, w) f(w) dw, \quad (1.1)$$

for all $f \in L_a^2$, where $K(z, w)$ is the so-called Bergman reproducing kernel of L_a^2 .

The theory of Bergman spaces has a long history. It goes back to the work of Bergman [3], who gave the first treatment of $L_a^2(\Omega)$. Today there are rich theories describing the Bergman spaces in various domains and their operators. Two of the most important classes of operators in the Bergman space theory are the Toeplitz and Hankel operators, which are defined in terms of the Bergman projection P . This theory was mainly developed in the late 1980s. For a very nice exposition of the $L^p(\mathbb{D})$ -theory of Bergman spaces, operators defined on them, and further historical references, we refer to Axler [1], Zhu [10], and a more modern approach in [2].

In this paper, we define weighted Bergman-type spaces $b_{\beta}^p(S_T)$ consisting of temperature functions on the cylinder $S_T = S^1 \times (0, T)$ and belonging to $L^p(\Omega_T, t^{\beta} dx dt)$, where $\Omega_T = (0, 2) \times (0, T)$. As in the holomorphic case, we prove that $b_{\beta}^p(S_T)$ is a Banach space. Therefore, there exist the Bergman projection and the corresponding reproducing kernel in this setting. Since the Bergman projection $P: L^2(\Omega_T) \rightarrow b^2(S_T)$ is an orthogonal projection, then it is

bounded, but the boundedness of P on L^p is not obvious at all. We will construct a family of reproducing kernels and bounded projections P_α of $L^p(\Omega_T, t^\beta dx dt)$ onto $b^p_\beta(S_T)$, for $p > 1$ and $p > (1 + \beta)/(1 + \alpha)$. The proof of the boundedness of these projections is based on a version of the Schur’s test, and the theory of Fourier multipliers. Using the fact that P_α is bounded we obtain the duality between the Bergman spaces.

The main results of this paper are the following theorems.

THEOREM 1.1. *Let $\alpha, \beta > -1$. If $p > \max(1, (1 + \beta)/(1 + \alpha))$, the operator $P_\alpha : L^p_\beta(\Omega_T) \rightarrow b^p_\beta(S_T)$ given by*

$$P_\alpha u(z) = \int_{\Omega_T} N_\alpha(z, w) u(w) \tau^\alpha dw, \quad z \in \Omega_T, \tag{1.2}$$

is a continuous projection onto $b^p_\beta(S_T)$, where

$$N_\alpha(z, w) = \frac{1 + \alpha}{2T^{1+\alpha}} + \sum_{m \in \mathbb{Z}^*} \frac{2^\alpha \pi^{2(1+\alpha)} m^{2(1+\alpha)}}{\gamma(1 + \alpha, 2\pi^2 m^2 T)} e^{-\pi^2 m^2 (t+\tau) + \pi m i(x-\gamma)}. \tag{1.3}$$

THEOREM 1.2. *Let $\alpha, \beta > -1$. If $p > \max(1, (1 + \beta)/(1 + \alpha))$, then $(b^p_\beta(S_T))^* = b^{p'}_{(\alpha-\beta/p)p'}(S_T)$ with respect to the duality*

$$\langle u, v \rangle_\alpha = \int_{\Omega_T} u(z) v(z) t^\alpha dx dt. \tag{1.4}$$

THEOREM 1.3. *Let $n, d \geq 1$ and $p > 1$. If $u \in b^p(S_T)$, then $t^{n/2}(\partial^n u / \partial x^n)$, $t^d(\partial^d u / \partial t^d) \in L^p(\Omega_T)$. Furthermore, there are constants $C_n, C_d > 0$ such that*

$$\begin{aligned} \left\| t^{n/2} \frac{\partial^n u}{\partial x^n} \right\|_{L^p(\Omega_T)} &\leq C_n \|u\|_{b^p(S_T)}, \\ \left\| t^d \frac{\partial^d u}{\partial t^d} \right\|_{L^p(\Omega_T)} &\leq C_d \|u\|_{b^p(S_T)}, \end{aligned} \tag{1.5}$$

for all $u \in b^p(S_T)$.

The paper is organized as follows. After some preliminaries in [Section 2](#), we define the Bergman space $b^p_\beta(S_T)$ in [Section 3](#). In [Section 4](#), we define a family of reproducing kernels and projections P_α . Finally, in [Section 5](#) we prove the boundedness of the projections P_α and the duality between the Bergman spaces.

2. Notation and preliminary results. Throughout this paper we will use the following notation: the conjugate exponent of $p > 1$ will be denoted by p' ,

we will write $z = (x, t)$, $w = (y, \tau)$, $dz = dxdt$, $dw = dyd\tau$, $S^1 = \{e^{\pi i\theta} : \theta \in [0, 2]\}$, and $\mathbb{Z}^* = \{n \in \mathbb{Z} : n \neq 0\}$. For $u \in L^1(\mathbb{R})$, we define its Fourier transform as

$$(Fu)(\zeta) = \int_{-\infty}^{\infty} u(x)e^{-2\pi i x \zeta} dx. \tag{2.1}$$

For $\Omega \subset \mathbb{R}_+^2$ an open set, let

$$H(\Omega) = \left\{ u \in C^2(\Omega) : \frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t} \text{ on } \Omega \right\}. \tag{2.2}$$

We will call the elements of $H(\Omega)$ temperature functions. $K(x, t)$ will denote the Gauss-Weierstrass kernel. For $t > 0$, let

$$\begin{aligned} \theta(x, t) &= \sum_{n \in \mathbb{Z}} K(x + 2n, t) = \frac{1}{2} \sum_{n \in \mathbb{Z}} e^{-\pi^2 n^2 t + \pi n i x}, \\ \varphi(x, t) &= -2 \frac{\partial \theta}{\partial x}(x, t), \end{aligned} \tag{2.3}$$

and for $t \leq 0$, let $K = \theta = \varphi = 0$ (see [8]). Moreover,

$$\int_0^2 \theta(x, t) dx = \int_{-\infty}^{\infty} K(x, t) dx = 1, \quad \forall t > 0. \tag{2.4}$$

Let $Q = (0, 1) \times (0, 1)$, $\Gamma = \Gamma_1 \cup \Gamma_2 \cup \Gamma_3$ (the parabolic boundary of Q), where $\Gamma_1 = \{0\} \times [0, 1)$, $\Gamma_2 = \{1\} \times [0, 1)$, and $\Gamma_3 = (0, 1) \times \{0\}$, and let λ be the one-dimensional Lebesgue measure on Γ .

We consider the heat kernel $\tilde{K}(x, t; \xi, \tau)$ on $Q \times \Gamma$ defined as follows:

$$\tilde{K}(x, t; \xi, \tau) = \begin{cases} \varphi(x, t - \tau), & \xi = 0, 0 \leq \tau < 1, \\ \varphi(1 - x, t - \tau), & \xi = 1, 0 \leq \tau < 1, \\ \theta(x - \xi, t) - \theta(x + \xi, t), & \tau = 0, 0 < \xi < 1. \end{cases} \tag{2.5}$$

It is well known that if $u \in H(Q) \cap C(\overline{Q})$, then (see [4])

$$u(x, t) = \int_{\Gamma} \tilde{K}(x, t; \xi, \tau) u(\xi, \tau) d\lambda(\xi, \tau), \quad \forall (x, t) \in Q. \tag{2.6}$$

Conversely, if $v \in C(\Gamma)$, then

$$u(x, t) = \int_{\Gamma} \tilde{K}(x, t; \xi, \tau) v(\xi, \tau) d\lambda(\xi, \tau) \tag{2.7}$$

is a temperature function on Q .

REMARK 2.1. Clearly $\tilde{K}(x, t; \cdot) \in L^1(\Gamma, d\lambda)$ for all $(x, t) \in Q$. This follows from (2.7) with $v \equiv 1$.

Let $R = (a, b) \times (c, d) \subset \mathbb{R}_+^2$ be a rectangle such that $d - c = (b - a)^2$. If $u \in H(R) \cap C(\bar{R})$ then $u \circ \Psi \in H(Q) \cap C(\bar{Q})$, where the mapping $\Psi : Q \rightarrow R$ is defined by

$$\Psi(\xi, \tau) = ((b - a)\xi + a, (d - c)\tau + c). \tag{2.8}$$

By (2.6) we have

$$u(x, t) = \int_{\Gamma_R} \tilde{K}(\Psi^{-1}(x, t); \Psi^{-1}(\xi, \tau)) u(\xi, \tau) d\lambda_R(\xi, \tau), \quad \forall (x, t) \in R, \tag{2.9}$$

where $\Gamma_R = \Psi(\Gamma) = \Psi(\Gamma_1) \cup \Psi(\Gamma_2) \cup \Psi(\Gamma_3)$ and λ_R is the one-dimensional Lebesgue measure normalized on each segment of Γ_R .

In [5] it was proved that there is a constant $C > 0$ such that

$$\theta(x, t) \leq C(1 + t)K(x, t), \quad -1 < x < 1. \tag{2.10}$$

In particular for $0 < t \leq T < \infty$ there is a constant $C_T > 0$ such that

$$\theta(x, t) \leq C_T K(x, t), \quad -1 < x < 1. \tag{2.11}$$

Now, suppose f is a 2-periodic continuous function on \mathbb{R} . Then u is a temperature function on \mathbb{R}_+^2 , 2-periodic in the variable x , and $u(x, t) \rightarrow f(x)$ as $t \rightarrow 0$ uniformly on $[0, 2]$ if and only if (see [8, Chapter 5, Theorem 8])

$$u(x, t) = \int_0^2 \theta(x - y, t) f(y) dy. \tag{2.12}$$

REMARK 2.2. Since $f(x)$ is continuous on \mathbb{R} and $u(x, t) \rightarrow f(x)$ uniformly on $[0, 2]$, the continuity of u at $t = 0$ follows.

On the other hand, if

$$u(x, t) = \int_0^2 \theta(x - y, t) f(y) dy = (K(\cdot, t) * f)(x), \tag{2.13}$$

by Minkowski's integral inequality, we have

$$\|u(\cdot, t)\|_{L^p(S^1)} \leq \|f\|_{L^p(S^1)}, \quad \forall t > 0. \tag{2.14}$$

The following result, proved in [6], will be useful in this paper.

LEMMA 2.3. *Let $R = (a, b) \times (c, d) \subset \mathbb{R}_+^2$ be such that $d - c = (b - a)^2$. If $u \in H(R) \cap C(\bar{R})$ and (x_0, t_0) is the midpoint of the upper boundary of R , then*

$$|u(x_0, t_0)|^p \leq \frac{C_p}{|R|} \iint_R |u(x, t)|^p dx dt, \tag{2.15}$$

where $|R|$ is the area of R and C_p is a constant depending only on $p > 0$.

Next, we prove a variant of Schur’s lemma (see [9]). This result provides a sufficient condition for the boundedness of an integral operator T defined on $L^p(\Omega, d\mu)$, $1 < p < \infty$.

LEMMA 2.4. *Let $1 < p < \infty$. Let (Ω, μ) be a measure space with μ a σ -finite measure, and let $N : \Omega \times \Omega \rightarrow \mathbb{C}$ and $G : \Omega \rightarrow \mathbb{R}_+$ be measurable functions. For a measurable function f , define*

$$Tf(w_1) = \int_{\Omega} N(w_1, w_2) f(w_2) d\mu(w_2). \tag{2.16}$$

Assume that there exist measurable functions $h, g : \Omega \rightarrow \mathbb{R}_+$ and constants $a, b \geq 0$ such that

$$\int_{\Omega} |N(w_1, w_2)| h(w_2)^{p'} d\mu(w_2) \leq (ag(w_1))^{p'}, \quad \mu\text{-a.e.}, \tag{2.17}$$

$$\int_{\Omega} |N(w_1, w_2)| g(w_1)^p G(w_1) d\mu(w_1) \leq (bh(w_2))^p G(w_2), \quad \mu\text{-a.e.} \tag{2.18}$$

Then $T : L^p(\Omega, Gd\mu) \rightarrow L^p(\Omega, Gd\mu)$ is a bounded operator and $\|T\| \leq ab$.

PROOF. By Hölder’s inequality and (2.17) we have

$$|Tf(w_1)| \leq ag(w_1) \left(\int_{\Omega} |N(w_1, w_2)| \left| \frac{f(w_2)}{h(w_2)} \right|^p d\mu(w_2) \right)^{1/p}. \tag{2.19}$$

By Tonelli’s theorem and (2.18) we have

$$\int_{\Omega} |Tf(w_1)|^p G(w_1) d\mu(w_1) \leq (ab)^p \int_{\Omega} |f(w_2)|^p G(w_2) d\mu(w_2). \tag{2.20}$$

That is, $\|Tf\|_{L^p(\Omega, Gd\mu)} \leq ab \|f\|_{L^p(\Omega, Gd\mu)}$. □

LEMMA 2.5. *Let $\alpha, \beta > -1$ and $p > \max(1, (1 + \beta)/(1 + \alpha))$. For every $\delta > 0$ such that*

$$\frac{\beta - \alpha}{p} < \delta < \min\left(\frac{1 + \alpha}{p'}, \frac{1 + \beta}{p}\right), \tag{2.21}$$

the function $h(t) = t^{-\delta}$ satisfies the following inequalities:

$$\begin{aligned} \int_0^T \frac{\tau^\alpha}{(t+\tau)^{1+\alpha}} h(\tau)^{p'} d\tau &\leq Ch(t)^{p'}, \quad \forall t \in (0, T), \\ \int_0^T \frac{\tau^\alpha}{(t+\tau)^{1+\alpha}} h(t)^p t^\beta dt &\leq Ch(\tau)^p \tau^\beta, \quad \forall \tau \in (0, T). \end{aligned} \tag{2.22}$$

PROOF. Let $y \in \mathbb{R}$. Making the change of variable $t = \tau s$, we have

$$\int_0^T \frac{t^{-y}}{(t+\tau)^{1+\alpha}} dt \leq \tau^{-y-\alpha} \int_0^\infty \frac{s^{-y}}{(s+1)^{1+\alpha}} ds = C_{\alpha,y} \tau^{-y-\alpha}, \tag{2.23}$$

where $C_{\alpha,y} = \int_0^\infty (s^{-y}/(s+1)^{1+\alpha}) ds < \infty$ whenever $-\alpha < y < 1$. We obtain (2.22) by letting $y = \delta p' - \alpha$ and $y = \delta p - \beta$, respectively. \square

3. The Bergman space $b_\beta^p(S_T)$. Let $W : \Omega \rightarrow \mathbb{R}^+$ be a measurable function such that $W^{-p'/p} \in L_{\text{loc}}^1(\Omega)$. Denote $L_W^p(\Omega) = L^p(\Omega, W dx dt)$ for $1 \leq p < \infty$.

DEFINITION 3.1. We define the Bergman space $b_W^p(\Omega)$ as the subspace of temperature functions in $L_W^p(\Omega)$. That is, $b_W^p(\Omega) = H(\Omega) \cap L_W^p(\Omega)$.

We will show that $b_W^p(\Omega)$ is a closed subspace of $L_W^p(\Omega)$ and therefore a Banach space. For this aim, we will need the following result.

PROPOSITION 3.2. *Given $1 \leq p < \infty$ and $\mathcal{K} \subset \Omega$ a compact set, there is a constant $C_{\mathcal{K}} > 0$ such that*

$$|u(x, t)| \leq C_{\mathcal{K}} \|u\|_{L_W^p(\Omega)}, \tag{3.1}$$

for all $(x, t) \in \mathcal{K}$, $u \in b_W^p(\Omega)$.

PROOF. Let $\delta = d(\mathcal{K}, \Omega^c) > 0$ and $\mathcal{K}_0 = \{z \in \Omega : d(z, \mathcal{K}) \leq \delta/2\}$. For every $z \in \mathcal{K}$, let R_z be a rectangle such that z is the midpoint of the upper boundary of R_z , with height(R_z) = {base(R_z)}² and $R_z \subset B(z, \delta/2) \subset \mathcal{K}_0$.

By Lemma 2.3 there is a constant $C > 0$ such that

$$\begin{aligned} |u(z)| &\leq \frac{C}{|R_z|} \iint_{R_z} |u(y, \tau)| dy d\tau \\ &\leq \frac{C}{|R_z|} \|u\|_{L_W^p(\Omega)} \left(\iint_{\mathcal{K}_0} W^{-p'/p} dy d\tau \right)^{1/p'}, \end{aligned} \tag{3.2}$$

for all $u \in b_W^p(\Omega)$.

We conclude the proof by choosing rectangles R_z congruent to one another, for every $z \in \mathcal{K}$. \square

REMARK 3.3. If $u_j \rightarrow u$ in $b_W^p(\Omega)$, then $u_j \rightarrow u$ uniformly on each compact subset of Ω .

THEOREM 3.4. For $1 \leq p < \infty$, $b_W^p(\Omega)$ is a closed subspace of $L_W^p(\Omega)$. Therefore $b_W^p(\Omega)$ is a Banach space.

PROOF. Given $u \in L_W^p(\Omega)$, let (u_j) be a sequence in $b_W^p(\Omega)$ such that

$$\|u - u_j\|_{L_W^p(\Omega)} \rightarrow 0. \tag{3.3}$$

We will show that u is a temperature function on Ω (except on a set of measure zero).

Pick $(x_0, t_0) \in \Omega$. Let $R = (a, b) \times (c, d)$, with $d - c = (b - a)^2$, such that $(x_0, t_0) \in R$ and $\bar{R} \subset \Omega$. By Proposition 3.2, there is a constant $C > 0$ such that

$$|u_j(x, t) - u_k(x, t)| \leq C \|u_j - u_k\|_{L_W^p(\Omega)}, \quad \forall (x, t) \in \bar{R}. \tag{3.4}$$

It follows that (u_j) converges uniformly on \bar{R} to a continuous function v .

Since $u_j \in H(R) \cap C(\bar{R})$, (2.9) implies that

$$u_j(x, t) = \int_{\Gamma_R} \tilde{K}(\Psi^{-1}(x, t); \Psi^{-1}(\xi, \tau)) u_j(\xi, \tau) d\lambda_R(\xi, \tau), \quad \text{for } (x, t) \in R. \tag{3.5}$$

By Remark 2.1 and the dominated convergence theorem we have

$$\begin{aligned} v(x, t) &= \lim_{j \rightarrow \infty} u_j(x, t) = \int_{\Gamma_R} \tilde{K}(\Psi^{-1}(x, t); \Psi^{-1}(\xi, \tau)) \lim_{j \rightarrow \infty} u_j(\xi, \tau) d\lambda_R(\xi, \tau) \\ &= \int_{\Gamma_R} \tilde{K}(\Psi^{-1}(x, t); \Psi^{-1}(\xi, \tau)) v(\xi, \tau) d\lambda_R(\xi, \tau). \end{aligned} \tag{3.6}$$

Since the function v is continuous on Γ_R then v is a temperature function on R . On the other hand, $u_j \rightarrow u$ in $L_W^p(\Omega)$ and then some subsequence of (u_j) converges to u almost everywhere on Ω . Therefore $u = v$ a.e. on R and $u \in H(R)$. Since $(x_0, t_0) \in \Omega$ was arbitrary, we conclude that $u \in H(\Omega)$. \square

We will write $b^p(\Omega) = b_W^p(\Omega)$ when $W \equiv 1$. Proposition 3.2 implies that the linear functional $\mathcal{F}_z : b^2(\Omega) \rightarrow \mathbb{C}$ defined by $\mathcal{F}_z(u) = u(z)$ is bounded for all $z \in \Omega$. Hence the Riesz representation theorem shows that there is a function $N : \Omega \times \Omega \rightarrow \mathbb{C}$ such that

$$u(z) = \langle u, N(z, \cdot) \rangle = \int_{\Omega} u(w) N(z, w) dw, \tag{3.7}$$

for all $u \in b^2(\Omega)$, $z \in \Omega$.

The function $N : \Omega \times \Omega \rightarrow \mathbb{C}$ is called the reproducing kernel of $b^2(\Omega)$, some of its properties are the following (see [2]):

- (1) $N(z, \cdot) \in b^2(\Omega)$ for all $z \in \Omega$;
- (2) N is real-valued and symmetric;

(3) if (u_n) is an orthonormal basis of $b^2(\Omega)$, then

$$N(z, w) = \sum_{n=1}^{\infty} u_n(z) \overline{u_n(w)}. \quad (3.8)$$

There is a unique orthogonal projection $P : L^2(\Omega) \rightarrow b^2(\Omega)$ called the Bergman projection. We have

$$Pu(z) = \langle Pu, N(z, \cdot) \rangle = \langle u, N(z, \cdot) \rangle = \int_{\Omega} u(w) N(z, w) dw. \quad (3.9)$$

For $T > 0$, we define

$$\begin{aligned} C(\overline{S_T}) &= \{u \in C(\mathbb{R} \times [0, T]) : u(x, t) = u(x+2, t)\}, \\ H(S_T) &= \{u \in H(\mathbb{R} \times (0, T)) : u(x, t) = u(x+2, t)\}. \end{aligned} \quad (3.10)$$

We will call the elements of $H(S_T)$ temperature functions on the cylinder S_T .

DEFINITION 3.5. For $1 \leq p < \infty$, we define the Bergman space of temperature functions on the cylinder S_T as

$$b_W^p(S_T) = \left\{ u \in H(S_T) : \int_{\Omega_T} |u|^p W dz < \infty \right\}, \quad (3.11)$$

where $\Omega_T = (0, 2) \times (0, T)$, and W is a measurable function such that $W^{-p'/p} \in L_{\text{loc}}^1(\Omega_T)$.

In $b_W^p(S_T)$ we define the following norm:

$$\|u\|_{b_W^p(S_T)} = \left(\int_{\Omega_T} |u|^p W dz \right)^{1/p}. \quad (3.12)$$

Notice that if $u \in b_W^p(S_T)$, then $u|_{\Omega_T} \in L_W^p(\Omega_T)$. Hence we write $b_W^p(S_T) \subset L_W^p(\Omega_T)$.

THEOREM 3.6. For $1 \leq p < \infty$, $b_W^p(S_T)$ is a closed subspace of $L_W^p(\Omega_T)$. Therefore $b_W^p(S_T)$ is a Banach space.

PROOF. Let $u \in L_W^p(\Omega_T)$ and let (u_j) be a sequence in $b_W^p(S_T)$ such that $\|u - u_j\|_{L_W^p(\Omega_T)} \rightarrow 0$. Consider the open set $\Omega = (-1, 3) \times (0, T)$ and the sequence $(u_j|_{\Omega})$. Extend the function $W : \Omega_T \rightarrow \mathbb{R}_+$ to be 2-periodic in the variable x . Then

$$\|u_i - u_j\|_{L_W^p(\Omega)} = 2\|u_i - u_j\|_{L_W^p(\Omega_T)}. \quad (3.13)$$

It follows that $(u_j|_{\Omega})$ is a Cauchy sequence in $b_W^p(\Omega)$. Since $b_W^p(\Omega)$ is a Banach space, there is $v \in b_W^p(\Omega)$ such that $\|v - u_j\|_{L_W^p(\Omega)} \rightarrow 0$. By [Remark 3.3](#), the sequence $(u_j|_{\Omega})$ converges uniformly on compact subsets of Ω . So,

$$v(0, t) = \lim_{j \rightarrow \infty} u_j(0, t) = \lim_{j \rightarrow \infty} u_j(2, t) = v(2, t), \quad (3.14)$$

for all $0 < t < T$. Extend the function v as a 2-periodic function in the variable x . This extension, which we still denote by v , belongs to $b_W^p(S_T)$ and $\|v - u_j\|_{L_W^p(\Omega_T)} \rightarrow 0$. Thus $u = v$ a.e. \square

As in (3.7), it is shown that there is a reproducing kernel of $b^2(S_T)$, that is, a function $N : \Omega_T \times \Omega_T \rightarrow \mathbb{R}$ satisfying

$$u(z) = \int_{\Omega_T} N(z, w) u(w) dw, \tag{3.15}$$

for all $u \in b^2(S_T)$, $z \in \Omega_T$.

If (u_n) is an orthonormal basis of $b^2(S_T)$, then

$$N(z, w) = \sum_{n=1}^{\infty} u_n(z) \overline{u_n(w)}. \tag{3.16}$$

The Bergman projection $P : L^2(\Omega_T) \rightarrow b^2(S_T)$ is the integral operator given by

$$Pu(z) = \int_{\Omega_T} N(z, w) u(w) dw, \quad \forall z \in \Omega_T. \tag{3.17}$$

We will prove that the extension of the Bergman projection P to $L^p(\Omega_T)$ is bounded for all $p > 1$. Actually, we will show that P is bounded on certain weighted Bergman spaces.

From now on, we will be working with weights consisting of powers of the distance of a point to the base of the cylinder, that is, we consider $W_\beta : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ given by $W_\beta(x, t) = t^\beta$. Clearly $W_\beta^{-p'/p} \in L_{\text{loc}}^1(\mathbb{R}_+^2)$.

Let $L_\beta^p(\Omega_T) = L_{W_\beta}^p(\Omega_T)$ and $b_\beta^p(S_T) = b_{W_\beta}^p(S_T)$. If $\beta > -1$, then $W_\beta dx dt$ is a finite measure on Ω_T and

$$L_\beta^p(\Omega_T) \subset L_\beta^1(\Omega_T), \quad b_\beta^p(S_T) \subset b_\beta^1(S_T), \tag{3.18}$$

for all $1 \leq p < \infty$.

We will show that the subspace $H(S_T) \cap C(\overline{S_T})$ is dense in $b_\beta^p(S_T)$, for all $\beta > -1$. Given $u \in b_\beta^p(S_T)$ and $0 < r < T$, we define the function u_r as follows:

$$u_r(x, t) = \int_0^2 \theta(x - y, t) u(y, r) dy. \tag{3.19}$$

From (2.12) and Remark 2.2, we have that $u_r \in H(S_T) \cap C(\overline{S_T})$ and $u_r(x, 0) = u(x, r)$.

The uniqueness of the solution of the heat equation on a finite cylinder (see [8]) yields

$$u_r(x, t) = u(x, t + r), \quad \text{for } 0 \leq t < T - r. \tag{3.20}$$

As before, we have

$$u(x, t+r) = \int_0^2 \theta(x-y, r) u(y, t) dy = (K(\cdot, r) * u(\cdot, t))(x), \quad \text{for } r > 0. \tag{3.21}$$

Minkowski's integral inequality implies

$$\int_0^2 |u(x, t+r)|^p dx \leq \int_0^2 |u(x, t)|^p dx, \quad \text{for } r > 0. \tag{3.22}$$

THEOREM 3.7. *Let $\beta > -1$, $1 \leq p < \infty$, and $u \in b_\beta^p(S_T)$. Then $\lim_{r \rightarrow 0} u_r = u$ in $b_\beta^p(S_T)$.*

PROOF. Given $\epsilon > 0$, there exists $v \in C_c(\Omega_T)$ such that $\|u - v\|_{L_\beta^p(\Omega_T)} < \epsilon$. We define $v_r : \Omega_T \rightarrow \mathbb{C}$ as

$$v_r(x, t) = \begin{cases} v(x, t+r), & 0 \leq x \leq 2, 0 \leq t \leq T-r, \\ 0, & 0 \leq x \leq 2, T-r \leq t \leq T. \end{cases} \tag{3.23}$$

Clearly $v_r \in C(\overline{\Omega_T})$. We have

$$\|u - u_r\|_{L_\beta^p(S_T)} \leq \|u - v\|_{L_\beta^p(\Omega_T)} + \|v - v_r\|_{L_\beta^p(\Omega_T)} + \|v_r - u_r\|_{L_\beta^p(\Omega_T)}. \tag{3.24}$$

Since v is uniformly continuous on $\overline{\Omega_T}$, it is easy to see that $\lim_{r \rightarrow 0} \|v - v_r\|_{L_\beta^p(\Omega_T)} = 0$.

The uniqueness of the solution of the heat equation on a finite cylinder and (2.12) allow to write

$$u_r(x, t) = \int_0^2 \theta\left(x-y, t - \frac{T}{2} + r\right) u\left(y, \frac{T}{2}\right) dy, \quad \text{for } \frac{T}{2} - r < t < T. \tag{3.25}$$

Hence

$$\begin{aligned} \|v_r - u_r\|_{L_\beta^p(\Omega_T)}^p &\leq \int_0^{T-r} \int_0^2 |v(x, t+r) - u(x, t+r)|^p t^\beta dz \\ &\quad + \left\| u\left(\cdot, \frac{T}{2}\right) \right\|_\infty^p \int_{T-r}^T \int_0^2 \left| \int_0^2 \theta\left(x-y, t - \frac{T}{2} + r\right) dy \right|^p t^\beta dz \\ &= A_r + B_r. \end{aligned} \tag{3.26}$$

By (2.4) we have that $\lim_{r \rightarrow 0} B_r = 2 \|u(\cdot, T/2)\|_\infty^p \lim_{r \rightarrow 0} \int_{T-r}^T t^\beta dt = 0$.

On the other hand, by (3.22) we have

$$\begin{aligned} \int_0^r \int_0^2 |v(x, t+r) - u(x, t+r)|^p t^\beta dz \\ \leq C_p \left\{ \|v\|_\infty^p \int_0^r t^\beta dt + \int_0^r \int_0^2 |u(x, t)|^p t^\beta dz \right\}. \end{aligned} \tag{3.27}$$

If $0 < r \leq t$, then

$$t^\beta \leq \begin{cases} (t+r)^\beta & \text{if } \beta \geq 0, \\ 2^{-\beta}(t+r)^\beta & \text{if } \beta < 0. \end{cases} \tag{3.28}$$

So,

$$\int_r^{T-r} \int_0^2 |v(x, t+r) - u(x, t+r)|^p t^\beta dz \leq C_\beta \|v - u\|_{L^p_\beta(\Omega_T)}^p < C_\beta \epsilon^p. \tag{3.29}$$

From (3.27) and (3.29) it follows that $\limsup_{r \rightarrow 0} A_r \leq C_\beta \epsilon^p$. Hence $\limsup_{r \rightarrow 0} \|u - u_r\|_{L^p_\beta(\Omega_T)} \leq C_\beta \epsilon$. \square

COROLLARY 3.8. *Let $\beta > -1$ and $1 \leq p < \infty$. Then $H(S_T) \cap C(\overline{S_T})$ is dense in $b^p_\beta(S_T)$.*

Now, to compute the reproducing kernel we need to find an orthonormal basis of $b^2(S_T)$, so if we define

$$u_n(x, t) = e^{-\pi^2 n^2 t + \pi n i x}, \tag{3.30}$$

then $u_n \in b^p_\beta(S_T)$ for all $n \in \mathbb{Z}, \beta > -1$. We will show that (u_n) is an orthogonal basis of $b^2(S_T)$.

LEMMA 3.9. *Let $\beta > -1$ and $1 \leq p < \infty$. The linear space generated by $(u_n)_{n \in \mathbb{Z}}$ is dense in $b^p_\beta(S_T)$.*

PROOF. Let $u \in b^p_\beta(S_T)$ and $\epsilon > 0$, by the previous theorem there exists $v \in C(\overline{S_T}) \cap H(S_T)$ such that

$$\|u - v\|_{b^p_\beta(S_T)} < \epsilon. \tag{3.31}$$

Since the set of trigonometric polynomials is dense in $L^p(S^1)$ and $v(\cdot, 0) \in C(S^1)$, there exists a trigonometric polynomial $q(x) = \sum_{|n| \leq N} a_n e^{\pi n i x}$ such that $\|v(\cdot, 0) - q\|_{L^p(S^1)} < \epsilon$.

On the other hand, by (2.12) with $f(x) = e^{\pi n i x}$, we have

$$u_n(x, t) = \int_0^2 \theta(x - y, t) e^{\pi n i y} dy = (K(y, t) * e^{\pi n i y})(x). \tag{3.32}$$

By (3.19) it follows that

$$v(x, t) = \int_0^2 \theta(x - y, t) v(y, 0) dy = (K(\cdot, t) * v(\cdot, 0))(x). \tag{3.33}$$

By (2.14) we obtain

$$\left\| v(\cdot, t) - \sum_{|n| \leq N} a_n u_n(\cdot, t) \right\|_{L^p(S^1)}^p \leq \|v(\cdot, 0) - q\|_{L^p(S^1)}^p, \tag{3.34}$$

Multiplying the previous inequality by t^β and integrating on $(0, T)$, we get

$$\left\| v - \sum_{|n| \leq N} a_n u_n \right\|_{b_\beta^p(S_T)} \leq C_{T,\beta} \|v(\cdot, 0) - q\|_{L^p(S^1)}. \tag{3.35}$$

Therefore, $\|u - \sum_{|n| \leq N} a_n u_n\|_{b_\beta^p(S_T)} < C'_{T,\beta} \epsilon$. Since $\epsilon > 0$ is arbitrary the result follows. \square

Clearly, the sequence $(u_n)_{n \in \mathbb{Z}}$ defined in (3.30) is an orthogonal set in $b^2(S_T)$. It follows that the sequence $(\|u_n\|_{b^2(S_T)}^{-1} u_n)$ is an orthonormal basis of $b^2(S_T)$. We have

$$\|u_n\|_{b^2(S_T)}^2 = \frac{1}{\pi^2 n^2} \gamma(1, 2\pi^2 n^2 T), \tag{3.36}$$

for all $n \in \mathbb{Z}^*$, where γ is defined by

$$\gamma(\alpha, z) = \int_0^z t^{\alpha-1} e^{-t} dt \quad \text{if } \alpha, z > 0. \tag{3.37}$$

Moreover, $\|u_0\|_{b^2(S_T)}^2 = 2T$. By (3.16) it follows that

$$N(z, w) = \frac{1}{2T} + \sum_{n \in \mathbb{Z}^*} \frac{\pi^2 n^2}{\gamma(1, 2\pi^2 n^2 T)} e^{-\pi^2 n^2 (t+\tau) + \pi n i(x-y)}. \tag{3.38}$$

4. The projections P_α . Before proving the continuity of the Bergman projection, we will study certain integral operators P_α with kernel N_α . The operators P_α will turn out to be continuous projections on $b_\beta^p(S_T)$.

DEFINITION 4.1. Given $\alpha > -1$, we define $N_\alpha : \Omega_T \times \Omega_T \rightarrow \mathbb{C}$ as

$$N_\alpha(z, w) = \sum_{m \in \mathbb{Z}} c_{m,\alpha} e^{-\pi^2 m^2 (t+\tau) + \pi m i(x-y)}, \tag{4.1}$$

where $c_{m,\alpha} = 2^\alpha \pi^{2(1+\alpha)} m^{2(1+\alpha)} / \gamma(1 + \alpha, 2\pi^2 m^2 T)$, for all $m \in \mathbb{Z}^*$ and $c_{0,\alpha} = (1 + \alpha) / 2T^{1+\alpha}$.

Since

$$\gamma(1 + \alpha, 2\pi^2 m^2 T) \geq \int_0^{2\pi^2 T} t^\alpha e^{-t} dt = C_\alpha > 0, \tag{4.2}$$

we have that $c_{m,\alpha} \leq C_\alpha m^{2(1+\alpha)}$, for all $m \in \mathbb{Z}^*$.

Using the fact that $e^{-x} \leq C'_\lambda x^{-\lambda}$ for $x > 0$ and $\lambda \geq 0$, we get

$$\left| \sum_{m \in \mathbb{Z}^*} m^k e^{-\pi^2 m^2 (t+\tau) + \pi m i(x-y)} \right| \leq \frac{C_k}{t^{k/2+1}} \sum_{m \in \mathbb{Z}^*} \frac{|m|^k}{(m^2)^{k/2+1}} \leq \frac{C_k}{t_0^{k/2+1}}, \tag{4.3}$$

for $t \geq t_0 > 0, k \geq 0$.

Therefore, the series defining N_α converges absolutely and uniformly on $\Omega' \times \Omega_T$ provided $\Omega' \subset \Omega_T$ is compact, furthermore, the function N_α is bounded on $\Omega' \times \Omega_T$. So $N_\alpha \in C^\infty(\Omega_T \times \Omega_T)$ and $N_\alpha(\cdot, w) \in H(S_T)$ for all $w \in \Omega_T$. Since $c_{m,\alpha} = c_{-m,\alpha}$, the function N_α is real valued and symmetric. If $\alpha = 0$, N_α coincides with the reproducing kernel of $b^2(S_T)$.

DEFINITION 4.2. For $\alpha > -1$, P_α is the integral operator given by

$$P_\alpha u(z) = \int_{\Omega_T} N_\alpha(z, w) u(w) \tau^\alpha dw, \quad z \in \Omega_T. \tag{4.4}$$

This integral is well defined for all $u \in C_c^\infty(\Omega_T)$. If $\alpha = 0$, P_α is the Bergman projection.

It is easy to see that

$$P_\alpha(e^{-\pi^2 n^2 t + \pi n i x}) = e^{-\pi^2 n^2 t + \pi n i x}, \tag{4.5}$$

for all $n \in \mathbb{Z}$. Therefore, P_α is a projection on the linear space generated by $\{e^{-\pi^2 n^2 t + \pi n i x}\}$.

We want to show the continuity of P_α on $L^p_\beta(\Omega_T)$. In order to do so, we analyze the following operator:

$$T_\alpha u(z) = \int_{\Omega_T} \Theta_\alpha(z, w) u(w) \tau^\alpha dw, \quad z \in \Omega_T, \tag{4.6}$$

where

$$\Theta_\alpha(z, w) = \theta_\alpha(x - y, t + \tau) = \frac{1}{2} \pi^{2(1+\alpha)} \sum_{m \in \mathbb{Z}} m^{2(1+\alpha)} e^{-\pi^2 m^2 (t+\tau) + \pi m i (x-y)}. \tag{4.7}$$

The series defining Θ_α has the same properties of convergence as N_α .

REMARK 4.3. If $\alpha \in \mathbb{N}$, then $\theta_\alpha(x, t) = (-1)^{1+\alpha} (\partial^{1+\alpha} / \partial t^{1+\alpha}) \theta(x, t)$.

Let $K_\alpha(x, t)$ be the function defined as

$$\begin{aligned} K_\alpha(x, t) &= \frac{1}{2} F^{-1} \left(\pi^{2(1+\alpha)} \zeta^{2(1+\alpha)} e^{-\pi^2 \zeta^2 t} \right) \left(\frac{x}{2} \right) \\ &= \frac{1}{\sqrt{\pi t}^{1+\alpha}} K(x, t) \int_{-\infty}^{\infty} \left(\sigma + i \frac{x}{2\sqrt{t}} \right)^{2(1+\alpha)} e^{-\sigma^2} d\sigma, \end{aligned} \tag{4.8}$$

where F^{-1} is the inverse Fourier transform with respect to the variable ζ .

We have the following estimate:

$$\begin{aligned} |K_\alpha(x, t)| &\leq \frac{C_\alpha}{t^{1+\alpha}} K(x, t) \int_{-\infty}^{\infty} \left(\sigma^{2(1+\alpha)} + \frac{x^{2(1+\alpha)}}{t^{1+\alpha}} \right) e^{-\sigma^2} d\sigma \\ &\leq \frac{C_\alpha}{t^{1+\alpha}} K(x, t) \left[1 + \frac{x^{2(1+\alpha)}}{t^{1+\alpha}} \right]. \end{aligned} \tag{4.9}$$

Since $x^\lambda e^{-x} \leq C_\lambda e^{-x/2}$ for $x, \lambda > 0$, we have

$$\begin{aligned} \frac{x^{2(1+\alpha)}}{(4t)^{1+\alpha}} K(x, t) &= \frac{1}{\sqrt{4\pi t}} \left(\frac{x^{2(1+\alpha)}}{(4t)^{1+\alpha}} e^{-x^2/4t} \right) \\ &\leq \frac{C_\alpha}{\sqrt{4\pi t}} e^{-x^2/8t} = C_\alpha K(x, 2t). \end{aligned} \tag{4.10}$$

Therefore,

$$|K_\alpha(x, t)| \leq \frac{C_\alpha}{t^{1+\alpha}} \{K(x, t) + K(x, 2t)\}. \tag{4.11}$$

LEMMA 4.4. For $\alpha > -1$, $K_\alpha(x, t) \in C(\mathbb{R}_+^2)$.

PROOF. We write $K_\alpha(x, t) = (1/\sqrt{\pi t}^{1+\alpha})K(x, t)\psi_\alpha(x, t)$, where

$$\psi_\alpha(x, t) = \int_{-\infty}^{\infty} \left(\sigma + i \frac{x}{2\sqrt{t}} \right)^{2(1+\alpha)} e^{-\sigma^2} d\sigma. \tag{4.12}$$

The function $f(z) = z^{2(1+\alpha)}$ is analytic on $\text{Im } z > 0$. Also

$$\begin{aligned} \int_{-\infty}^{\infty} \left| \left(\sigma + i \frac{x}{2\sqrt{t}} \right)^{2(1+\alpha)} \right| e^{-\sigma^2} d\sigma \\ \leq \int_{-\infty}^{\infty} \left(|\sigma| + \frac{|x|}{2\sqrt{t}} \right)^{2(1+\alpha)} e^{-\sigma^2} d\sigma < \infty, \end{aligned} \tag{4.13}$$

for all $(x, t) \in \mathbb{R}_+^2$. From the dominated convergence theorem it follows that $\psi_\alpha(x, t) \in C(\mathbb{R}_+^2)$. □

REMARK 4.5. If $\alpha \in \mathbb{N}$, then $K_\alpha(x, t) = (-1)^{1+\alpha}(\partial^{1+\alpha}/\partial t^{1+\alpha})K(x, t)$.

Now we get an alternate expression for the function θ_α in terms of the function K_α .

PROPOSITION 4.6. For $\alpha > -1$, $\theta_\alpha(x, t) = \sum_{m \in \mathbb{Z}} K_\alpha(x + 2m, t)$.

PROOF. From (4.11) we have

$$\left| \sum_{m \in \mathbb{Z}} K_\alpha(x + 2m, t) \right| \leq \frac{C_\alpha}{t^{1+\alpha}} \{\theta(x, t) + \theta(x, 2t)\}. \tag{4.14}$$

Hence, the series converges uniformly on compact subsets of \mathbb{R}_+^2 and therefore it is continuous. Since the series is 2-periodic in x , it admits a representation as a Fourier series,

$$\sum_{m \in \mathbb{Z}} K_\alpha(x + 2m, t) = \sum_{m \in \mathbb{Z}} a_m(t) e^{\pi m i x}, \tag{4.15}$$

where convergence is in $L^2(S^1)$. Moreover,

$$\int_0^2 \sum_{m \in \mathbb{Z}} |K_\alpha(x + 2m, t)| dx \leq \frac{C_\alpha}{t^{1+\alpha}} \int_{-\infty}^\infty (K(x, t) + K(x, 2t)) dx = \frac{C_\alpha}{t^{1+\alpha}}. \tag{4.16}$$

By the dominated convergence theorem we have

$$\begin{aligned} a_m(t) &= \frac{1}{2} \int_0^2 \left[\sum_{n \in \mathbb{Z}} K_\alpha(x + 2n, t) \right] e^{-\pi m i x} dx \\ &= \frac{1}{2} \sum_{n \in \mathbb{Z}} \int_{2n}^{2n+2} K_\alpha(x, t) e^{-\pi m i x} dx \\ &= \frac{1}{2} \int_{-\infty}^\infty K_\alpha(x, t) e^{-\pi m i x} dx \\ &= F(K_\alpha(2x, t))(m) \\ &= \frac{1}{2} \pi^{2(1+\alpha)} m^{2(1+\alpha)} e^{-\pi^2 m^2 t}. \end{aligned} \tag{4.17}$$

□

The following result is the key to prove the boundedness of the operator P_α . The proof is based on the Schur’s test.

THEOREM 4.7. *Let $\alpha, \beta > -1$. If $p > \max(1, (1 + \beta)/(1 + \alpha))$, the operator $T_\alpha : L^p_\beta(\Omega_T) \rightarrow b^p_\beta(S_T)$ given by*

$$T_\alpha u(z) = \int_{\Omega_T} \Theta_\alpha(z, w) u(w) \tau^\alpha dw, \quad z \in \Omega_T, \tag{4.18}$$

is bounded.

PROOF. Since $\Theta_\alpha(\cdot, w) \in H(S_T)$, it follows that $T_\alpha u \in H(S_T)$ for $u \in C^\infty_c(\Omega_T)$. By Lemma 2.4, it is enough to prove that for each p there is a positive measurable function $h(x, t)$ such that

$$\begin{aligned} \int_{\Omega_T} |\theta_\alpha(x - y, t + \tau)| h(y, \tau)^{p'} \tau^\alpha dy d\tau &\leq C_\alpha h(x, t)^{p'}, \quad (x, t) \in \Omega_T, \\ \int_{\Omega_T} |\theta_\alpha(x - y, t + \tau)| h(x, t)^p \tau^\alpha t^\beta dx dt &\leq C_\alpha h(y, \tau)^p t^\beta, \quad (y, \tau) \in \Omega_T. \end{aligned} \tag{4.19}$$

By Proposition 4.6 and (4.11) we have that

$$|\theta_\alpha(x, t)| \leq \frac{C_\alpha}{t^{1+\alpha}} (\theta(x, t) + \theta(x, 2t)). \tag{4.20}$$

Using (2.11) we get for $0 < x, y < 2$ that

$$\begin{aligned} |\theta_\alpha(x - y, t + \tau)| &\leq \frac{C_\alpha}{(t + \tau)^{1+\alpha}} \{K(x - y + 2, t + \tau) + K(x - y, t + \tau) \\ &\quad + K(x - y - 2, t + \tau) + K(x - y + 2, 2(t + \tau)) \\ &\quad + K(x - y, 2(t + \tau)) + K(x - y - 2, 2(t + \tau))\}. \end{aligned} \tag{4.21}$$

Let $\delta > 0$ be such that $(\beta - \alpha)/p < \delta < \min((1 + \alpha)/p', (1 + \beta)/p)$. Set $h(x, t) = t^{-\delta}$. From [Lemma 2.5](#) we have

$$\begin{aligned} & \int_0^T \frac{\tau^{-\delta p' + \alpha}}{(t + \tau)^{1 + \alpha}} \int_0^2 K(x - y \pm 2, \lambda(t + \tau)) dy d\tau \\ & \leq \int_0^T \frac{\tau^{-\delta p' + \alpha}}{(t + \tau)^{1 + \alpha}} \int_{-\infty}^{\infty} K(y, \lambda(t + \tau)) dy d\tau \\ & = \int_0^T \frac{\tau^{-\delta p' + \alpha}}{(t + \tau)^{1 + \alpha}} d\tau \\ & \leq C_\alpha t^{-\delta p'}. \end{aligned} \tag{4.22}$$

From the estimate in [\(4.21\)](#), together with the above calculations and letting $\lambda = 1, 2$ we conclude that

$$\int_{\Omega_T} |\theta_\alpha(x - y, t + \tau)| \tau^{-\delta p'} \tau^\alpha dy d\tau \leq C_\alpha t^{-\delta p'}. \tag{4.23}$$

The proof of the second inequality is similar. □

The theory of Fourier multipliers is another useful tool that it will help to define an isomorphism \mathcal{M}_α connecting the operators T_α and P_α .

DEFINITION 4.8. Given $1 \leq p \leq \infty$, a bounded sequence (μ_n) is a multiplier on $L^p(S^1)$ if there is $c > 0$ such that

$$\left\| \sum \mu_n \hat{f}(n) e^{inx} \right\|_{L^p(S^1)} \leq c \|f\|_{L^p(S^1)} \tag{4.24}$$

for all trigonometric polynomials f .

The following result about multipliers is due to Hirschman [\[7\]](#).

THEOREM 4.9. Let (μ_n) be a bounded sequence such that $|\mu_n| = O(|n|^{-\epsilon})$, $0 < \epsilon < 1$. Then for all p satisfying $(1 - \epsilon)/2 < 1/p < (1 + \epsilon)/2$, (μ_n) is a multiplier on $L^p(S^1)$.

LEMMA 4.10. Let (λ_n) be a bounded sequence such that $\lambda_n = C_1 + O(|n|^{-\epsilon})$ for some $C_1 \neq 0$ and $0 < C_2 \leq |\lambda_n|$ for $n \in \mathbb{Z}$ and all $\epsilon > 0$. Then (λ_n) induces an isomorphism \mathcal{M} of $b_p^p(S_T)$ onto itself for all $p > 1$.

PROOF. By [Lemma 3.9](#) it is enough to define the operator \mathcal{M} on the elements $e^{-\pi^2 n^2 t + \pi n i x}$, $n \in \mathbb{Z}$:

$$\mathcal{M}(e^{-\pi^2 n^2 t + \pi n i x}) = \lambda_n e^{-\pi^2 n^2 t + \pi n i x}. \tag{4.25}$$

We note that \mathcal{M} can be written as $\mathcal{M} = C_1 I + \mathcal{M}'$, where I is the identity operator and \mathcal{M}' is a multiplier operator by [Theorem 4.9](#). Thus, there exists $C > 0$ such that $\|\mathcal{M}u\|_{L^p(S^1)} \leq C \|u\|_{L^p(S^1)}$ for all trigonometric polynomials u .

By hypothesis (λ_n^{-1}) is a bounded sequence and $\lambda_n^{-1} = C_1^{-1} + O(|n|^{-\epsilon})$, as before we can see that there exists a constant $c > 0$ such that $c\|\mathcal{M}^{-1}u\|_{L^p_t(S^1)} \leq \|u\|_{L^p(S^1)}$ for all trigonometric polynomials u . So,

$$c \left\| \sum_{|n| \leq N} a_n e^{-\pi^2 n^2 t + \pi n i x} \right\|_{L^p(S^1)}^p \leq \left\| \sum_{|n| \leq N} \lambda_n a_n e^{-\pi^2 n^2 t + \pi n i x} \right\|_{L^p(S^1)}^p \leq C \left\| \sum_{|n| \leq N} a_n e^{-\pi^2 n^2 t + \pi n i x} \right\|_{L^p(S^1)}^p. \tag{4.26}$$

Multiplying the inequality by t^β and integrating on $(0, T)$, we have

$$c \left\| \sum_{|n| \leq N} a_n e^{-\pi^2 n^2 t + \pi n i x} \right\|_{b_\beta^p(S_T)} \leq \left\| \sum_{|n| \leq N} \lambda_n a_n e^{-\pi^2 n^2 t + \pi n i x} \right\|_{b_\beta^p(S_T)} \leq C \left\| \sum_{|n| \leq N} a_n e^{-\pi^2 n^2 t + \pi n i x} \right\|_{b_\beta^p(S_T)}. \tag{4.27}$$

□

5. Proof of the main theorems. By the previous work, it is easy to prove that the operator P_α is bounded on $L_\beta^p(\Omega_T)$.

PROOF OF THEOREM 1.1. Since $N_\alpha(\cdot, w) \in H(S_T)$, it follows that $P_\alpha u \in H(S_T)$, for $u \in C_c^\infty(\Omega_T)$.

Let $\lambda_{n,\alpha} = 2^{-\alpha-1} \gamma(1 + \alpha, 2\pi^2 n^2 T)$ for every $n \in \mathbb{Z}^*$ and $\lambda_{0,\alpha} = 2T^{(1+\alpha)} / (1 + \alpha)$. Then

$$0 < c_\alpha = 2^{-\alpha-1} \int_0^{2\pi^2 T} t^\alpha e^{-t} dt \leq \lambda_{n,\alpha} \leq 2^{-\alpha-1} \int_0^\infty t^\alpha e^{-t} dt = C_\alpha, \tag{5.1}$$

for all $n \in \mathbb{Z}^*$. Thus, $c_\alpha \leq \lambda_{n,\alpha} \leq C_\alpha$, for all $n \in \mathbb{Z}$.

Using the fact that $e^{-t} \leq C'_\sigma t^{-\sigma}$ for $t, \sigma > 0$, we have that

$$\int_{2\pi^2 n^2 T}^\infty t^\alpha e^{-t} dt \leq C'_\sigma \int_{2\pi^2 n^2 T}^\infty t^{\alpha-\sigma} dt = \frac{C_{\sigma,T}}{\sigma - \alpha - 1} n^{2(\alpha-\sigma+1)} \tag{5.2}$$

provided $\alpha - \sigma < -1$. Letting $\sigma = \alpha + 1 + \epsilon/2$ we have

$$\lambda_{n,\alpha} = C_\alpha + O(|n|^{-\epsilon}), \quad \forall \epsilon > 0. \tag{5.3}$$

Let \mathcal{M}_α be the isomorphism induced by the sequence $(\lambda_{n,\alpha})$ on $b_\beta^p(S_T)$ (see [Lemma 4.10](#)). That is,

$$\mathcal{M}_\alpha \left(\sum a_n e^{-\pi^2 n^2 t + \pi n i x} \right) = \sum \lambda_{n,\alpha} a_n e^{-\pi^2 n^2 t + \pi n i x}. \tag{5.4}$$

In particular, we have

$$\mathcal{M}_\alpha N_\alpha(z, w) = 1 + \theta_\alpha(x - y, t + \tau), \tag{5.5}$$

where \mathcal{M}_α is acting on the variable z .

Since the series defining $N_\alpha(z, w)$ and $\Theta_\alpha(z, w)$ converges uniformly on the set $\{z\} \times \Omega_T$ we have

$$(\mathcal{M}_\alpha \circ P_\alpha)u(z) = (\mathbf{1}_\alpha + T_\alpha)u(z), \quad \forall u \in C_c^\infty(\Omega_T), \tag{5.6}$$

where $\mathbf{1}_\alpha u(z) = \int_{\Omega_T} u(w) \tau^\alpha dw$ is a bounded operator on $L_\beta^p(\Omega_T)$ if $p > \max(1, (1 + \beta)/(1 + \alpha))$. So, P_α is continuous on $L_\beta^p(\Omega_T)$. The result follows from (4.5) and Lemma 3.9. \square

COROLLARY 5.1. *The Bergman projection $P : L^p(\Omega_T) \rightarrow b^p(S_T)$ is continuous for all $p > 1$.*

5.1. The dual space of $b_\beta^p(S_T)$. This section is devoted to the study of the dual space of $b_\beta^p(S_T)$. For $\beta > -1, p > 1$, we denote by $\langle \cdot, \cdot \rangle_\beta$ the weighted duality between $L_\beta^p(\Omega_T)$, and $L_\beta^{p'}(\Omega_T)$, that is, $\langle u, v \rangle_\beta = \int_{\Omega_T} u(z)v(z)t^\beta dxdt$.

PROOF OF THEOREM 1.2. Let $v \in b_{(\alpha-\beta/p)p'}^{p'}(S_T)$. We define

$$\Phi(u) = \int_{\Omega_T} u(z)v(z)t^\alpha dxdt, \tag{5.7}$$

by Hölder's inequality $\Phi \in (b_\beta^p(S_T))^*$ and $\|\Phi\| \leq \|v\|_{b_{(\alpha-\beta/p)p'}^{p'}(S_T)}$.

Conversely, let $\Phi \in (b_\beta^p(S_T))^*$. Theorem 1.1 implies that $\Phi_\alpha = \Phi \circ P_\alpha \in (L_\beta^p(\Omega_T))^*$. Therefore, there is $v_1 \in L_\beta^{p'}(\Omega_T)$ such that

$$\Phi_\alpha(u) = \langle u, v_1 \rangle_\beta = \int_{\Omega_T} u(z)v_1(z)t^\beta dxdt, \tag{5.8}$$

for all $u \in L_\beta^p(\Omega_T)$.

Since P_α is a projection onto $b_\beta^p(S_T)$ then $\Phi(u) = \Phi_\alpha(u)$ for all $u \in b_\beta^p(\Omega_T)$. Also,

$$\Phi_\alpha(u) = \Phi_\alpha(P_\alpha u) = \int_{\Omega_T} \left(\int_{\Omega_T} N_\alpha(z, w)u(w)\tau^\alpha dyd\tau \right) v_1(z)t^\beta dxdt, \tag{5.9}$$

for all $u \in L_\beta^p(\Omega_T)$.

Consider $u \in C_c^\infty(\Omega_T)$, then

$$\begin{aligned} \Phi_\alpha(u) &= \int_{\Omega_T} \left(\int_{\Omega_T} N_\alpha(z, w)v_1(z)t^\beta dxdt \right) u(w)\tau^\alpha dyd\tau \\ &= \int_{\Omega_T} v(w)u(w)\tau^\alpha dyd\tau. \end{aligned} \tag{5.10}$$

It was proved in Section 4 that $N_\alpha(z, w)$ and all of its partial derivatives are bounded on $\Omega_T \times \Omega'$ with $\Omega' \subset \Omega_T$ compact. Furthermore, $N_\alpha(z, \cdot) \in H(S_T)$. Since $v_1 \in L_\beta^{p'}(\Omega_T) \subset L_\beta^1(\Omega_T)$ then v is a well-defined function satisfying the heat equation.

Since $C_c^\infty(\Omega_T)$ is dense in $L_\beta^p(\Omega_T)$ and $\Phi_\alpha \in (L_\beta^p(\Omega_T))^*$ then $v(w)\tau^{\alpha-\beta} \in L_\beta^{p'}(\Omega_T)$, which implies that $v \in L_{(\alpha-\beta/p)p'}^{p'}(\Omega_T)$. Hence $v \in b_{(\alpha-\beta/p)p'}^{p'}(S_T)$ and it represents Φ .

Finally, we need to prove that the correspondence $v \rightarrow \Phi$ is injective. Since $p' > (1/(1+\alpha))(1+(\alpha-\beta/p)p')$, Theorem 1.1 implies that P_α is a continuous projection from $L_{(\alpha-\beta/p)p'}^{p'}(\Omega_T)$ onto $b_{(\alpha-\beta/p)p'}^{p'}(S_T)$.

Assume that $\Phi = 0$ is represented by $v \in b_{(\alpha-\beta/p)p'}^{p'}(S_T)$. Let $u \in C_c^\infty(\Omega_T)$. By Fubini's theorem we have

$$\begin{aligned} \int_{\Omega_T} u(z)v(z)t^\alpha dxdt &= \int_{\Omega_T} u(z)(P_\alpha v)(z)t^\alpha dxdt \\ &= \int_{\Omega_T} v(w) \left(\int_{\Omega_T} N_\alpha(z, w)u(z)t^\alpha dxdt \right) \tau^\alpha dyd\tau \\ &= \int_{\Omega_T} v(w)(P_\alpha u)(w)\tau^\alpha dyd\tau = \Phi(P_\alpha u) = 0. \end{aligned} \tag{5.11}$$

The density of the space $C_c^\infty(\Omega_T)$ implies that $v = 0$. By the open mapping theorem we have that the norms $\|\Phi_\alpha\|$ and $\|v\|_{b_{(\alpha-\beta/p)p'}^{p'}(S_T)}$ are equivalent. \square

COROLLARY 5.2. *If $p > 1$, then $b^p(S_T)^* = b^{p'}(S_T)$ with the usual duality.*

COROLLARY 5.3. *Let $p > 1$. If $p \leq (1+\beta)/(1+\alpha)$, then $P_\alpha : L_\beta^p(\Omega_T) \rightarrow b_\beta^p(S_T)$ is not bounded.*

PROOF. Note that the adjoint operator $P_\alpha^* : b_{-(\beta/p)p'}^{p'}(S_T) \rightarrow L_{-(\beta/p)p'}^{p'}(\Omega_T)$ under the usual integral pairing is given by

$$P_\alpha^* u(z) = t^\alpha \int_{\Omega_T} N_\alpha(z, w)u(w)dw. \tag{5.12}$$

Since the function 1 is in $b_{-(\beta/p)p'}^{p'}(S_T)$ and $P_\alpha^* 1 = c_{0,\alpha}t^\alpha$ is not in $L_{-(\beta/p)p'}^{p'}(\Omega_T)$ we have that P_α^* is unbounded. Thus, P_α is not bounded. \square

Finally, we give an application of the continuity of P_α^* . We show that there is a constant $C_n > 0$ such that $\|t^{n/2}(\partial^n u/\partial x^n)\|_{L^p(\Omega_T)} \leq C_n \|u\|_{b^p(S_T)}$, for all $u \in b^p(S_T)$.

PROOF OF THEOREM 1.3. If $u \in b^p(S_T)$, then

$$u(z) = \int_{\Omega_T} N(z, w)u(w)dw. \tag{5.13}$$

Differentiating under the integral sign leads to

$$\frac{\partial^n \mathbf{u}}{\partial x^n}(z) = \int_{\Omega_T} \frac{\partial^n N}{\partial x^n}(z, w) \mathbf{u}(w) dw. \tag{5.14}$$

We have

$$\frac{\partial^n N}{\partial x^n}(z, w) = \sum_{m \in \mathbb{Z}^*} \frac{i^n \pi^{2+n} m^{2+n}}{\gamma(1, 2\pi^2 m^2 T)} e^{-\pi^2 m^2(t+\tau) + \pi m i(x-y)}. \tag{5.15}$$

Let D_n be the operator defined by

$$D_n \mathbf{u}(z) = \frac{\partial^n \mathbf{u}}{\partial x^n}(z) = \int_{\Omega_T} \frac{\partial^n N}{\partial x^n}(z, w) \mathbf{u}(w) dw. \tag{5.16}$$

On the other hand, by letting $\alpha = n/2$ in (4.1) we define the operator T_n as follows:

$$T_n \mathbf{u}(z) = \int_{\Omega_T} \left(-\frac{1+n/2}{2T^{1+n/2}} + N_{n/2}(z, w) \right) \mathbf{u}(w) dw. \tag{5.17}$$

Let $\lambda_{m,n} = 2^{n/2} \gamma(1, 2\pi^2 m^2 T) / i^n \gamma(1+n/2, 2\pi^2 m^2 T)$ for every $m \in \mathbb{Z}^*$ and $\lambda_{0,n} = (1+n/2)/2T^{1+n/2}$. As in (5.3) we have

$$\lambda_{m,n} = i^n C_n + i^n O(|m|^{-\epsilon}), \tag{5.18}$$

for all $\epsilon > 0$.

Let \mathcal{M}_n be the isomorphism induced by the sequence $(\lambda_{m,n})$ (see Lemma 4.10). That is,

$$\mathcal{M}_n \left(\sum a_m e^{-\pi^2 m^2 t + \pi m i x} \right) = \sum \lambda_{m,n} a_m e^{-\pi^2 m^2 t + \pi m i x}. \tag{5.19}$$

It is easy to see that

$$(\mathcal{M}_n \circ D_n)(e^{-\pi^2 m^2 t + \pi m i x}) = T_n(e^{-\pi^2 m^2 t + \pi m i x}), \tag{5.20}$$

for all $m \in \mathbb{Z}$.

By the inequality in (4.26), we have that there is a constant $C_n > 0$ such that

$$\|D_n \mathbf{u}\|_{L^p(S^1)}^p = \|\mathcal{M}_n^{-1}(T_n \mathbf{u})\|_{L^p(S^1)}^p \leq C_n \|T_n \mathbf{u}\|_{L^p(S^1)}^p, \tag{5.21}$$

for all $\mathbf{u} \in b^p(S_T)$.

By multiplying this inequality by $t^{np/2}$, integrating on $(0, T)$, and using Tonelli's theorem we have

$$\|t^{n/2} D_n \mathbf{u}\|_{L^p(\Omega_T)}^p \leq C_n \|t^{n/2} T_n \mathbf{u}\|_{L^p(\Omega_T)}^p. \tag{5.22}$$

From (5.12), it follows that

$$P_{n/2}^* \mathbf{u}(z) = t^{n/2} (T_n \mathbf{u})(z) + \frac{1+n/2}{2T^{1+n/2}} t^{n/2} \int_{\Omega_T} \mathbf{u}(w) dw. \quad (5.23)$$

By [Theorem 1.1](#), $P_{n/2} : L^p(\Omega_T) \rightarrow b^p(S_T)$ is a bounded projection, for all $p > 1$. Hence $P_{n/2}^* : b^p(S_T) \rightarrow L^p(\Omega_T)$ is a bounded operator, for all $p > 1$. It follows that

$$\begin{aligned} \|t^{n/2} D_n \mathbf{u}\|_{L^p(\Omega_T)} &\leq C_n \left(\|P_{n/2}^* \mathbf{u}\|_{L^p(\Omega_T)} + \|\mathbf{u}\|_{b^1(S_T)} \|t^{n/2}\|_{L^p(\Omega_T)} \right) \\ &\leq C_n \|\mathbf{u}\|_{b^p(S_T)}. \end{aligned} \quad (5.24)$$

The proof of the other inequality is similar. \square

ACKNOWLEDGMENTS. This work is part of my doctoral thesis at the Universidad Nacional Autónoma de México (UNAM). The thesis was supported by the Consejo Nacional de Ciencia y Tecnología (CONACyT) and the Dirección General de Estudios de Posgrado (DGEP, UNAM). I would like to thank my thesis advisor Salvador Pérez-Esteva. I am grateful to the referees for helpful suggestions.

REFERENCES

- [1] S. Axler, *Bergman spaces and their operators*, Surveys of Some Recent Results in Operator Theory, Vol. I (J. B. Conway and B. B. Morrel, eds.), Pitman Res. Notes Math. Ser., vol. 171, Longman Sci. Tech., Harlow, 1988, pp. 1–50.
- [2] S. Axler, P. Bourdon, and W. Ramey, *Harmonic Function Theory*, Graduate Texts in Mathematics, vol. 137, Springer-Verlag, New York, 1992.
- [3] S. Bergman, *The Kernel Function and Conformal Mapping*, Mathematical Surveys, no. 5, American Mathematical Society, New York, 1950.
- [4] J. R. Cannon, *The One-Dimensional Heat Equation*, Encyclopedia of Mathematics and Its Applications, vol. 23, Addison-Wesley Publishing, Massachusetts, 1984.
- [5] J. R. Cannon and S. Pérez-Esteva, *On the continuous dependence of the solution of a linear parabolic partial differential equation on the boundary data and the solution at an interior spatial point*, Partial Differential Equations and Applications, Lecture Notes in Pure and Appl. Math., vol. 177, Marcel Dekker, New York, 1996, pp. 57–68.
- [6] M. Guzmán-Partida, *Hardy spaces of conjugate temperatures*, Studia Math. 122 (1997), no. 2, 153–165.
- [7] A. Torchinsky, *Real-Variable Methods in Harmonic Analysis*, Pure and Applied Mathematics, vol. 123, Academic Press, Florida, 1986.
- [8] D. V. Widder, *The Heat Equation*, Academic Press, New York, 1975.
- [9] P. Wojtaszczyk, *Banach Spaces for Analysts*, Cambridge Studies in Advanced Mathematics, vol. 25, Cambridge University Press, Cambridge, 1991.
- [10] K. H. Zhu, *Operator Theory in Function Spaces*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 139, Marcel Dekker, New York, 1990.

Marcos López-García: Instituto de Matemáticas, Universidad Nacional Autónoma de México, Circuito Exterior Ciudad Universitaria, México, D.F. 04510, Mexico

E-mail address: flopez@matem.unam.mx



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

