EXAMPLES OF FRIEDRICHS MODEL OPERATORS WITH A CLUSTER POINT OF EIGENVALUES

SERGUEI I. IAKOVLEV

Received 26 February 2002

A family of selfadjoint operators of the Friedrichs model is considered. These symmetric type operators have one singular point, zero of order m . For every $m > 3/2$, we construct a rank 1 perturbation from the class Lip 1 such that the corresponding operator has a sequence of eigenvalues converging to zero. Thus, near the singular point, there is no singular spectrum finiteness condition in terms of a modulus of continuity of a perturbation for these operators in case of $m > 3/2$.

2000 Mathematics Subject Classification: 47B06, 47B25.

1. Statement of the problem and main result. Consider selfadjoint operators S_m , $m > 0$, given by

$$S_m = \operatorname{sgn} t \cdot |t|^m \cdot + (\cdot, \varphi) \varphi \quad (1.1)$$

on the domain of functions $u(t) \in L_2(\mathbb{R})$ such that $|t|^m u(t) \in L_2(\mathbb{R})$. Here $\varphi \in L_2(\mathbb{R})$ and t is the independent variable. The action of the operator S_m can be written as follows:

$$(S_m u)(t) = \operatorname{sgn} t \cdot |t|^m u(t) + \varphi(t) \int_{\mathbb{R}} u(x) \overline{\varphi(x)} dx. \quad (1.2)$$

The function φ is assumed to satisfy the smoothness condition

$$|\varphi(t+h) - \varphi(t)| \leq \omega(|h|), \quad |h| \leq 1, \quad (1.3)$$

where the function $\omega(t)$ (the modulus of continuity of the function φ) is monotone and satisfies a Dini condition

$$\omega(t) \downarrow 0 \quad \text{as } t \downarrow 0, \quad \int_0^1 \frac{\omega(t)}{t} dt < \infty. \quad (1.4)$$

For the operators S_m , the absolutely continuous spectrum fills the real axis $\mathbb{R}$. The behavior of the singular spectrum of the operators S_m is of interest. Note that we define the singular spectrum as the union of the point spectrum and the singular continuous one. The structure of the spectrum $\sigma_{\text{sing}}(S_1)$ (the singular spectrum of the operator $S_1 = t \cdot + (\cdot, \varphi) \varphi$) has been studied in detail in [1, 2, 3, 4, 5, 6, 7, 8, 9]. It is shown in [1, 7] that for this operator,

there exists an exact condition of the singular spectrum finiteness. Namely, if $\omega(t) = O(\sqrt{t})$ as $t \rightarrow 0^+$, $\sigma_{\text{sing}}(S_1)$ consists of at most a finite number of eigenvalues of finite multiplicity (the singular continuous spectrum is missing). But if $\liminf \omega(t)/\sqrt{t} = +\infty$ as $t \rightarrow 0^+$, then we construct examples showing that a nontrivial singular spectrum appears, in particular, the operator S_1 has accumulation points of eigenvalues. Note that the real appearance of a nontrivial singular spectrum in the Friedrichs model was, for the first time, shown by Pavlov and Petras [8].

Using the simple change of variables $\text{sgn } t \cdot |t|^m = x$, we can show that outside any neighborhood of the origin, the structure of the spectrum $\sigma_{\text{sing}}(S_m)$ is identical with the one of the operator S_1 . This is due to the smoothness of this change of variables outside any neighborhood of the origin, and also to the local character of the main results of [1, 2, 3, 4, 5, 6, 7, 8, 9]. Namely, suppose that conditions (1.3), (1.4), and also some additional conditions on the function φ are fulfilled only in a certain interval $(c, d) \subset \mathbb{R}$, then, the main results of [1, 2, 3, 4, 5, 6, 7, 8, 9], concerning the structure of $\sigma_{\text{sing}}(S_1)$, will remain true in any closed subinterval $\Delta \subset (c, d)$. At the same time, as shown in this paper, for the operator S_m , $m > 3/2$, the behavior of the singular spectrum has quite different character in a neighborhood of the origin. In this case, it turns out that the singular spectrum can appear for any modulus of continuity $\omega(t)$. Hence, near zero, there is no condition of the singular spectrum finiteness in terms of the modulus of continuity of $\varphi(t)$ like for the operator S_1 . Here, we can also use the pointed change of variables but, since, for instance, $(\text{sgn } t \cdot |t|^m)'_{|t=0} = 0$ for $m > 1$, it is not smooth (that is, not a diffeomorphism) near zero. In this sense, zero is a singular point of the operators S_m , $m \neq 1$, and needs a special attention.

We start with a formulation of the main theorem of this paper on the construction of a function $\varphi(t)$ such that the corresponding operator S_m , $m > 3/2$, has a nontrivial singular spectrum near the singular point zero and, in particular, a sequence of eigenvalues converging to the origin. The proof of this theorem will be obtained as a combination of a sequence of lemmas. Observe that the actual modulus of continuity $\widetilde{\omega}(h) := \sup\{|\varphi(t_1) - \varphi(t_2)| : |t_1 - t_2| < h\}$ of the function φ always satisfies the additional constraint of semiadditivity $\widetilde{\omega}(t_1 + t_2) \leq \widetilde{\omega}(t_1) + \widetilde{\omega}(t_2)$ for all $t_1, t_2 \geq 0$.

THEOREM 1.1 (main result). *Let a nonnegative, monotone function $\omega(t)$, $t \geq 0$, be semiadditive such that $\omega(t_1 + t_2) \leq \omega(t_1) + \omega(t_2)$ for all $t_1, t_2 \geq 0$. Then for any $m > 3/2$, a compactly supported function φ satisfying the smoothness condition $|\varphi(t+h) - \varphi(t)| \leq \omega(|h|)$, $h \in \mathbb{R}$, is constructed and such that the corresponding operator $S_m = \text{sgn } t \cdot |t|^m \cdot +(\cdot, \varphi)\varphi$ has a sequence of eigenvalues converging to zero.*

Note that the result of Theorem 1.1 can be formulated in terms of real zeros of some analytic functions. Define in the upper half plane an analytic function

$M_m(z)$ in the following way:

$$M_m(z) = 1 + \int_{-\infty}^{+\infty} \frac{|\varphi^2(t)|}{\operatorname{sgn} t \cdot |t|^m - z} dt, \quad \operatorname{Im} z > 0. \quad (1.5)$$

It is easily shown that under conditions (1.3) and (1.4), the function $M_m(z)$ is continuously extended up to the real axis on the intervals $(-\infty; 0)$ and $(0; +\infty)$. We define, for $\lambda \in \mathbb{R} \setminus \{0\}$, the value $M_m(\lambda) := M_m(\lambda + i0)$ and the roots set $N := \{\lambda \in \mathbb{R} \setminus \{0\} : M_m(\lambda) = 0\}$. Then, we have the following inclusion $\sigma_{\text{sing}}(S_m) \subseteq N \cup \{0\}$. (See, e.g., [10] where similar assertions are proved for the function $1 + \int_{-\infty}^{+\infty} (|\varphi^2(t)|/(t^2 - z))dt$ or [8] for the function $1 + \int_{-\infty}^{+\infty} (|\varphi^2(t)|/(t - z))dt$.) Further, the exact condition $\omega(t) = O(\sqrt{t})$, as $t \rightarrow 0^+$, appears to guarantee that outside any neighborhood of the origin, there is at most a finite number of zeros of the function $M_m(\lambda)$. At the same time, Theorem 1.1 means that, for $m > 3/2$, the function $M_m(\lambda)$ can have a sequence of zeros converging to the origin for any monotone, nonnegative, and semiadditive function ω satisfying condition (1.4).

2. Construction of the function φ

LEMMA 2.1. *Let the function φ belong to $L_2(\mathbb{R})$. Then a point $\lambda \in \mathbb{R}$ is an eigenvalue of the operator $S_m = \operatorname{sgn} t \cdot |t|^m \cdot +(\cdot, \varphi)\varphi$ if and only if the following conditions hold:*

$$\frac{\varphi(t)}{\operatorname{sgn} t \cdot |t|^m - \lambda} \in L_2(\mathbb{R}), \quad (2.1)$$

$$1 + \int_{-\infty}^{+\infty} \frac{|\varphi^2(t)|}{\operatorname{sgn} t \cdot |t|^m - \lambda} dt = 0. \quad (2.2)$$

PROOF. Solving the eigenvalue problem $S_m v = \lambda v$, we find that $v(t) = C\varphi(t)/(\operatorname{sgn} t \cdot |t|^m - \lambda)$. (Remark that from now on, we denote by C various constants which may be different even in a single chain of inequalities.) This expression will be an eigenfunction if it is from the space $L_2(\mathbb{R})$ and satisfies the equation $S_m v - \lambda v = 0$, that is,

$$(\operatorname{sgn} t \cdot |t|^m - \lambda) \frac{C\varphi(t)}{\operatorname{sgn} t \cdot |t|^m - \lambda} + \varphi(t) \int_{-\infty}^{+\infty} \frac{C|\varphi^2(x)|}{\operatorname{sgn} x \cdot |x|^m - \lambda} dx = 0, \quad (2.3)$$

or

$$\left(1 + \int_{-\infty}^{+\infty} \frac{|\varphi^2(x)|}{\operatorname{sgn} x \cdot |x|^m - \lambda} dx\right) \varphi(t) = 0. \quad (2.4)$$

The proof is complete. $\square$

LEMMA 2.2. *Suppose that a nonnegative, monotone function $\omega(t)$ is semiadditive $\omega(t_1 + t_2) \leq \omega(t_1) + \omega(t_2)$ for all $t_1, t_2 \geq 0$. If $\liminf_{t \rightarrow 0^+} \omega(t)/t = 0$, then $\omega(t) = 0$ for all $t \geq 0$.*

PROOF. According to the semiadditivity, $\omega(nt) \leq n\omega(t)$, $n \in \mathbb{N}$. Therefore, using the notation $[t]$ for the integer part of t , by the monotonicity of $\omega(t)$, we have for all $t, x > 0$,

$$\begin{aligned} \omega(t) &= \omega\left(\left(\frac{t}{x}\right)x\right) \leq \omega\left(\left(\left[\frac{t}{x}\right] + 1\right)x\right) \leq \left(\left[\frac{t}{x}\right] + 1\right)\omega(x) \\ &\leq \left(\frac{t}{x} + 1\right)\omega(x) = (t+x)\frac{\omega(x)}{x}. \end{aligned} \quad (2.5)$$

Thus,

$$\begin{aligned} \omega(t) &\leq \liminf_{x \rightarrow 0^+} (t+x)\frac{\omega(x)}{x} = \lim_{\delta \rightarrow 0^+} \inf_{0 < x < \delta} (t+x)\frac{\omega(x)}{x} \\ &\leq \lim_{\delta \rightarrow 0^+} (t+\delta) \inf_{0 < x < \delta} \frac{\omega(x)}{x} = t \cdot \liminf_{x \rightarrow 0^+} \frac{\omega(x)}{x} = 0. \end{aligned} \quad (2.6)$$

The lemma is proved. $\square$

COROLLARY 2.3. *If additionally $\omega(t) \rightarrow 0^+$ as $t \rightarrow 0^+$, then for any $t_0 > 0$, there exists a constant $C > 0$ such that*

$$Ct \leq \omega(t), \quad t \in [0; t_0]. \quad (2.7)$$

PROOF. It suffices to show that there exists a positive ε less than t_0 such that the inequality $Ct \leq \omega(t)$ is fulfilled with a certain constant $C > 0$ for all $t \in [0; \varepsilon]$. If this is the case, put $\tilde{C} := \min\{C, \omega(\varepsilon)/t_0\}$. Then, the inequality $\tilde{C}t \leq \omega(t)$ already holds for all $t \in [0; t_0]$. Indeed, using the monotonicity of ω , we have for $t \in [\varepsilon; t_0]$

$$\tilde{C}t \leq \frac{\omega(\varepsilon)}{t_0} \cdot t \leq \inf_{x \in [\varepsilon; t_0]} \left(\frac{\omega(x)}{x}\right) \cdot t \leq \frac{\omega(t)}{t} \cdot t = \omega(t). \quad (2.8)$$

We prove that there are such constants ε and C . By assuming the converse, we construct two positive sequences C_k and t_k tending to 0 such that $\omega(t_k) \leq C_k t_k$. Hence, $\lim_{k \rightarrow +\infty} \omega(t_k)/t_k = 0$. Since

$$\liminf_{t \rightarrow 0^+} \frac{\omega(t)}{t} \leq \liminf_{k \rightarrow +\infty} \frac{\omega(t_k)}{t_k} = 0, \quad (2.9)$$

by Lemma 2.2, $\omega = 0$. This completes the proof. $\square$

Remark that the nonnegativity of ω is a consequence of its semiadditivity and monotonicity as $\omega(t+0) \leq \omega(t) + \omega(0)$, and hence $0 \leq \omega(0) \leq \omega(t)$.

Corollary 2.3 shows that it is sufficient to prove **Theorem 1.1** for $\omega(t) = C_\omega t$ with an arbitrary constant $C_\omega > 0$.

Let $\{u_n\}_{n=0}^{+\infty}$ and $\{\varepsilon_n\}_{n=1}^{+\infty}$ be two sequences from the interval $(0; 10^{-1})$ satisfying the condition

$$u_n < \varepsilon_n < \frac{u_{n-1}}{8}, \quad n = 1, 2, \dots \quad (2.10)$$

On the real axis, we define a sequence of functions φ_n as follows:

$$\varphi_n(t) := \begin{cases} \omega\left(t - \frac{u_n}{2}\right), & t \in \left[\frac{1}{2}u_n; \frac{3}{4}u_n\right], \\ \omega(u_n - t), & t \in \left[\frac{3}{4}u_n; u_n\right], \\ 0, & t \notin \left[\frac{1}{2}u_n; u_n\right], \end{cases} \quad (2.11)$$

where $\omega(t) = C_\omega t$.

We will show that for any real-valued Lipschitz function $y(t)$ compactly supported in the interval $(-\infty; -1)$, it is possible to select the sequences u_n and ε_n , and a bounded sequence of nonnegative numbers $\{c^n\}_{n=1}^{+\infty}$ such that the points $\lambda_n := (u_n + \varepsilon_n)^m$ will be eigenvalues of the operator $S_m = \operatorname{sgn} t \cdot |t|^m \cdot +(\cdot, \varphi)\varphi$ with the function

$$\varphi(t) := K \cdot \sum_{k=1}^{+\infty} (c^k)^{1/2} \varphi_k(t) + y(t). \quad (2.12)$$

Here $K > 0$ is a parameter. It will be shown that for all K large enough, the function φ satisfies the required smoothness condition $|\varphi(t+h) - \varphi(t)| \leq \omega(|h|)$, $t, h \in \mathbb{R}$.

LEMMA 2.4. *For the points λ_n to be eigenvalues of the operator S_m defined by (1.1), it is necessary and sufficient that*

$$\int_{-\infty}^{+\infty} \frac{|\varphi^2(t)|}{\operatorname{sgn} t \cdot |t|^m - \lambda_1} dt = -1, \quad (2.13)$$

$$\int_{-\infty}^{+\infty} \frac{|\varphi^2(t)|}{(\operatorname{sgn} t \cdot |t|^m - \lambda_n)(\operatorname{sgn} t \cdot |t|^m - \lambda_{n+1})} dt = 0, \quad n = 1, 2, \dots \quad (2.14)$$

PROOF. We need to verify conditions (2.1) and (2.2) for $\lambda = \lambda_n$, $n = 1, 2, \dots$. As $u_n < u_{n-1}/8$, the supports of the functions φ_n are disjoint. Thus, the function φ is bounded and compactly supported. According to (2.10), $\lambda_n = (u_n + \varepsilon_n)^m < t^m$ for $t \geq u_{n-1}/2$. Therefore, φ vanishes identically in a neighborhood of λ_n . Hence, condition (2.1) is fulfilled for any $\lambda = \lambda_n$, $n = 1, 2, \dots$.

Now putting in (2.2) $\lambda = \lambda_n$, $n = 1, 2, \dots$ we then pass from the obtained system of equations to an equivalent one by subtracting the n th equation from the $(n+1)$ st for $n = 1, 2, \dots$. As a result, we get system (2.13) and (2.14). The lemma is proved. $\square$

Since $\varphi(t) = 0$ for $t > \lambda_1$, it follows that

$$\alpha_m := \int_{-\infty}^{+\infty} \frac{\varphi^2(t)}{\operatorname{sgn} t \cdot |t|^m - \lambda_1} dt < 0. \quad (2.15)$$

Therefore, after solving the homogeneous system (2.14), the first equality (2.13) will be satisfied by replacing the function φ by $\varphi/\sqrt{|\alpha_m|}$.

Substituting expression (2.12) for $\varphi(t)$ in (2.14), we obtain a system of linear equations for the unknowns c^n ,

$$\sum_{k=1}^{n-1} (-d_{nk}c^k) + d_{nn}c^n + \sum_{k=n+1}^{+\infty} (-d_{nk}c^k) = y_n, \quad n = 1, 2, \dots, \quad (2.16)$$

with the coefficients

$$d_{nk} := K^2 \int_{u_k/2}^{u_k} \frac{\varphi_k^2(t)}{|(t^m - \lambda_n)(t^m - \lambda_{n+1})|} dt, \quad (2.17)$$

$$y_n := \int_{-\infty}^{-1} \frac{y^2(t)}{(|t|^m + \lambda_n)(|t|^m + \lambda_{n+1})} dt. \quad (2.18)$$

In the next section we show that the linear system (2.16) has a nonnegative solution in the space l_∞ of bounded sequences.

3. Solution of the linear system

LEMMA 3.1. *The coefficients d_{nk} of the linear system (2.16) satisfy the following inequalities:*

$$d_{nn} \geq K^2 \frac{C_1}{\varepsilon_n^m u_n^{m-3}}, \quad (3.1)$$

$$\sum_{k=1}^{n-1} d_{nk} \leq K^2 \frac{C_2}{u_{n-1}^{2m}}, \quad (3.2)$$

$$\sum_{k=n+1}^{+\infty} d_{nk} \leq K^2 \frac{C_3 u_{n+1}^3}{\varepsilon_n^m \varepsilon_{n+1}^m}, \quad (3.3)$$

with some positive constants C_1 , C_2 , and C_3 .

PROOF. For arbitrary $a, b > 0$ and $m \geq 1$, the following inequality obviously holds:

$$|a - b| \cdot \max\{a^{m-1}, b^{m-1}\} \leq |a^m - b^m| \leq |a - b| \cdot \max\{a^{m-1}, b^{m-1}\} m. \quad (3.4)$$

Thus,

$$\begin{aligned} \frac{1}{|a-b| \cdot \max\{a^{m-1}, b^{m-1}\}m} &\leq \frac{1}{|a^m - b^m|} \\ &\leq \frac{1}{|a-b| \cdot \max\{a^{m-1}, b^{m-1}\}}. \end{aligned} \quad (3.5)$$

Using inequalities (3.5) and (2.10) with elementary estimates and a simple change of variables $x := u_n - t$, we get

$$\begin{aligned} d_{nn} &\geq K^2 \int_{3u_n/4}^{u_n} \frac{\omega^2(u_n - t)dt}{((u_n + \varepsilon_n)^m - t^m) \cdot (t^m - (u_{n+1} + \varepsilon_{n+1})^m)} \\ &\geq K^2 \int_{3u_n/4}^{u_n} \omega^2(u_n - t)dt \\ &\quad \cdot \frac{1}{((u_n + \varepsilon_n)^m - (3u_n/4)^m) \cdot (u_n^m - (u_{n+1} + \varepsilon_{n+1})^m)} \\ &\geq K^2 \int_0^{u_n/4} \omega^2(x)dx \\ &\quad \cdot \frac{1}{(m(u_n + \varepsilon_n - (3/4)u_n)(u_n + \varepsilon_n)^{m-1}) \cdot (m(u_n - u_{n+1} - \varepsilon_{n+1})u_n^{m-1})} \\ &\geq K^2 \frac{C_\omega^2}{3} \left(\frac{u_n}{4}\right)^3 \cdot \frac{1}{(m(u_n/4 + \varepsilon_n)(u_n + \varepsilon_n)^{m-1}) \cdot (mu_n^m)} \\ &\geq K^2 C \frac{u_n^3}{(u_n + \varepsilon_n)^m \cdot u_n^m} \geq K^2 \frac{C_1}{\varepsilon_n^m \cdot u_n^{m-3}}. \end{aligned} \quad (3.6)$$

Inequality (3.2) is established analogously,

$$\begin{aligned} \sum_{k=1}^{n-1} d_{nk} &= \sum_{k=1}^{n-1} K^2 \int_{u_k/2}^{u_k} \frac{\varphi_k^2(t)dt}{(t^m - (u_n + \varepsilon_n)^m) \cdot (t^m - (u_{n+1} + \varepsilon_{n+1})^m)} \\ &\leq K^2 \sum_{k=1}^{n-1} \int_{u_k/2}^{u_k} \frac{\varphi_k^2(t)dt}{((u_{n-1}/2)^m - (u_n + \varepsilon_n)^m) \cdot ((u_{n-1}/2)^m - (u_{n+1} + \varepsilon_{n+1})^m)} \\ &\leq K^2 \int_0^{u_1} \left(\sum_{k=1}^{+\infty} \varphi_k^2(t) \right) dt \frac{1}{((u_{n-1}/2 - u_n - \varepsilon_n)(u_{n-1}/2)^{m-1})} \\ &\quad \times \frac{1}{((u_{n-1}/2 - u_{n+1} - \varepsilon_{n+1})(u_{n-1}/2)^{m-1})} \\ &\leq K^2 C \frac{1}{(u_{n-1}/2 - u_{n-1}/8 - u_{n-1}/8)^2 \cdot (u_{n-1}/2)^{2(m-1)}} \leq K^2 \frac{C_2}{u_{n-1}^{2m}}. \end{aligned} \quad (3.7)$$

We took into account that the functions $\varphi_k(t)$ are uniformly bounded and have disjoint supports, therefore $\int_0^{u_1} (\sum_{k=1}^{+\infty} \varphi_k^2(t)) dt \leq C$. Finally, we have

$$\begin{aligned}
 & \sum_{k=n+1}^{+\infty} d_{nk} \\
 &= K^2 \sum_{k=n+1}^{+\infty} \int_{u_{k/2}}^{u_k} \frac{\varphi_k^2(t) dt}{((u_n + \varepsilon_n)^m - t^m) \cdot ((u_{n+1} + \varepsilon_{n+1})^m - t^m)} \\
 &\leq K^2 \sum_{k=n+1}^{+\infty} \int_{u_{k/2}}^{u_k} \frac{\varphi_k^2(t) dt}{((u_n + \varepsilon_n)^m - u_{n+1}^m) \cdot ((u_{n+1} + \varepsilon_{n+1})^m - u_{n+1}^m)} \\
 &\leq K^2 \sum_{k=n+1}^{+\infty} \int_{u_{k/2}}^{u_k} \frac{\omega^2(u_k - t) dt}{((u_n + \varepsilon_n - u_{n+1})(u_n + \varepsilon_n)^{m-1}) \cdot (\varepsilon_{n+1}(u_{n+1} + \varepsilon_{n+1})^{m-1})} \\
 &\leq K^2 \sum_{k=n+1}^{+\infty} \int_{u_{k/2}}^{u_k} \frac{\omega^2(u_{n+1} - t) dt}{((u_n + \varepsilon_n - u_{n+1}/8)(u_n + \varepsilon_n)^{m-1}) \cdot \varepsilon_{n+1}(u_{n+1} + \varepsilon_{n+1})^{m-1}} \\
 &\leq K^2 \int_0^{u_{n+1}} \omega^2(u_{n+1} - t) dt \frac{C}{(u_n + \varepsilon_n)^m \cdot \varepsilon_{n+1}(u_{n+1} + \varepsilon_{n+1})^{m-1}} \\
 &\leq K^2 u_{n+1}^3 \frac{C_3}{\varepsilon_n^m \cdot \varepsilon_{n+1}^m}.
 \end{aligned} \tag{3.8}$$

The lemma is proved. $\square$

We rewrite system (2.16) in matrix form

$$(I + A)\vec{c} = f, \tag{3.9}$$

where the column vectors $\vec{c} = (c^1, c^2, \dots)^T$, $f = (y_1/d_{11}, y_2/d_{22}, \dots)^T$, and the infinite matrix A has the entries $(A)_{nk} = (\delta_{nk} - 1) \cdot d_{nk}/d_{nn}$. Equation (3.9) will be considered in the Banach space l_∞ .

In the sequel, we consider the sequences u_n and ε_n defined as follows:

$$u_n = u_{n-1}^\alpha, \quad \varepsilon_n = u_{n-1}^\beta, \quad n = 1, 2, \dots, \tag{3.10}$$

with some $u_0 \in (0; 10^{-1})$ and $\alpha > \beta > 2$. It is evident that inequality (2.10) is fulfilled.

LEMMA 3.2. *For every $m > 3/2$, the numbers α and β can be found satisfying the inequality $\alpha > \beta > 2$ such that the following estimate:*

$$d_{nn} \geq K^2 C_1 \tag{3.11}$$

holds for all $n = 1, 2, \dots$

PROOF. Substituting (3.10) in (3.1), we get

$$d_{nn} \geq K^2 \frac{C_1}{u_{n-1}^{\beta m + \alpha(m-3)}}. \quad (3.12)$$

As $u_{n-1} \in (0; 1)$ for $m \geq 3$, the assertion of the lemma is true with any $\alpha > \beta > 2$. Now let $m \in (3/2; 3)$. Then, the inequality $\beta m + \alpha(m-3) > 0$ is fulfilled if $\alpha < \beta m / (3-m)$. It must be $\alpha > \beta > 2$, therefore we require that $\beta m / (3-m) > \beta$. As the last inequality holds for any $\beta > 0$, we can take for $m \in (3/2; 3)$ any

$$\beta > 2, \quad \alpha \in \left(\beta; \frac{\beta m}{(3-m)} \right). \quad (3.13)$$

This completes the proof. $\square$

By (2.18), $y_n > 0$, and

$$y_n = \int_{-\infty}^{-1} \frac{|y(t)|^2 dt}{(|t|^m + \lambda_n)(|t|^m + \lambda_{n+1})} \leq \int_{-\infty}^{-1} |y(t)|^2 dt \equiv \|y\|_2^2. \quad (3.14)$$

Hence, for the l_∞ -norm of the vector f equal to $\|f\|_\infty = \sup_n (y_n / d_{nn})$, we obtain the following estimate:

$$\|f\|_\infty \leq \frac{\|y\|_2^2}{K^2 C_1}. \quad (3.15)$$

LEMMA 3.3. *For every $m > 3/2$, the numbers α and β can be found satisfying the inequality $\alpha > \beta > 2$ such that $\|A\| < 1$ for all u_0 small enough.*

PROOF. By the definition,

$$\begin{aligned} \|A\| &= \sup_{\|s\|_\infty=1} \|As\|_\infty \leq \sup_{\|s\|_\infty=1} \sup_n \sum_{k=1}^{+\infty} |A_{nk} s_k| \\ &\leq \sup_n \sum_k |A_{nk}| = \sup_n \sum_{k \neq n} \frac{d_{nk}}{d_{nn}} \\ &= \sup_n \left(\sum_{k=1}^{n-1} \frac{d_{nk}}{d_{nn}} + \sum_{k=n+1}^{+\infty} \frac{d_{nk}}{d_{nn}} \right). \end{aligned} \quad (3.16)$$

Hence, it suffices to require that

$$S_1 := \frac{(\sum_{k=1}^{n-1} d_{nk})}{d_{nn}} < \frac{1}{4}, \quad S_2 := \frac{(\sum_{k=n+1}^{+\infty} d_{nk})}{d_{nn}} < \frac{1}{4}. \quad (3.17)$$

Using (3.1), (3.2), (3.3), and (3.10), we get

$$\begin{aligned} S_1 &\leq \frac{C_2}{C_1} \cdot \frac{\varepsilon_n^m u_n^{m-3}}{u_{n-1}^{2m}} = \frac{C_2}{C_1} \cdot u_{n-1}^{\alpha(m-3)+(\beta-2)m}, \\ S_2 &\leq \frac{C_3}{C_1} \cdot \frac{u_n^{m-3} u_{n+1}^3}{\varepsilon_{n+1}^m} = \frac{C_3}{C_1} \cdot u_n^{3\alpha-m\beta+m-3}. \end{aligned} \quad (3.18)$$

Thus, if there exist $\alpha > \beta > 2$ such that

$$\alpha(m-3) + (\beta-2)m > 0, \quad 3\alpha - m\beta + m - 3 > 0, \quad (3.19)$$

then

$$S_1 \leq \left(\frac{C_2}{C_1}\right) \cdot u_0^{\alpha(m-3)+(\beta-2)m}, \quad S_2 \leq \left(\frac{C_3}{C_1}\right) \cdot u_0^{3\alpha-m\beta+m-3}. \quad (3.20)$$

Therefore, (3.17) will be true for all u_0 small enough.

Let $m \geq 3$. The first inequality in (3.19) holds for all $\alpha > \beta > 2$. The second one is true for $\alpha > (\beta m + 3 - m)/3$. It follows that we can take any $\beta > 2$ and $\alpha > \max\{\beta, (\beta m + 3 - m)/3\} = (\beta m + 3 - m)/3$. Note that this choice of α and β is consistent with the choice made for $m \geq 3$ in Lemma 3.2.

Now let $m \in (3/2; 3)$. Then, (3.19) is equivalent to

$$\alpha < \frac{(\beta m - 2m)}{(3 - m)}, \quad \alpha > \frac{(\beta m + 3 - m)}{3}. \quad (3.21)$$

As it must be $\alpha > \beta$, find β such that

$$\frac{(\beta m + 3 - m)}{3} < \beta < \frac{(\beta m - 2m)}{(3 - m)}. \quad (3.22)$$

The inequality $(\beta m + 3 - m)/3 < \beta$ is fulfilled for all $\beta > 1$. The inequality $\beta < (\beta m - 2m)/(3 - m)$ is equivalent to $\beta > 2m/(2m - 3)$. Hence, for $m \in (3/2; 3)$, we can take any $\beta > \max\{2; 2m/(2m - 3)\} = 2m/(2m - 3)$ and $\alpha \in (\beta; (\beta m - 2m)/(3 - m))$. We also see that α and β satisfy (3.13). This completes the proof. $\square$

By virtue of the inequality $\|A\| < 1$, equation (3.9) has a unique solution in l_∞ ,

$$\tilde{c} = (I + A)^{-1} f. \quad (3.23)$$

It is easily seen that all the components c^n of the vector $\vec{c} = (c^1, c^2, \dots)^T$ are nonnegative. Indeed, $f_n = y_n/d_{nn} > 0$ since, by (2.17) and (2.18), $y_n, d_{nn} > 0$. The Neumann series $\vec{c} = (I + (-A) + (-A)^2 + \dots)f$ includes powers of a matrix with nonnegative entries such that $(-A)_{nk} = (1 - \delta_{nk}) \cdot d_{nk}/d_{nn} \geq 0$ since $d_{nk} \geq 0$. Consequently, the vector $\vec{c}$ is obtained as a result of the action of a matrix with nonnegative entries on a vector whose components are also nonnegative.

4. Smoothness of the function φ . Introducing the convenient notations

$$\begin{aligned} u_n^1 &:= u_n, & u_n^2 &:= \frac{u_n}{2}, & \Delta_n &:= [u_n^2; u_n^1], \\ \Delta_n^2 &:= \left[u_n^2; \frac{(u_n^1 + u_n^2)}{2} \right], & \Delta_n^1 &:= \left[\frac{(u_n^1 + u_n^2)}{2}; u_n^1 \right], \end{aligned} \quad (4.1)$$

we can write (2.11) as follows:

$$\varphi_n(t) := \begin{cases} \omega(t - u_n^2), & t \in \Delta_n^2, \\ \omega(u_n^1 - t), & t \in \Delta_n^1, \\ 0, & t \notin \Delta_n. \end{cases} \quad (4.2)$$

LEMMA 4.1. *Suppose that ω is monotone and semiadditive (and thus non-negative). Then, the function*

$$\psi(t) := \sum_{k=1}^{+\infty} (c^k)^{1/2} \varphi_k(t) \quad (4.3)$$

satisfies the following smoothness condition:

$$|\psi(t+h) - \psi(t)| \leq \sup_k (c^k)^{1/2} \omega(|h|), \quad t, h \in \mathbb{R}. \quad (4.4)$$

Inequality (4.4) remains true if $\{\Delta_n\}_{n=1}^{+\infty}$ is an arbitrary sequence of disjoint finite intervals.

PROOF. If $t_1 \in \Delta_n$ and $t_2 \notin \Delta_n$, then, as the intervals Δ_k are disjoint for different k , for some $i \in \{1, 2\}$, we have

$$\psi(t_1) = (c^n)^{1/2} \cdot \omega(|t_1 - u_n^i|) \leq (c^n)^{1/2} \cdot \omega(|t_1 - t_2|). \quad (4.5)$$

Thus,

$$|\psi(t_2) - \psi(t_1)| \leq \max\{\psi(t_2), \psi(t_1)\} \leq \sup_k (c^k)^{1/2} \cdot \omega(|t_1 - t_2|). \quad (4.6)$$

For $t, \delta > 0$, $\omega(t + \delta) - \omega(t) \leq \omega(\delta)$ according to the semiadditivity of ω . Therefore, for $t_1, t_2 \in \Delta_n^i$, we obtain

$$\begin{aligned} |\psi(t_1) - \psi(t_2)| &= (c^n)^{1/2} \cdot |\omega(|t_1 - u_n^i|) - \omega(|t_2 - u_n^i|)| \\ &\leq (c^n)^{1/2} \cdot \omega(|t_1 - t_2|). \end{aligned} \quad (4.7)$$

Now, let $t_1 \in \Delta_n^i$ and $t_2 \in \Delta_n^j$, $i \neq j$. Assume that $|t_1 - u_n^i| \geq |t_2 - u_n^j|$, then

$$\begin{aligned} |\psi(t_1) - \psi(t_2)| &= (c^n)^{1/2} \cdot (\omega(|t_1 - u_n^i|) - \omega(|t_2 - u_n^j|)) \\ &\leq (c^n)^{1/2} \cdot (\omega(|t_1 - u_n^j|) - \omega(|t_2 - u_n^j|)) \\ &\leq (c^n)^{1/2} \cdot \omega(|t_1 - t_2|). \end{aligned} \quad (4.8)$$

The lemma is proved. $\square$

By (2.12), the function $\varphi(t) = K\psi(t) + \gamma(t)$ where the function γ is assumed to satisfy a Lipschitz condition $|\gamma(t+h) - \gamma(t)| \leq C_\gamma|h|$, $h \in \mathbb{R}$. Since the functions ψ and γ have disjoint supports and $\omega(t) = C_\omega t$, we see that the following inequality:

$$\begin{aligned} |\varphi(t+h) - \varphi(t)| &\leq K \sup_k (c^k)^{1/2} \cdot \omega(|h|) + C_\gamma|h| \\ &\leq \left(K \sup_k (c^k)^{1/2} + \frac{C_\gamma}{C_\omega} \right) \omega(|h|) \end{aligned} \quad (4.9)$$

holds for all $t, h \in \mathbb{R}$.

After finding the solution of system (2.14), we satisfy (2.13) replacing the function φ by $\varphi/\sqrt{|\alpha_m|}$. This replacement corresponds to a passage from the functions ψ and γ to $\psi/\sqrt{|\alpha_m|}$ and $\gamma/\sqrt{|\alpha_m|}$, respectively. Therefore, for the new function φ , the following smoothness condition will be fulfilled:

$$|\varphi(t+h) - \varphi(t)| \leq \left(K \sup_k (c^k)^{1/2} |\alpha_m|^{-1/2} + \frac{C_\gamma |\alpha_m|^{-1/2}}{C_\omega} \right) \omega(|h|), \quad h \in \mathbb{R}. \quad (4.10)$$

LEMMA 4.2. *The constant*

$$M := K \sup_k (c^k)^{1/2} |\alpha_m|^{-1/2} + \frac{C_\gamma |\alpha_m|^{-1/2}}{C_\omega}, \quad (4.11)$$

in the smoothness condition (4.10), satisfies the inequality $M \leq C/K$, and hence, it is less than one for all K large enough.

PROOF. Since

$$\begin{aligned} |\alpha_m| &= \int_{-\infty}^{+\infty} \frac{\varphi^2(t)}{|\operatorname{sgn} t \cdot |t|^m - \lambda_1|} dt \\ &\geq K^2 \int_0^1 \frac{\psi^2(t)}{\lambda_1 - t^m} dt \geq K^2 \int_0^1 \psi^2(t) dt \equiv K^2 \|\psi\|_2^2, \end{aligned} \quad (4.12)$$

it follows that $|\alpha_m|^{-1/2} \leq 1/(K\|\psi\|_2)$. From (3.23), we obtain the inequality

$$\|\tilde{c}\|_\infty \leq (1 - \|A\|)^{-1} \|f\|_\infty. \quad (4.13)$$

In view of relation (3.15), then

$$\|\tilde{c}\|_\infty \leq (1 - \|A\|)^{-1} \frac{\|y\|_2^2}{K^2 C_1}. \quad (4.14)$$

Therefore, for the first summand in (4.11), we get

$$K \sup_k (c^k)^{1/2} |\alpha_m|^{-1/2} \leq K (1 - \|A\|)^{-1/2} \frac{\|y\|_2}{K C_1^{1/2}} \cdot \frac{1}{K \|\psi\|_2} \leq \frac{C}{K}. \quad (4.15)$$

Likewise, we find that

$$\frac{C_y |\alpha_m|^{-1/2}}{C_\omega} \leq \frac{C_y}{C_\omega} \cdot \frac{1}{K \|\psi\|_2} \leq \frac{C}{K}. \quad (4.16)$$

The lemma is proved. $\square$

As a result, the function φ satisfies the smoothness condition

$$|\varphi(t+h) - \varphi(t)| \leq \omega(|h|), \quad h \in \mathbb{R}, \quad (4.17)$$

for all K large enough. Theorem 1.1 is, thus, completely proved.

ACKNOWLEDGMENT. The author expresses a deep gratitude to S. N. Naboko for his attention to this work and for useful remarks.

REFERENCES

- [1] E. M. Dyn'kin, S. N. Naboko, and S. I. Yakovlev, *A finiteness bound for the singular spectrum in a selfadjoint Friedrichs model*, Algebra i Analiz **3** (1991), no. 2, 77–90.
- [2] L. D. Faddeev, *On a model of Friedrichs in the theory of perturbations of the continuous spectrum*, Trudy Mat. Inst. Steklov **73** (1964), 292–313.
- [3] S. I. Iakovlev, *Perturbations of a singular spectrum in a selfadjoint Friedrichs model*, Vestnik Leningrad. Univ. Mat. Mekh. Astronom. (1990), no. 1, 116–117.
- [4] Ya. V. Mikityuk, *The singular spectrum of selfadjoint operators*, Dokl. Akad. Nauk SSSR **303** (1988), no. 1, 33–36.
- [5] S. N. Naboko, *Uniqueness theorems for operator-valued functions with positive imaginary part and the singular spectrum in the selfadjoint Friedrichs model*, Dokl. Akad. Nauk SSSR **275** (1984), no. 6, 1310–1313.

- [6] ———, *Uniqueness theorems for operator-valued functions with positive imaginary part, and the singular spectrum in the selfadjoint Friedrichs model*, Ark. Mat. **25** (1987), no. 1, 115–140.
- [7] S. N. Naboko and S. I. Yakovlev, *Conditions for the finiteness of the singular spectrum in a selfadjoint Friedrichs model*, Funktsional. Anal. i Prilozhen. **24** (1990), no. 4, 88–89.
- [8] B. S. Pavlov and S. V. Petras, *The singular spectrum of a weakly perturbed multiplication operator*, Funkcional. Anal. i Prilozhen. **4** (1970), no. 2, 54–61.
- [9] S. I. Yakovlev, *On the structure of the singular spectrum in selfadjoint Friedrichs model*, 1991, Depon. VINITI, pp. 1–25.
- [10] ———, *A finiteness bound for the singular spectrum in a neighborhood of a singular point of operators of the Friedrichs model*, Algebra i Analiz **10** (1998), no. 4, 210–237.

Serguei I. Iakovlev: Departamento de Matemáticas, Universidad Simón Bolívar,
Apartado Postal 89000, Caracas 1080-A, Venezuela

E-mail address: serguei@usb.ve