FIXED-POINT ITERATION PROCESSES FOR NON-LIPSCHITZIAN MAPPINGS OF ASYMPTOTICALLY QUASI-NONEXPANSIVE TYPE

DAYA RAM SAHU and JONG SOO JUNG

Received 5 February 2003

We study convergences of Mann and Ishikawa iteration processes for mappings of asymptotically quasi-nonexpansive type in Banach spaces.

2000 Mathematics Subject Classification: 47H09, 47H10.

1. Introduction and preliminaries. Let D be a nonempty subset of a real Banach space X and $T : D \rightarrow D$ a nonlinear mapping. The mapping T is said to be asymptotically quasi-nonexpansive (see [5]) if $F(T) \neq \emptyset$ and there exists a sequence $\{k_n\}$ in $[0, \infty)$ with $\lim_{n \rightarrow \infty} k_n = 0$ such that

$$\|T^n x - p\| \leq (1 + k_n) \|x - p\| \quad (1.1)$$

for all $x \in D$, $p \in F(T)$, and $n \in \mathbb{N}$. The mapping T is said to be asymptotically nonexpansive (see [3]) if there exists a sequence $\{k_n\}$ in $[0, \infty)$ with $\lim_{n \rightarrow \infty} k_n = 0$ such that

$$\|T^n x - T^n y\| \leq (1 + k_n) \|x - y\| \quad (1.2)$$

for all $x, y \in D$ and $n \in \mathbb{N}$. The mapping T is said to be a mapping of asymptotically nonexpansive type [4] if

$$\limsup_{n \rightarrow \infty} \sup_{x \in D} (\|T^n x - T^n y\| - \|x - y\|) \leq 0 \quad (1.3)$$

for any $y \in D$.

In 1973, Petryshyn and Williamson [7] gave necessary and sufficient conditions for Mann iterative sequence to converge to fixed points of quasi-nonexpansive mappings. In 1997, Ghosh and Debnath [2] extended the results of [7] and gave the necessary and sufficient conditions for Ishikawa iterative sequence to converge to fixed points for quasi-nonexpansive mappings.

Recently, Liu [5] extended results of [2, 7] and gave the necessary and sufficient conditions for Ishikawa iterative sequence to converge to fixed points of asymptotically quasi-nonexpansive mappings.

First, we introduce the concept of class of mappings of asymptotically quasi-nonexpansive type: the mapping T is said to be a mapping of asymptotically quasi-nonexpansive type if $F(T) \neq \emptyset$ and

$$\limsup_{n \rightarrow \infty} \sup_{x \in D} (\|T^n x - p\| - \|x - p\|) \leq 0 \quad \text{for any } p \in F(T). \quad (1.4)$$

REMARK 1.1. If T is a mapping of asymptotically nonexpansive type with $F(T) \neq \emptyset$, then T is a mapping of asymptotically quasi-nonexpansive type.

REMARK 1.2. If D is bounded and T is an asymptotically quasi-nonexpansive mapping, then T is a mapping of asymptotically quasi-nonexpansive type. In fact, if T is an asymptotically quasi-nonexpansive mapping, then there exists a sequence $\{k_n\}$ in $[0, \infty)$ with $\lim_{n \rightarrow \infty} k_n = 0$ such that

$$\|T^n x - p\| \leq (1 + k_n) \|x - p\| \quad (1.5)$$

for all $x \in D$, $p \in F(T)$, and $n \in \mathbb{N}$, which implies

$$\sup_{x \in D} \{ \|T^n x - T^n y\| - \|x - y\| \} \leq k_n \cdot \text{diam } D \quad (1.6)$$

for any $y \in F(T)$ and $n \in \mathbb{N}$. Hence

$$\limsup_{n \rightarrow \infty} \sup_{x \in D} (\|T^n x - T^n y\| - \|x - y\|) \leq 0 \quad \text{for any } y \in F(T). \quad (1.7)$$

We observe from Remarks 1.1 and 1.2 that the class of mappings of asymptotically nonexpansive type is an intermediate class between the class of mappings of asymptotically quasi-nonexpansive type and that of mappings of asymptotically nonexpansive type with nonempty fixed-point sets. Let

$$\begin{aligned} C_1 &= \{T : T : D \rightarrow D \text{ is a nonexpansive mapping}\}, \\ C_2 &= \{T : T : D \rightarrow D \text{ is a quasi-nonexpansive mapping}\}, \\ C_3 &= \{T : T : D \rightarrow D \text{ is an asymptotically nonexpansive mapping}\}, \\ C_4 &= \{T : T : D \rightarrow D \text{ is an asymptotically quasi-nonexpansive mapping}\}, \\ C_5 &= \{T : T : D \rightarrow D \text{ is a mapping of asymptotically nonexpansive type}\}, \\ C_6 &= \{T : T : D \rightarrow D \text{ is a mapping of asymptotically quasi-nonexpansive type}\}. \end{aligned} \quad (1.8)$$

Then we have the following implications:

$$\begin{array}{ccc} C_1 & \implies & C_2 \\ \Downarrow & & \Downarrow \\ C_3 & \implies & C_4 \\ \Downarrow & & \Downarrow \\ C_5 & \implies & C_6. \end{array} \quad (1.9)$$

In this paper, we are mainly interested in the problem of approximation of fixed points of the more general class of mappings of asymptotically quasi-nonexpansive type than that of asymptotically quasi-nonexpansive mappings. The purpose of this paper is to continue discussion concerning convergence of Mann and Ishikawa iteration processes for mappings of asymptotically quasi-nonexpansive type in Banach spaces. We give necessary and sufficient conditions for the Mann and Ishikawa iteration processes to converge to fixed points of mappings of asymptotically quasi-nonexpansive type. Further, we obtain extensions of various results obtained quite recently by Deng [1], Ghosh and Denath [2], Liu [5], and Tan and Xu [9, 10] to more general types of space as well as families of operators.

We say that a Banach space X satisfies *Opial's condition* [6] if, for each sequence $\{x_n\}$ in X weakly convergent to a point x and for all $y \neq x$,

$$\liminf_{n \rightarrow \infty} \|x_n - x\| < \liminf_{n \rightarrow \infty} \|x_n - y\|. \quad (1.10)$$

The examples of Banach spaces which satisfy Opial's condition are Hilbert spaces, and all $L^p[0, 2\pi]$ with $1 < p \neq 2$ fail to satisfy Opial's condition [6].

Let D be a nonempty closed convex subset of a Banach space X . Then $I - T$ is *demiclosed at zero* if, for any sequence $\{x_n\}$ in D , condition $x_n \rightarrow x$ weakly and $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ implies $(I - T)x = 0$.

2. Main results. In this section, we establish some weak and strong convergences for mappings of asymptotically quasi-nonexpansive type in Banach spaces.

LEMMA 2.1. *Let D be a nonempty subset of a normed space X and let $T : D \rightarrow E$ be a mapping of asymptotically quasi-nonexpansive type. For two given real sequences $\{\alpha_n\}$ and $\{\beta_n\}$ in $[0, 1]$, let a sequence $\{x_n\}$ in D be defined by*

$$\begin{aligned} x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T y_n, \\ y_n &= (1 - \beta_n)x_n + \beta_n T x_n, \quad n = 1, 2, \dots \end{aligned} \quad (2.1)$$

If p is a fixed point of T , then

- (a) $\|x_{n+1} - p\| \leq \|x_n - p\| + (1 + \beta_n) \sup_{x \in D} (\|T^n x - p\| - \|x - p\|)$, $n = 1, 2, \dots$,
- (b) $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists.

PROOF. Let p be a fixed point of T .

(a) From (2.1), we have

$$\begin{aligned} \|x_{n+1} - p\| &\leq (1 - \alpha_n)\|x_n - p\| + \alpha_n\|T^n y_n - p\| \\ &\leq (1 - \alpha_n)\|x_n - p\| + \alpha_n(\|T^n y_n - p\| \\ &\quad - \|y - p\|) + \alpha_n\|y_n - p\| \end{aligned}$$

$$\begin{aligned}
&\leq (1 - \alpha_n) \|x_n - p\| + (\|T^n y_n - p\| - \|y_n - p\|) \\
&\quad + \alpha_n ((1 - \beta_n) \|x_n - p\| + \beta_n \|T^n x_n - p\|) \\
&\leq \|x_n - p\| + (\|T^n y_n - p\| - \|y_n - p\|) \\
&\quad + \beta_n (\|T^n x_n - p\| - \|x_n - p\|) \\
&\leq \|x_n - p\| + (1 + \beta_n) \sup_{x \in D} (\|T^n x - p\| - \|x - p\|).
\end{aligned} \tag{2.2}$$

(b) For $m, n \in \mathbb{N}$, we have

$$\begin{aligned}
\|x_{n+m} - p\| &\leq \|x_{n+m-1} - p\| + 2 \sup_{x \in D} (\|T^{n+m-1} x - p\| - \|x - p\|) \\
&\leq \|x_{n+m-1} - p\| + 2 \sup_{x \in D} (\|T^m x - p\| - \|x - p\|) \\
&\leq \|x_{n+m-2} - p\| + 4 \sup_{x \in D} (\|T^m x - p\| - \|x - p\|) \\
&\leq \cdots \leq \|x_n - p\| + 2n \sup_{x \in D} (\|T^m x - p\| - \|x - p\|).
\end{aligned} \tag{2.3}$$

Hence, for $n \in \mathbb{N}$,

$$\begin{aligned}
\limsup_{m \rightarrow \infty} \|x_m - p\| &\leq \|x_n - p\| + 2n \limsup_{m \rightarrow \infty} (\|T^m x - p\| - \|x - p\|) \\
&\leq \|x_n - p\|.
\end{aligned} \tag{2.4}$$

It follows that

$$\limsup_{m \rightarrow \infty} \|x_m - p\| \leq \liminf_{n \rightarrow \infty} \|x_n - p\|. \tag{2.5}$$

Thus $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists. $\square$

LEMMA 2.2. Let D and T be as in [Lemma 2.1](#). For a given real sequence $\{\alpha_n\}$ in $[0, 1]$, let a sequence $\{x_n\}$ in D be defined by

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n, \quad n = 1, 2, \dots \tag{2.6}$$

If p is a fixed point of T , then

- (a) $\|x_{n+1} - p\| \leq \|x_n - p\| + \sup_{x \in D} (\|T^n x - p\| - \|x - p\|)$, $n = 1, 2, \dots$,
- (b) $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists.

THEOREM 2.3. Let X be a Banach space which satisfies Opial's condition and let D be a weakly compact subset of X . Let T and $\{x_n\}$ be as in [Lemma 2.1](#). Suppose that T has a fixed point, $I - T$ is demiclosed at zero, and $\{x_n\}$ is an approximating fixed-point sequence for T , that is, $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$. Then $\{x_n\}$ converges weakly to a fixed point of T .

PROOF. First, we show that $\omega_w(x_n) \subset F(T)$. Let $x_{n_k} \rightarrow x$ weakly. By assumption, we have $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$. Since $I - T$ is demiclosed at zero,

$x \in F(T)$. By Opial's condition, $\{x_n\}$ possesses only one weak limit point, that is, $\{x_n\}$ converges weakly to a fixed point of T . $\square$

THEOREM 2.4. *Let X be a Banach space which satisfies Opial's condition and let D be a weakly compact subset of X . Let T and $\{x_n\}$ be as in [Lemma 2.2](#). Suppose that T has a fixed point, $I - T$ is demiclosed at zero, and $\{x_n\}$ is an approximating fixed-point sequence for T , that is, $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$. Then $\{x_n\}$ converges weakly to a fixed point of T .*

REMARK 2.5. [Theorem 2.3](#) improves Theorem 2 of Deng [[1](#)] for mappings of asymptotically quasi-nonexpansive type. [Theorem 2.4](#) generalizes Theorem 2.1 of Schu [[8](#)].

THEOREM 2.6. *Let D be a closed subset of Banach space, let $T : D \rightarrow D$ be a mapping of asymptotically quasi-nonexpansive type, and $F(T)$ be nonempty closed set. For two given real sequences $\{\alpha_n\}$ and $\{\beta\}$ in $[0, 1]$, let the Ishikawa iterative sequence $\{x_n\}$ in D be defined by [\(2.1\)](#). Then $\{x_n\}$ converges strongly to a fixed point of T if and only if $\liminf_n d(x_n, F(T)) = 0$.*

PROOF. Let $\{x_n\}$ converge strongly to a point $z \in F(T)$. Then $\lim_n d(x_n, F(T)) = 0$. Conversely, suppose $\liminf_n d(x_n, F(T)) = 0$. From [Lemma 2.1\(a\)](#),

$$\|x_{n+1} - p\| \leq \|x_n - p\| + 2 \sup_{x \in D} (\|T^n x - p\| - \|x - p\|) \quad (2.7)$$

for any $n \in \mathbb{N}$ and $p \in F(T)$. Since T is a mapping of asymptotically quasi-nonexpansive type, we have

$$\limsup_n \left\{ \sup_{k \geq n} (\|T^k x - p\| - \|x - p\|) \right\} \leq 0. \quad (2.8)$$

Hence, there exists a positive integer n_0 and a sequence $\{a_n\}$ of positive real numbers with $\lim_n a_n = 0$ such that

$$\sup_{k \geq n} \left\{ \sup_{x \in D} (\|T^k x - p\| - \|x - p\|) \right\} \leq a_n \quad (2.9)$$

for any $n \geq n_0$. Without loss of generality, we can assume that $a_n = 1/2n^2$. Hence,

$$\sup_{k \geq n} \left\{ \sup_{x \in D} (\|T^k x - p\| - \|x - p\|) \right\} \leq \frac{1}{2n^2} \quad (2.10)$$

for any $n \geq n_0$. It follows from [\(2.7\)](#) that

$$\|x_{n+1} - p\| \leq \|x_n - p\| + \frac{1}{n^2} \quad (2.11)$$

for all $n \geq n_0$, that is,

$$d(x_{n+1}, F(T)) \leq d(x_n, F(T)) + \frac{1}{n^2} \quad (2.12)$$

for all $n \geq n_0$. Hence for $n, m \geq n_0$, we have

$$d(x_{n+m}, F(T)) \leq d(x_n, F(T)) + \sum_{i=n}^{n+m-1} \frac{1}{i^2}. \quad (2.13)$$

Using [10, Lemma 1, page 303], we obtain that $\lim_n d(x_n, F(T))$ exists, and it follows from $\liminf_n d(x_n, F(T)) = 0$ that $\lim_n d(x_n, F(T)) = 0$. Thus, $\lim_n d(x_n, F(T)) = 0$. For each $\varepsilon > 0$, there exists a natural number m_0 such that

$$d(x_n, F(T)) < \frac{\varepsilon}{3} \quad (2.14)$$

for all $n \geq m_0$. Then there exists a $p' \in F(T)$ such that $d(x_n, p') < \varepsilon/2$ for all $n \geq m_0$. If $n, m \geq m_0$, then

$$d(x_n, x_m) \leq d(x_n, p') + d(p', x_m) \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \quad (2.15)$$

This shows that $\{x_n\}$ is a Cauchy sequence in D . Let $\lim_n x_n = v \in D$. Since $F(T) \subset D$ is closed and $\lim_n d(x_n, F(T)) = 0$, we conclude that $v \in F(T)$. This completes the proof. $\square$

As a consequence of [Theorem 2.6](#), we obtain the following result.

THEOREM 2.7. *Let D be a closed subset of Banach space, let $T : D \rightarrow D$ be a mapping of asymptotically quasi-nonexpansive type, and let $F(T)$ be a nonempty closed set. For a given sequence $\{\alpha_n\}$ in $[0, 1]$, let the Mann iterative sequence $\{x_n\}$ in D be defined by (2.6). Then $\{x_n\}$ converges strongly to a fixed point of T if and only if $\liminf_n d(x_n, F(T)) = 0$.*

REMARK 2.8. Theorems 2.6 and 2.7 extend corresponding results of Ghosh and Debnath [2], Liu [5], and Petryshyn and Williamson [7] from quasi-nonexpansive or asymptotically quasi-nonexpansive mapping to large class of non-Lipschitzian mappings.

ACKNOWLEDGMENT. The first author wishes to acknowledge the financial support of the Department of Science and Technology, India, made in the program year 2002–2003, Project No. SR/FTP/MS-15. The second author was supported by Korea Research Foundation Grant KRF-2000-DP0013.

REFERENCES

- [1] L. Deng, *Convergence of the Ishikawa iteration process for nonexpansive mappings*, J. Math. Anal. Appl. **199** (1996), no. 3, 769–775.

- [2] M. K. Ghosh and L. Debnath, *Convergence of Ishikawa iterates of quasi-nonexpansive mappings*, J. Math. Anal. Appl. **207** (1997), no. 1, 96–103.
- [3] K. Goebel and W. A. Kirk, *A fixed point theorem for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. **35** (1972), 171–174.
- [4] W. A. Kirk, *Fixed point theorems for non-Lipschitzian mappings of asymptotically nonexpansive type*, Israel J. Math. **17** (1974), 339–346.
- [5] Q. Liu, *Iterative sequences for asymptotically quasi-nonexpansive mappings*, J. Math. Anal. Appl. **259** (2001), no. 1, 1–7.
- [6] Z. Opial, *Weak convergence of the sequence of successive approximations for non-expansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591–597.
- [7] W. V. Petryshyn and T. E. Williamson Jr., *Strong and weak convergence of the sequence of successive approximations for quasi-nonexpansive mappings*, J. Math. Anal. Appl. **43** (1973), 459–497.
- [8] J. Schu, *Weak and strong convergence to fixed points of asymptotically nonexpansive mappings*, Bull. Austral. Math. Soc. **43** (1991), no. 1, 153–159.
- [9] K.-K. Tan and H. K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, J. Math. Anal. Appl. **178** (1993), no. 2, 301–308.
- [10] ———, *Fixed point iteration processes for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. **122** (1994), no. 3, 733–739.

Daya Ram Sahu: Department of Applied Mathematics, Shri Shankaracharya College of Engineering, Junwani, Bhilai 490 020, India

E-mail address: sahudr@rediffmail.com

Jong Soo Jung: Department of Mathematics, Dong-A University, Pusan 604-714, Korea

E-mail address: jungjs@mail.donga.ac.kr