ABSOLUTELY CONTINUOUS MEASURES AND COMPACT COMPOSITION OPERATOR ON SPACES OF CAUCHY TRANSFORMS

YUSUF ABU MUHANNA and EL-BACHIR YALLAOUI

Received 5 October 2003

The analytic self-map of the unit disk $\mathbf{D}$, φ is said to induce a composition operator C_φ from the Banach space X to the Banach space Y if $C_\varphi(f) = f \circ \varphi \in Y$ for all $f \in X$. For $z \in \mathbf{D}$ and $\alpha > 0$, the families of weighted Cauchy transforms F_α are defined by $f(z) = \int_{\mathbf{T}} K_x^\alpha(z) d\mu(x)$, where $\mu(x)$ is complex Borel measure, x belongs to the unit circle $\mathbf{T}$, and the kernel $K_x(z) = (1 - \bar{x}z)^{-1}$. In this paper, we will explore the relationship between the compactness of the composition operator C_φ acting on F_α and the complex Borel measures $\mu(x)$.

2000 Mathematics Subject Classification: 30E20, 30D99.

1. Background. Let $\mathbf{T}$ be the unit circle and $\mathbf{M}$ the set of all complex-valued Borel measures on $\mathbf{T}$. For $\alpha > 0$ and $z \in \mathbf{D}$, we define the space of weighted Cauchy transforms F_α to be the family of all functions $f(z)$ such that

$$f(z) = \int_{\mathbf{T}} K_x^\alpha(z) d\mu(x), \quad (1.1)$$

where the Cauchy kernel $K_x(z)$ is given by

$$K_x(z) = \frac{1}{1 - \bar{x}z} \quad (1.2)$$

and where μ in (1.1) varies over all measures in $\mathbf{M}$. The class F_α is a Banach space with respect to the norm

$$\|f\|_{F_\alpha} = \inf \|\mu\|_{\mathbf{M}}, \quad (1.3)$$

where the infimum is taken over all Borel measures μ satisfying (1.1). $\|\mu\|$ denotes the total variation norm of μ . The family F_1 has been studied extensively in the Soviet literature. The generalizations for $\alpha > 0$ were defined by MacGregor [7]. The Banach spaces F_α have been well studied in [2, 3, 5, 7]. Among the properties of F_α we list the following:

- (i) $F_\alpha \subset F_\beta$ whenever $0 < \alpha < \beta$,
- (ii) F_α is Möbius invariant,
- (iii) $f \in F_\alpha$ if and only if $f' \in F_{1+\alpha}$ and $\|f'\|_{F_{1+\alpha}} \leq \alpha \|f\|_{F_\alpha}$,
- (iv) if $g \in F_{\alpha+1}$, then $f(z) = \int_0^z g(w) dw \in F_\alpha$ and $\|f\|_{F_\alpha} \leq (2/\alpha) \|g\|_{F_{1+\alpha}}$.

The space F_α may be identified with $\mathbf{M}/H_0^1$, the quotient of the Banach space $\mathbf{M}$ of Borel measures by H_0^1 , the subspace of L^1 consisting of functions with mean value zero

whose conjugate belongs to the Hardy space H^1 . Hence, F_α is isometrically isomorphic to $\mathbf{M}/\overline{H_0^1}$. Furthermore, $\mathbf{M}$ admits a decomposition $\mathbf{M} = L^1 \oplus \mathbf{M}_s$, where $\mathbf{M}_s$ is the space of Borel measures, which are singular with respect to Lebesgue measure, and $\overline{H_0^1} \subset L^1$. According to the Lebesgue decomposition theorem, any $\mu \in \mathbf{M}$ can be written as $\mu = \mu_a + \mu_s$, where μ_a is absolutely continuous with respect to the Lebesgue measure and μ_s is singular with respect to the Lebesgue measure ($\mu_a \perp \mu_s$). Since $|x| = 1$ in (1.1), if we let $x = e^{it}$, then we can decompose μ in such a way that $d\mu(e^{it}) = h(e^{it})dt + d\mu_s(e^{it})$, where $h(e^{it}) \in L^1$. Consequently, F_α is isomorphic to $L^1/\overline{H_0^1} \oplus \mathbf{M}_s$. Hence, F_α can be written as $F_\alpha = F_{\alpha a} \oplus F_{\alpha s}$, where $F_{\alpha a}$ is isomorphic to $L^1/\overline{H_0^1}$, the closed subspace of $\mathbf{M}$ of absolutely continuous measures, and $F_{\alpha s}$ is isomorphic to $\mathbf{M}_s$, the subspace of $\mathbf{M}$ of singular measures. If $f \in F_{\alpha a}$, then the singular part is null and the measure μ for which (1.1) holds reduces to $d\mu(t) = d\mu(e^{it}) = h(e^{it})dt$, where $h(e^{it}) \in L^1$ and dt is the Lebesgue measure on $\mathbf{T}$, see [1]. The functions in $F_{\alpha a}$ may be then written as, for $x = e^{it}$,

$$f(z) = \int_{-\pi}^{\pi} K_x^\alpha(z) h(e^{it}) dt. \quad (1.4)$$

Furthermore, if $h(e^{it})$ is nonnegative, then

$$\|f\|_{F_\alpha} = \|h(e^{it})\|_{L^1}. \quad (1.5)$$

REMARK 1.1. For simplicity, we will adopt the following notation throughout the paper. We will reserve μ for the Borel measures of $\mathbf{M}$, and since in (1.1) $|x| = 1$, we can write $x = e^{it}$, where $t \in [-\pi, \pi)$. We will reserve dt for the normalized Lebesgue of the unit circle $\mathbf{T}$, and $d\sigma$ for the singular part of $d\mu$. Hence, instead of writing $d\mu(e^{it}) = d\mu_a(e^{it}) + d\mu_s(e^{it}) = h(e^{it})dt + d\mu_s(e^{it})$, we may simply write $d\mu(t) = hdt + d\sigma(t)$.

2. Introduction. If X and Y are Banach spaces, and L is a linear operator from X to Y , we say that L is bounded if there exists a positive constant A such that $\|L(f)\|_Y \leq A\|f\|_X$ for all f in X . We denote by $C(X, Y)$ the set of all bounded linear operators from X to Y . If $L \in C(X, Y)$, we say that L is a compact operator from X to Y if the image of every bounded set of X is relatively compact (i.e., has compact closure) in Y . Equivalently, a linear operator L is a compact operator from X to Y if and only if for every bounded sequence $\{f_n\}$ of X , $\{L(f_n)\}$ has a convergent subsequence in Y . We will denote by $K(X, Y)$ the subset of $C(X, Y)$ of compact linear operators from X into Y .

Let $H(\mathbf{D})$ denote the set of all analytic functions on the unit disk $\mathbf{D}$ and map $\mathbf{D}$ into $\mathbf{D}$. If X and Y are Banach spaces of functions on the unit disk $\mathbf{D}$, we say that $\varphi \in H(\mathbf{D})$ induces a bounded composition operator $C_\varphi(f) = f(\varphi)$ from X to Y , if $C_\varphi \in C(X, Y)$ or equivalently $C_\varphi(X) \subseteq Y$, and there exists a positive constant A for all $f \in X$ and $\|C_\varphi(f)\|_Y \leq A\|f\|_X$. In case $X = Y$, then we say that φ induces a composition operator C_φ on X . If $f \in X$, then $C_\varphi(f) = f(\varphi) \in X$. Similarly, we say that $\varphi \in H(\mathbf{D})$ induces a compact composition operator if $C_\varphi \in K(X, Y)$.

A fundamental problem that has been studied concerning composition operators is to relate function theoretic properties of φ to operator theoretic properties of the restriction of C_φ to various Banach spaces of analytic functions. However, since the

spaces of Cauchy transforms are defined in terms of Borel measures, it seems natural to investigate the relation between the behavior of the composition operator and the measure. The work in this paper was motivated by the work of Cima and Matheson in [1], who showed that C_φ is compact on F_1 if and only if $C_\varphi(F_1) \subset F_{1a}$. In our work, we will generalize this result for $\alpha > 1$.

Now, if $C_\varphi \in C(F_\alpha, F_\alpha)$, then $C_\varphi(f) = (f \circ \varphi) = f(\varphi) \in F_\alpha$ for all $f \in F_\alpha$ and there exists a positive constant A such that

$$\|C_\varphi(f)\|_{F_\alpha} = \|f(\varphi)\|_{F_\alpha} = \|[\mu] \| \leq A \|f\|_{F_\alpha}. \quad (2.1)$$

Since F_α can be identified with the quotient space $\mathbf{M}/\overline{H_0^1}$, we can view C_φ as a map,

$$\begin{aligned} C_\varphi : \mathbf{M}/\overline{H_0^1} &\longrightarrow \mathbf{M}/\overline{H_0^1}, \\ f &\longmapsto f(\varphi). \end{aligned} \quad (2.2)$$

The equivalence class of a complex measure μ will be written as

$$\begin{aligned} [\mu] &= \mu + \overline{H_0^1} = \{\mu + \bar{h} : h \in H_0^1\}, \\ \|[\mu] \| &= \inf_h \|\mu + \bar{h}\|. \end{aligned} \quad (2.3)$$

The space $C(F_\alpha, F_\alpha)$ has been studied in [4], where Hibschweiler showed that

- (1) if $\alpha \geq 1$, then $C_\varphi \in C(F_\alpha, F_\alpha)$ for any analytic self map φ of the unit disc,
- (2) $C_\varphi \in C(F_\alpha, F_\alpha)$ if and only if $\{K_X^\alpha(\varphi) : |x| = 1\}$ is a norm bounded subset of F_α ,
- (3) if $C_\varphi \in C(F_\alpha, F_\alpha)$, then $C_\varphi \in C(F_\beta, F_\beta)$ for $0 < \alpha < \beta$,
- (4) if $C_\varphi \in C(F_\alpha, F_\alpha)$, then the operator $\varphi' C_\varphi \in C(F_{\alpha+1}, F_{\alpha+1})$.

In this paper, we will investigate necessary and sufficient conditions for C_φ to be compact on F_α for $\alpha \geq 1$. Since F_α is Mobius invariant, then there is no loss of generality in assuming that $\varphi(0) = 0$.

3. Compactness and absolutely continuous measures. In this section, we will show that the compactness of the composition operator C_φ on F_α is strongly tied with the absolute continuity of the measure that supports it. First, we state this lemma due to [6].

LEMMA 3.1. *If $0 < \alpha < \beta$, then $F_\alpha \subset F_{\beta a}$ and the inclusion map is a compact operator of norm one.*

Next, we show that the bounded function of F_α belongs to $F_{\alpha a}$ for $\alpha \geq 1$.

PROPOSITION 3.2. *$H^\infty \cap F_\alpha \subset F_{\alpha a}$ for $\alpha \geq 1$.*

PROOF. Suppose that $f \in H^\infty \cap F_\alpha$, where $\alpha \geq 1$. Then, using the previous lemma and the fact that $H^\infty \subset F_{1a}$, we get that for any $z \in \mathbf{D}$, $f(z) \in H^\infty \cap F_\alpha \subset F_{1a} \cap F_\alpha \subseteq F_{\alpha a}$. $\square$

In the following theorem, $g_x(e^{it})$ is L^1 continuous function of x means,

$$\|g_x(e^{it}) - g_y(e^{it})\|_{L^1} \rightarrow 0 \quad \text{as } x \rightarrow y, \quad (3.1)$$

$g_x(e^{it})$ is M^+ continuous, if

$$\left\| \int_x g_x(e^{it}) d\mu_n(x) - \int_x g_x(e^{it}) d\mu(x) \right\|_{L^1} \rightarrow 0, \quad (3.2)$$

for any sequence $\{\mu_n\}$ of nonnegative Borel measures that are weak* convergent to μ .

THEOREM 3.3. *For a holomorphic self-map φ of the unit disc $\mathbf{D}$ and $\alpha \geq 1$, if C_φ is compact on F_α , then $(C_\varphi \circ K_x^\alpha)(z) \in F_{\alpha\alpha}$ and*

$$(C_\varphi \circ K_x^\alpha)(z) = \int_{-\pi}^{\pi} g_x(e^{it}) K_x^\alpha(z) dt, \quad (3.3)$$

where $g_x(e^{it})$ is a function of two variables x , with $|x| = 1$ and e^{it} , $\|g_x(e^{it})\|_{L^1} \leq a < \infty$, $g_x(e^{it})$ is nonnegative, and M^+ is a continuous function of x .

PROOF. Assume that C_φ is compact and let $\{f_j\}_{j=1}^\infty$ be a sequence of functions such that

$$f_j(z) = K_x^\alpha(\rho_j z) = \frac{1}{(1 - \rho_j \bar{x}z)^\alpha}, \quad (3.4)$$

where $0 < \rho_j < 1$ and $\lim_{j \rightarrow \infty} \rho_j = 1$. Then, it is known from [2] that $f_j(z) \in F_\alpha$ for every j , and $\|f_j(z)\|_{F_\alpha} = 1$. Furthermore, there exist $\mu_j \in \mathbf{M}$, such that $\|\mu_j\| = 1$, $d\mu_j > 0$, and

$$\begin{aligned} f_j(z) &= \frac{1}{(1 - \rho_j \bar{x}z)^\alpha} \\ &= \int_{\mathbf{T}} K_{e^{it}}^\alpha(z) d\mu_j(e^{it}) \\ &= \int_{\mathbf{T}} \frac{1}{(1 - e^{-it}z)^\alpha} d\mu_j(e^{it}). \end{aligned} \quad (3.5)$$

Since C_φ is compact on F_α , then $(C_\varphi \circ f_j) \in F_\alpha$ and $\|C_\varphi(f_j)\| \leq \|C_\varphi\| \|f_j\|_{F_\alpha} = \|C_\varphi\|$ for all j . Furthermore, $C_\varphi \circ f_j \in H^\infty$, therefore using the previous result, we get that $(C_\varphi \circ f_j) \in H^\infty \cap F_\alpha \subset F_{\alpha\alpha}$ for every j . Therefore, there exists L^1 a nonnegative function $g_x^j(e^{it})$ such that

$$\begin{aligned} d\mu_j(e^{it}) &= g_x^j(e^{it}) dt, \quad \|g_x^j\|_{L^1} \leq \|C_\varphi\|, \\ (f_j \circ \varphi)(z) &= K_x^\alpha(\rho_j \varphi) = \int_{-\pi}^{\pi} g_x^j(e^{it}) K_{e^{it}}^\alpha(z) dt. \end{aligned} \quad (3.6)$$

Now, because $F_{\alpha\alpha}$ is closed and C_φ is compact, the sequence $\{f_j \circ \varphi\}_{j=1}^\infty$ has a convergent subsequence $\{f_{j_k} \circ \varphi\}$ that converges to $(K_x^\alpha \circ \varphi)(z) \in F_{\alpha\alpha}$. Therefore,

$$\begin{aligned} \lim_{k \rightarrow \infty} (f_{j_k} \circ \varphi)(z) &= \lim_{k \rightarrow \infty} K_x^\alpha(\rho_{j_k} \varphi) \\ &= \lim_{k \rightarrow \infty} \int_{-\pi}^{\pi} g_x^{j_k}(e^{it}) K_{e^{it}}^\alpha(z) dt \\ &= \int_{-\pi}^{\pi} g_x(e^{it}) K_{e^{it}}^\alpha(z) dt \\ &= (K_x^\alpha \circ \varphi)(z) = \frac{1}{(1 - \bar{x}\varphi(z))^\alpha} \in F_{\alpha\alpha}, \end{aligned} \quad (3.7)$$

where the function $g_x(e^{it})$ is an L^1 nonnegative continuous function of x , and $\|g_x\|_{L^1} \leq \|C_\varphi\|$. The continuity of $g_x(e^{it})$ in M^+ follows from the compactness and the fact that if $f(z) = \int_{\mathbf{T}} K_x^\alpha(z) d\mu(x)$, with $d\mu(x) > 0$, then $\|f\|_{F_\alpha} = \|\mu\|$.

This concludes the proof. $\square$

COROLLARY 3.4. *Let $g_x(e^{it})$ be as in Theorem 3.3. Then, for $h(x) \in \overline{H_0^1}$, the operator $\int g_x(e^{it})h(x)dx = u(e^{it}) \in \overline{H_0^1}$ is bounded on $\overline{H_0^1}$.*

PROOF. For the operator to be well defined, $\int (h(x)dx/(1-\bar{x}\varphi(z))^\alpha) = 0$ for all $h(x) \in \overline{H_0^1}$. Hence, $\int g_x(e^{it})h(x)dx = u(e^{it}) \in \overline{H_0^1}$. $\square$

In the opposite direction, we have the following lemma.

LEMMA 3.5. *Let $g_x(e^{it})$ be a nonnegative M^+ continuous function such that $\|g_x\|_{L^1} \leq a < \infty$ and $g_x(e^{it})$ defines a bounded operator on $\overline{H_0^1}$. If $f(z) = \int (1/(1-\bar{x}z)^\alpha) d\mu(x)$, let L be the operator given by*

$$L[f(z)] = \iint \frac{g_x(e^{it})}{(1-e^{-it}z)^\alpha} dt d\mu(x), \quad (3.8)$$

then L is compact operator on F_α , $\alpha \geq 1$.

PROOF. First, note that the condition that $g_x(e^{it})$ defines a bounded operator on $\overline{H_0^1}$ implies that the L operator is a well-defined function on F_α . Let $\{f_n(z)\}$ be a bounded sequence in F_α and let $\{\mu_n\}$ be the corresponding norm bounded sequence of measures in $\mathbf{M}$. Since every norm bounded sequence of measures in $\mathbf{M}$ has a weak* convergent subsequence, let $\{\mu_n\}$ be such subsequence that is convergent to $\mu \in \mathbf{M}$. We want to show that $\{L(f_n)\}$ has a convergent subsequence in F_α .

First, we assume that $d\mu_n(x) > 0$ for all n , and let $w_n(t) = \int g_x(e^{it})d\mu_n(x)$ and $w(t) = \int g_x(e^{it})d\mu(x)$, then, as $g_x(e^{it})$ is an M^+ continuous function, we have $w_n(t), w(t) \in L^1$ for all n , and $w_n(t) \rightarrow w(t)$ in L^1 . Now, since $g_x(e^{it})$ is a nonnegative continuous function in x and $\{\mu_n\}$ is weak* convergent to μ , then

$$\begin{aligned} L(f_n(z)) &= \iint \frac{g_x(e^{it})d(t)}{(1-e^{-it}z)^\alpha} d\mu_n(x) = \int \frac{w_n(t)}{(1-e^{-it}z)^\alpha} dt, \\ L(f(z)) &= \iint \frac{g_x(e^{it})d(t)}{(1-e^{-it}z)^\alpha} d\mu(x) = \int \frac{w(t)}{(1-e^{-it}z)^\alpha} dt. \end{aligned} \quad (3.9)$$

Furthermore, because $w_n(t)$ is nonnegative, then

$$\begin{aligned} \|L(f_n)\|_{F_\alpha} &= \|w_n\|_{L^1}, \\ \|L(f)\|_{F_\alpha} &= \|w\|_{L^1}. \end{aligned} \quad (3.10)$$

Now, since $\|w_n - w\|_{L^1} \rightarrow 0$, then $\|L(f_n) - L(f)\|_{F_\alpha} \rightarrow 0$, which shows that $\{L(f_n)\}$ has a convergent subsequence in F_α and thus L is a compact operator for the case where μ is a positive measure.

In the case where μ is complex measure, we write

$$d\mu_n(x) = (d\mu_n^1(x) - d\mu_n^2(x)) + i(d\mu_n^3(x) - d\mu_n^4(x)), \quad (3.11)$$

where each $d\mu_n^j(x) > 0$, and define $w_n^j(t) = \int g_x(e^{it})d\mu_n^j(x)$, then

$$w_n(t) = \int g_x(e^{it})d\mu_n(x) = (w_n^1(t) - w_n^2(t)) + i(w_n^3(t) - w_n^4(t)). \quad (3.12)$$

Using an argument similar to the one above, we get that $w_n^j(t), w^j(t) \in L^1$, and $\|w_n^j - w^j\|_{L^1} \rightarrow 0$. Consequently, $\|w_n - w\|_{L^1} \rightarrow 0$, where $w(t) = (w^1(t) - w^2(t)) + i(w^3(t) - w^4(t)) = \int g_x(e^{it})d\mu(x)$.

Hence, $\|L(f_n) - L(f)\|_{F_\alpha} \leq \|w_n - w\|_{L^1} \rightarrow 0$.

Finally, we conclude that the operator is compact. $\square$

The following is the converse of [Theorem 3.3](#).

THEOREM 3.6. *For a holomorphic self-map φ of the unit disc $\mathbf{D}$, if*

$$(K_x^\alpha \circ \varphi)(z) = \frac{1}{(1 - \bar{x}\varphi(z))^\alpha} = \int \frac{g_x(e^{it})}{(1 - e^{-it}z)^\alpha} dt, \quad g_x \in L^1 \quad (3.13)$$

is nonnegative, $\|g_x\|_{L^1} \leq a < \infty$ for all $x \in \mathbf{T}$ and g_x is an M^+ continuous function, then C_φ is compact on F_α .

PROOF. We want to show that C_φ is compact on F_α . Let $f(z) \in F_\alpha$. Then, there exists a measure μ in $\mathbf{M}$ such that for every z in D ,

$$f(z) = \int \frac{1}{(1 - \bar{x}z)^\alpha} d\mu(x). \quad (3.14)$$

Using the assumption of the theorem, we get that

$$(f \circ \varphi)(z) = \int \frac{1}{(1 - \bar{x}\varphi(z))^\alpha} d\mu_n(x) = \iint \frac{g_x(e^{it})}{(1 - e^{-it}z)^\alpha} dt d\mu_n(x), \quad (3.15)$$

which by the previous lemma was shown to be compact on F_α . $\square$

Now, we give some examples.

COROLLARY 3.7. *Let $\varphi \in H(\mathbf{D})$, with $\|\varphi\|_\infty < 1$. Then, C_φ is compact on F_α , $\alpha \geq 1$.*

PROOF. $(C_\varphi \circ K_x^\alpha)(z) = 1/(1 - \bar{x}\varphi(z))^\alpha \in H^\infty \cap F_\alpha \subset F_{\alpha a}$ and is subordinate to $1/(1 - z)^\alpha$, hence,

$$(C_\varphi \circ K_x^\alpha)(z) = \int K_x^\alpha(z) g_x(e^{it}) dt \quad (3.16)$$

with $g_x(e^{it}) \geq 0$ and since $1 = (C_\varphi \circ K_x^\alpha)(0) = \int g_x(e^{it}) dt$, we get that $\|g_x(e^{it})\|_1 = 1$. $\square$

REMARK 3.8. In fact, one can show that C_φ , as in the above corollary, is compact from F_α , $\alpha \geq 1$, into F_1 . In other words, a contraction.

COROLLARY 3.9. *If C_φ is compact on F_α , $\alpha \geq 1$, and $\lim_{r \rightarrow 1} |\varphi(re^{i\theta})| = 1$, then $|1/\varphi'(e^{i\theta})| = 0$.*

PROOF. If C_φ is compact, then

$$(C_\varphi \circ K_x^\alpha)(z) = \int K_x^\alpha(z) g_x(e^{it}) dt. \quad (3.17)$$

Hence, if $z = e^{i\theta}$ and $\varphi(e^{i\theta}) = x$, then

$$\lim_{r \rightarrow 1} \frac{(e^{i\theta} - re^{i\theta})^\alpha}{(1 - \bar{x}\varphi(re^{i\theta}))^\alpha} = 0. \quad (3.18)$$

□

COROLLARY 3.10. *If $C_\varphi \in K(F_\alpha, F_\alpha)$ for $\alpha \geq 1$, then C_φ is a contraction.*

4. Miscellaneous results. We first start by giving another characterization of compactness on F_α .

LEMMA 4.1. *Let $\varphi \in C(F_\alpha, F_\alpha)$, $\alpha > 0$, then $\varphi \in K(F_\alpha, F_\alpha)$ if and only if for any bounded sequence (f_n) in F_α , with $f_n \rightarrow 0$ uniformly on compact subsets of $\mathbf{D}$ as $n \rightarrow \infty$, $\|C_\varphi(f_n)\|_{F_\alpha} \rightarrow 0$ as $n \rightarrow \infty$.*

PROOF. Suppose $C_\varphi \in K(F_\alpha, F_\alpha)$ and let (f_n) be a bounded sequence (f_n) in F_α with $\lim_{n \rightarrow \infty} f_n \rightarrow 0$ uniformly on compact subsets of $\mathbf{D}$. If the conclusion is false, then there exists an $\epsilon > 0$ and a subsequence $n_1 < n_2 < n_3 < \dots$ such that

$$\|C_\varphi(f_{n_j})\|_{F_\alpha} \geq \epsilon, \quad \forall j = 1, 2, 3, \dots \quad (4.1)$$

Since (f_n) is bounded and C_φ is compact, one can find another subsequence $n_{j_1} < n_{j_2} < n_{j_3} < \dots$ and f in F_α such that

$$\lim_{k \rightarrow \infty} \|C_\varphi(f_{n_{j_k}}) - f\|_{F_\alpha} = 0. \quad (4.2)$$

Since point functional evaluations are continuous in F_α , then for any $z \in \mathbf{D}$ there exists $A > 0$ such that

$$|(C_\varphi(f_{n_{j_k}}) - f)(z)| \leq A \|C_\varphi(f_{n_{j_k}}) - f\|_{F_\alpha} \rightarrow 0 \quad \text{as } k \rightarrow \infty. \quad (4.3)$$

Hence,

$$\lim_{k \rightarrow \infty} [C_\varphi(f_{n_{j_k}}) - f] \rightarrow 0 \quad (4.4)$$

uniformly on compact subsets of $\mathbf{D}$. Moreover, since $f_{n_{j_k}} \rightarrow 0$ uniformly on compact subsets of $\mathbf{D}$, then $f = 0$, that is, $C_\varphi(f_{n_{j_k}}) \rightarrow 0$ on compact subsets of F_α . Hence,

$$\lim_{k \rightarrow \infty} \|C_\varphi(f_{n_{j_k}})\|_{F_\alpha} = 0, \quad (4.5)$$

which contradicts our assumption. Thus, we must have

$$\lim_{n \rightarrow \infty} \|C_\varphi(f_n)\|_{F_\alpha} = 0. \quad (4.6)$$

Conversely, let (f_n) be a bounded sequence in the closed unit ball of F_α . We want to show that $C_\varphi(f_n)$ has a norm convergent subsequence. The closed unit ball of F_α is a compact subset of F_α in the topology of uniform convergence on the compact subsets of $\mathbf{D}$. Therefore, there is a subsequence (f_{n_k}) such that

$$f_{n_k} \rightarrow f \quad (4.7)$$

uniformly on compact subsets of D . Hence, by hypothesis,

$$\|C_\varphi(f_{n_k}) - C_\varphi(f)\|_{F_\alpha} \rightarrow 0 \quad \text{as } k \rightarrow \infty, \quad (4.8)$$

which completes the proof. $\square$

PROPOSITION 4.2. *If $C_\varphi \in C(F_\alpha, F_\alpha)$, then $C_\varphi \in K(F_\alpha, F_\beta)$ for all $\beta > \alpha > 0$.*

PROOF. Let (f_n) be a bounded sequence in the closed unit ball of F_α . Then, $(f_n \circ \varphi)$ is bounded in F_α and since the inclusion map $i : F_\alpha \rightarrow F_\beta$ is compact, $(f_n \circ \varphi)$ has a convergent subsequence in F_β . $\square$

PROPOSITION 4.3. *$C_\varphi(f) = (f \circ \varphi)$ is compact on F_α if and only if the operator $\varphi' C_\varphi(g) = \varphi'(g \circ \varphi)$ is compact on $F_{\alpha+1}$.*

PROOF. Suppose that $C_\varphi(f) = (f \circ \varphi)$ is compact on F_α . It is known from [4] that $\varphi' C_\varphi(g) = \varphi'(g \circ \varphi)$ is bounded on $F_{\alpha+1}$. Let (g_n) be a bounded sequence in $F_{\alpha+1}$, with $g_n \rightarrow 0$ uniformly on compact subsets of $\mathbf{D}$ as $n \rightarrow \infty$. We want to show that $\lim_{n \rightarrow \infty} \|\varphi'(g_n \circ \varphi)\|_{F_{\alpha+1}} = 0$. Let (f_n) be the sequence defined by $f_n(z) = \int_0^z g_n(w) dw$. Then, $f_n \in F_\alpha$ and $\|f_n\|_{F_\alpha} \leq (2/\alpha) \|g_n\|_{F_{\alpha+1}}$, thus (f_n) is a bounded sequence in F_α . Furthermore, using the Lebesgue dominated convergence theorem we get that $f_n \rightarrow 0$ uniformly on compact subsets of $\mathbf{D}$. Thus,

$$\begin{aligned} \|\varphi'(g_n \circ \varphi)\|_{F_{\alpha+1}} &= \|\varphi'(f_n' \circ \varphi)\|_{F_{\alpha+1}} \\ &= \|(f_n \circ \varphi)'\|_{F_{\alpha+1}} \\ &\leq \alpha \|(f_n \circ \varphi)\|_{F_\alpha} \rightarrow 0 \quad \text{as } n \rightarrow \infty, \end{aligned} \quad (4.9)$$

which shows that $\varphi' C_\varphi(g) = \varphi'(g \circ \varphi)$ is compact on $F_{\alpha+1}$.

Conversely, assume that $\varphi' C_\varphi(g) = \varphi'(g \circ \varphi)$ is compact on $F_{\alpha+1}$. Then, in particular, $\varphi' C_\varphi(f') = \varphi'(f' \circ \varphi) = (f \circ \varphi)'$ is a compact for every $f \in F_\alpha$. Now, since $\|(f \circ \varphi)\|_{F_\alpha} \leq (2/\alpha) \|(f \circ \varphi)'\|_{F_{\alpha+1}}$. Let (f_n) be a bounded sequence in F_α with $f_n \rightarrow 0$ uniformly on compact subsets of $\mathbf{D}$ as $n \rightarrow \infty$. We want to show that $\lim_{n \rightarrow \infty} \|(f_n \circ \varphi)\|_{F_\alpha} = 0$. Since any bounded sequence of F_α is also a bounded sequence of $F_{\alpha+1}$, then $\|(f_n \circ \varphi)\|_{F_\alpha} \leq (2/\alpha) \|(f_n \circ \varphi)'\|_{F_{\alpha+1}} \rightarrow 0$ as $n \rightarrow \infty$ and the proof is complete. $\square$

ACKNOWLEDGMENT. The authors would like to thank the referee for his many helpful suggestions and comments.

REFERENCES

- [1] J. A. Cima and A. Matheson, *Cauchy transforms and composition operators*, Illinois J. Math. **42** (1998), no. 1, 58–69.
- [2] D. J. Hallenbeck, T. H. MacGregor, and K. Samotij, *Fractional Cauchy transforms, inner functions and multipliers*, Proc. London Math. Soc. (3) **72** (1996), no. 1, 157–187.
- [3] D. J. Hallenbeck and K. Samotij, *On Cauchy integrals of logarithmic potentials and their multipliers*, J. Math. Anal. Appl. **174** (1993), no. 2, 614–634.
- [4] R. A. Hirschweiler, *Composition operators on spaces of Cauchy transforms*, Studies on Composition Operators (Laramie, Wyo, 1996), Contemp. Math., vol. 213, American Mathematical Society, Rhode Island, 1998, pp. 57–63.
- [5] R. A. Hirschweiler and T. H. MacGregor, *Closure properties of families of Cauchy-Stieltjes transforms*, Proc. Amer. Math. Soc. **105** (1989), no. 3, 615–621.
- [6] R. A. Hirschweiler and E. Nordgren, *Cauchy transforms of measures and weighted shift operators on the disc algebra*, Rocky Mountain J. Math. **26** (1996), no. 2, 627–654.
- [7] T. H. MacGregor, *Analytic and univalent functions with integral representations involving complex measures*, Indiana Univ. Math. J. **36** (1987), no. 1, 109–130.

Yusuf Abu Muhanna: Department of Mathematics, American University of Sharjah, P.O. Box 26666, Sharjah, United Arab Emirates

E-mail address: ymuhanna@aus.ac.ae

El-Bachir Yallaoui: College of Arts and Sciences, Abu Dhabi University, P.O. Box 1790, Al Ain, United Arab Emirates

E-mail address: elbachir.yallaoui@adu.ae