## ISHIKAWA ITERATIVE SEQUENCE WITH ERRORS FOR STRONGLY PSEUDOCONTRACTIVE OPERATORS IN ARBITRARY BANACH SPACES

## **YONGJIN LI**

Received 25 May 2003

The Ishikawa iterative sequences with errors are studied for Lipschitzian strongly pseudocontractive operators in arbitrary real Banach spaces; some well-known results of Chidume (1998) and Zeng (2001) are generalized.

2000 Mathematics Subject Classification: 47H10.

**1. Introduction.** Let *E* be an arbitrary real Banach space with norm  $\|\cdot\|$  and let  $E^*$  be the dual space of *E*. The duality mapping  $J: E \to 2^{E^*}$  is defined by

$$Jx = \{ f \in E^* : \langle x, f \rangle = \|x\| \cdot \|f\|, \|f\| = \|x\| \},$$
(1.1)

where  $\langle x, f \rangle$  denotes the value of the continuous linear function  $f \in E^*$  at  $x \in E$ . It is well known that if  $E^*$  is strictly convex, then *J* is single valued.

An operator  $T: D(T) \subset E \rightarrow E$  is said to be *accretive* if the inequality

$$\|x - y\| \le \|x - y + s(Tx - Ty)\|$$
(1.2)

holds for every  $x, y \in D(T)$  and for all s > 0.

An operator *T* with domain D(T) and range R(T) in *E* is said to be a *strong pseudo-contraction* if there exists t > 1 such that for all  $x, y \in D(T)$  and r > 0, the following inequality holds:

$$\|x - y\| \le \|(1 + r)(x - y) - rt(Tx - Ty)\|.$$
(1.3)

If t = 1 in inequality (1.3), then *T* is called *pseudocontractive*.

As a consequence of the result of Kato [3], *T* is pseudocontractive if and only if for each  $x, y \in D(T)$ , there exists  $j(x - y) \in J(x - y)$  such that

$$\left\langle (I-T)x - (I-T)y, j(x-y) \right\rangle \ge 0. \tag{1.4}$$

Furthermore, *T* is strongly pseudocontractive if and only if there exists k > 0 such that

$$\langle (I-T)x - (I-T)y, j(x-y) \rangle \ge k ||x-y||^2.$$
 (1.5)

Chidume [2] proved that if *E* is a real uniformly smooth Banach space, *K* is a nonempty closed convex bounded subset of *E*, and  $T: K \to K$  is a strongly pseudocontraction with a fixed point  $x^*$  in *K*, then both the Mann and Ishikawa iteration schemes converge strongly to  $x^*$  for an arbitrary initial point  $x_0 \in K$ . Zeng [6] and Li and Liu [4] consider an iterative process for Lipschitzian strongly pseudocontractive operator in arbitrary real Banach spaces. In [2], Chidume proved the following theorem.

**THEOREM 1.1** [2]. Suppose *E* is a real uniformly smooth Banach space and *K* is a bounded closed convex and nonempty subset of *E*. Suppose  $T: E \to E$  is a strongly pseudocontractive map such that  $Tx^* = x^*$  for some  $x^* \in K$ . Let  $\{\alpha_n\}, \{\beta_n\}$  be real satisfying the following conditions:

- (i)  $0 \le \alpha_n, \beta_n \le 1$  for all  $n \ge 0$ ;
- (ii)  $\lim_{n\to\infty} \alpha_n = 0$ ,  $\lim_{n\to\infty} \beta_n = 0$ ;
- (iii)  $\sum_{n=1}^{\infty} \alpha_n = \infty$ .

Then, for arbitrary  $x_0 \in K$ , the sequence  $\{x_n\}$  defined iteratively by

$$x_{n+1} = (1 - \alpha_n) x_n + \alpha_n T y_n,$$
  

$$y_n = (1 - \beta_n) x_n + \beta_n T x_n, \quad n \ge 0,$$
(1.6)

converges strongly to  $x^*$ ; moreover,  $x^*$  is unique.

Our objective in this note is to consider an iterative sequence with errors for Lipschitzian strongly pseudocontractive operators in arbitrary real Banach spaces. Our results improve and extend the results of Chidume [2] and Zeng [6].

The following lemmas play an important role in proving our main results.

**LEMMA 1.2** [5]. Let  $\{a_n\}, \{b_n\}, \{c_n\}$  be a nonnegative sequence satisfying

$$a_{n+1} \le (1-t_n)a_n + b_n + c_n. \tag{1.7}$$

With  $\{t_n : n = 0, 1, 2, ...\} \subset [0, 1]$ ,  $\sum_{n=1}^{\infty} t_n = \infty$ ,  $b_n = o(t_n)$ , and  $\sum_{n=1}^{\infty} c_n < \infty$ , then  $\lim_{n \to \infty} a_n = 0$ .

2. Main results. Now, we state and prove the following theorems.

**THEOREM 2.1.** Suppose *E* is an arbitrary real Banach space and  $T : E \to E$  is a Lipschitzian strongly pseudocontractive map such that  $Tx^* = x^*$  for some  $x^* \in E$ . Suppose  $\{u_n\}, \{v_n\}$  are sequences in *E* and  $\{\alpha_n\}, \{\beta_n\}$  are sequences in [0,1] such that

(1) 
$$\sum_{n=1}^{\infty} \|u_n\| < \infty$$
,  $\sum_{n=1}^{\infty} \|v_n\| < \infty$ ;

- (2)  $\sum_{n=1}^{\infty} \alpha_n = \infty$ ,  $\alpha_n \to 0$  as  $n \to \infty$ ;
- (3)  $\beta_n \to 0 \text{ as } n \to \infty$ .

Then for any  $x_0 \in E$ , the Ishikawa iteration sequence  $\{x_n\}$  with errors defined by

$$y_n = (1 - \beta_n) x_n + \beta_n T x_n + v_n, x_{n+1} = (1 - \alpha_n) x_n + \alpha_n T y_n + u_n,$$
(2.1)

converges strongly to  $x^*$ ; moreover,  $x^*$  is unique.

**PROOF.** Since  $T: E \rightarrow E$  is strongly pseudocontractive, we have that (I - T) is strongly accretive, so for any  $x, y \in E$ , (1.5) holds, where k = (t-1)/t and  $t \in (1, \infty)$ . Thus

$$\langle ((1-k)I-T)x - ((1-k)I-T)y, j(x-y) \rangle \ge 0$$
 (2.2)

and so it follows from [3, Lemma 1.1] that

$$\|x - y\| \le \|x - y + r[((1 - k)I - T)x - ((1 - k)I - T)y]\|$$
(2.3)

for all  $x, y \in E$  and r > 0.

From  $x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n + u_n$ , we obtain

$$x_{n} = x_{n+1} + \alpha_{n} x_{n} - \alpha_{n} T y_{n} - u_{n}$$
  
=  $(1 + \alpha_{n}) x_{n+1} + \alpha_{n} [(I - T) x_{n+1} - k x_{n+1}] - (1 - k) \alpha_{n} x_{n}$   
+  $(2 - k) \alpha_{n}^{2} (x_{n} - T y_{n}) + \alpha_{n} (T x_{n+1} - T y_{n}) - [(2 - k) \alpha_{n} + 1] u_{n}.$  (2.4)

It is easy to see that

$$x^* = (1 + \alpha_n)x^* + \alpha_n [((1 - k)I - T)x^*] - (1 - k)\alpha_n x^*$$
(2.5)

so that

$$x_{n} - x^{*} = (1 + \alpha_{n})(x_{n+1} - x^{*}) + \alpha_{n}[((1 - k)I - T)x_{n+1} - ((1 - k)I - T)x^{*}]$$
  
- (1 - k)\alpha\_{n}(x\_{n} - x^{\*}) + (2 - k)\alpha\_{n}^{2}(x\_{n} - Ty\_{n}) + \alpha\_{n}(Tx\_{n+1} - Ty\_{n}) (2.6)  
- [(2 - k)\alpha\_{n} + 1]u\_{n}.

Hence

$$\begin{aligned} ||x_{n} - x^{*}|| &\geq (1 + \alpha_{n}) \left\| (x_{n+1} - x^{*}) + \frac{\alpha_{n}}{1 + \alpha_{n}} [((1 - k)I - T)x_{n+1} - ((1 - k)I - T)x^{*}] \right\| \\ &- (1 - k)\alpha_{n} ||x_{n} - x^{*}|| - (2 - k)\alpha_{n}^{2} ||x_{n} - Ty_{n}|| \\ &- \alpha_{n} ||Tx_{n+1} - Ty_{n}|| - [(2 - k)\alpha_{n} + 1]||u_{n}|| \\ &\geq (1 + \alpha_{n}) ||(x_{n+1} - x^{*})|| - (1 - k)\alpha_{n} ||x_{n} - x^{*}|| - (2 - k)\alpha_{n}^{2} ||x_{n} - Ty_{n}|| \\ &- \alpha_{n} ||Tx_{n+1} - Ty_{n}|| - [(2 - k)\alpha_{n} + 1]||u_{n}|| \end{aligned}$$

$$(2.7)$$

so that

$$||x_{n+1} - x^*|| \leq \left[\frac{1 + (1 - k)\alpha_n}{1 + \alpha_n}\right] ||x_n - x^*|| + \alpha_n^2 ||x_n - Ty_n|| + \alpha_n ||Tx_{n+1} - Ty_n|| + (2\alpha_n + 1)||u_n|| \leq \left(1 - \frac{k\alpha_n}{1 + \alpha_n}\right) ||x_n - x^*|| + \alpha_n^2 ||x_n - Ty_n|| + \alpha_n ||Tx_{n+1} - Ty_n|| + (2\alpha_n + 1)||u_n|| \leq \left(1 - \frac{k\alpha_n}{2}\right) ||x_n - x^*|| + \alpha_n^2 ||x_n - Ty_n|| + \alpha_n ||Tx_{n+1} - Ty_n|| + 3||u_n||.$$
(2.8)

Since T is a Lipschitzian operator and L is Lipschitz bound, we have

$$\begin{aligned} ||y_{n} - x^{*}|| &= ||(1 - \beta_{n})(x_{n} - x^{*}) + \beta_{n}(Tx_{n} - x^{*}) + v_{n}|| \\ &\leq [1 + \beta_{n}(L - 1)]||x_{n} - x^{*}|| + ||v_{n}|| \leq L||x_{n} - x^{*}|| + ||v_{n}||, \\ ||x_{n} - Ty_{n}|| &\leq ||x_{n} - x^{*}|| + L||y_{n} - x^{*}|| \leq (1 + L^{2})||x_{n} - x^{*}|| + L||v_{n}||, \\ ||Tx_{n+1} - Ty_{n}|| \leq L||(1 - \alpha_{n})(x_{n} - y_{n}) + \alpha_{n}(Ty_{n} - y_{n}) + u_{n}|| \\ &\leq L(1 - \alpha_{n})[\beta_{n}(1 + L)||x_{n} - x^{*}|| + ||v_{n}||] \\ &+ L\alpha_{n}(1 + L)[L||x_{n} - x^{*}|| + ||v_{n}||] + L||u_{n}|| \\ &\leq [L(1 + L)\beta_{n} + (1 + L)L^{2}\alpha_{n}]||x_{n} - x^{*}|| + L(1 + L)||v_{n}|| + L||u_{n}||. \end{aligned}$$

So there exist  $M_1 > 0$  and  $M_2 > 0$  such that

$$||x_{n+1} - x^*|| \le \left(1 - \frac{k\alpha_n}{2}\right)||x_n - x^*|| + [L(1+L)\beta_n + (L^3 + 3L^2 + 2)\alpha_n]\alpha_n||x_n - x^*|| + M_1||u_n|| + M_2||v_n||.$$
(2.10)

Since  $\alpha_n \to 0$  and  $\beta_n \to 0$ , there exists N > 0 such that for all n > N, we have

$$L(1+L)\beta_n + (L^3 + 3L^2 + 2)\alpha_n < \frac{k}{4}.$$
(2.11)

Thus

$$||x_{n+1} - x^*|| \le \left(1 - \frac{k\alpha_n}{4}\right)||x_n - x^*|| + M_1||u_n|| + M_2||v_n||.$$
(2.12)

Set

$$t_n = \frac{k\alpha_n}{4}, \qquad b_n = 0, \qquad c_n = M_1 ||u_n|| + M_2 ||v_n||.$$
 (2.13)

Then we have

$$a_{n+1} \le (1 - t_n)a_n + b_n + c_n. \tag{2.14}$$

According to the above argument, it is easily seen that

$$\sum_{k=0}^{\infty} t_n = \infty, \qquad b_n = o(t_n), \qquad \sum_{k=0}^{\infty} c_n < \infty$$
(2.15)

and so, by Lemma 1.2, we have  $\lim a_n = \lim ||x_n - x^*|| = 0$ . Uniqueness follows as in [1]. The proof of the theorem is complete.

**REMARK 2.2.** Our Theorem 2.1 generalized the theorem of Chidume [2] from uniformly smooth Banach space to arbitrary Banach space and from Ishikawa iteration to Ishikawa iteration with errors. In addition, our results extend, generalize, and improve the corresponding results obtained by Zeng [6] and Li and Liu [4].

**ACKNOWLEDGMENTS.** This work was supported by the Foundation of Sun Yat-sen University Advanced Research Centre and Lingnan Foundation. The author is grateful to the referees for careful reading of the manuscript, helpful comments, and valuable suggestions.

## REFERENCES

- C. E. Chidume, Iterative approximation of fixed points of Lipschitzian strictly pseudocontractive mappings, Proc. Amer. Math. Soc. 99 (1987), no. 2, 283-288.
- [2] \_\_\_\_\_, Global iteration schemes for strongly pseudo-contractive maps, Proc. Amer. Math. Soc. 126 (1998), no. 9, 2641–2649.
- [3] T. Kato, Nonlinear semigroups and evolution equations, J. Math. Soc. Japan 19 (1967), 508– 520.
- Y. Q. Li and L. W. Liu, *Iterative processes for Lipschitz strongly accretive operators*, Acta Math. Sinica 41 (1998), no. 4, 845–850.
- [5] L. S. Liu, Ishikawa and Mann iterative process with errors for nonlinear strongly accretive mappings in Banach spaces, J. Math. Anal. Appl. 194 (1995), no. 1, 114-125.
- [6] L. C. Zeng, Ishikawa type iterative sequences with errors for Lipschitzian strongly pseudocontractive mappings in Banach spaces, Chinese Ann. Math. Ser. A 22 (2001), no. 5, 639–644.

Yongjin Li: Department of Mathematics, Sun Yat-sen University, Guangzhou 510275, China *E-mail address*: stslyj@zsu.edu.cn



Advances in **Operations Research** 



**The Scientific** World Journal







Hindawi

Submit your manuscripts at http://www.hindawi.com



Algebra



Journal of Probability and Statistics



International Journal of Differential Equations





Complex Analysis





Mathematical Problems in Engineering



Abstract and Applied Analysis



Discrete Dynamics in Nature and Society



International Journal of Mathematics and Mathematical Sciences





Journal of **Function Spaces** 



International Journal of Stochastic Analysis

