METHOD OF SEMIDISCRETIZATION IN TIME TO NONLINEAR RETARDED DIFFERENTIAL EQUATIONS WITH NONLOCAL HISTORY CONDITIONS

S. AGARWAL and D. BAHUGUNA

Received 4 January 2004

This paper deals with the applications of the method of semidiscretization in time to a nonlinear retarded differential equation with a nonlocal history condition. We establish the existence and uniqueness of a strong solution. Finally, we consider some applications of the abstract results.

2000 Mathematics Subject Classification: 34K30, 34G20, 47H06.

1. Introduction. Consider the following nonlinear retarded differential equation in a real Hilbert space H :

$$\begin{aligned} u'(t) + Au(t) &= f(t, u(t), u(r_1(t)), u(r_2(t)), \dots, u(r_m(t))), \quad t \in (0, T], \\ h(u_{[-\tau, 0]}) &= \phi_0, \quad \text{on } [-\tau, 0], \end{aligned} \quad (1.1)$$

where $0 < \tau, T < \infty$, $\phi_0 \in \mathcal{C}_0 := C([-\tau, 0]; H)$, the nonlinear operator A is single valued and maximal monotone defined from the domain $D(A) \subset H$ into H , the nonlinear map f is defined from $[0, T] \times H^{m+1}$ into H , the map h is defined from $\mathcal{C}_0$ into $\mathcal{C}_0$, and $\psi_{[-\tau, 0]}$ is the restriction of $\psi \in \mathcal{C}_T := C([-\tau, T]; H)$ on $[-\tau, 0]$. Here $\mathcal{C}_t := C([-\tau, t]; H)$ for $t \in [0, T]$ is the Banach space of all continuous functions from $[-\tau, t]$ into H endowed with the supremum norm

$$\|\phi\|_t := \sup_{-\tau \leq \eta \leq t} \|\phi(\eta)\|, \quad \phi \in \mathcal{C}_t, \quad (1.2)$$

where $\|\cdot\|$ represents the norm in H and the functions $r_i : [0, T] \rightarrow [-\tau, T]$, $i = 1, 2, \dots, m$.

The existence and uniqueness results for (1.1) may also be applied to the particular case, namely, the retarded functional differential equation

$$\begin{aligned} u'(t) + Au(t) &= f(t, u(t), u(t - \tau_1), u(t - \tau_2), \dots, u(t - \tau_m)), \quad t \in (0, T], \\ u &= \phi_0, \quad \text{on } [-\tau, 0], \end{aligned} \quad (1.3)$$

where $0 < \tau_i < T$ and $\tau = \max\{\tau_1, \tau_2, \dots, \tau_m\}$.

For the earlier works on existence, uniqueness, and stability of various types of solutions of differential and functional differential equations with nonlocal conditions, we refer to Byszewski and Lakshmikantham [7], Byszewski [6], Balachandran and Chandrasekaran [4], Lin and Liu [11], and the references cited in these papers.

Our aim is to extend the application of the method of semidiscretization in time, also known as the method of lines, to (1.1). For the applications of the method of lines to nonlinear evolution and nonlinear functional evolution equations, we refer to Kartsatos and Parrott [9, 10], Kartsatos [8], Bahuguna and Raghavendra [3], Bahuguna [1], and the references cited in these papers.

Suppose that there is $\chi \in \mathcal{C}_T$ such that $h(\chi_{[-\tau,0]}) = \phi_0$ on $[-\tau, 0]$ and $\chi(0) \in D(A)$. We prove the existence of a *strong solution* u of (1.1) under the assumptions of Theorem 2.4, stated in the next section, in the sense that there exists a unique function $u \in \mathcal{C}_T$ such that $u(t) \in D(A)$ for a.e. $t \in [0, T]$, u is differentiable a.e. on $[0, T]$, and

$$\begin{aligned} u'(t) + Au(t) &= f(t, u(t), u(r_1(t)), \dots, u(r_m(t))), \quad \text{a.e. } t \in [0, T], \\ u_{[-\tau,0]} &= \chi_{[-\tau,0]}, \quad \text{on } [-\tau, 0]. \end{aligned} \quad (1.4)$$

Finally, we show that u is unique if and only if $\chi \in \mathcal{C}_T$ satisfying $h(\chi_{[-\tau,0]}) = \phi_0$ is unique up to $[-\tau, 0]$. We also consider some applications of the abstract results.

2. Preliminaries and main result. Let H be a real Hilbert space. Let (x, y) be the inner product of $x, y \in H$. We assume the following conditions.

(A1) The operator $A : D(A) \subset H \rightarrow H$ is maximal monotone, that is,

$$(Ax - Ay, x - y) \geq 0, \quad \forall x, y \in D(A), \quad R(I + A) = H, \quad (2.1)$$

where $R(\cdot)$ is the range of an operator.

(A2) The map $h : \mathcal{C}_0 \rightarrow \mathcal{C}_0$ and there exists $\chi \in \mathcal{C}_T$ such that $h(\chi_{[-\tau,0]}) = \phi_0$ and $\chi(0) \in D(A)$.

(A3) The nonlinear map $f : [0, T] \times H^{m+1} \rightarrow H$ satisfies a local Lipschitz-like condition

$$\|f(t, u_1, u_2, \dots, u_{m+1}) - f(s, v_1, v_2, \dots, v_{m+1})\| \leq L_f(r) \left[|t - s| + \sum_{i=1}^{m+1} \|u_i - v_i\| \right], \quad (2.2)$$

for all $(u_1, u_2, \dots, u_{m+1})$ and $(v_1, v_2, \dots, v_{m+1})$ in $B_r(H^{m+1}, (\chi(0), \dots, \chi(0)))$ and $t \in [0, T]$, where $L_f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a nondecreasing function and, for $r > 0$,

$$B_r(H^{m+1}, (\chi(0), \dots, \chi(0))) = \left\{ (u_1, \dots, u_{m+1}) \in H^{m+1} : \sum_{i=1}^{m+1} \|u_i - \chi(0)\| \leq r \right\}. \quad (2.3)$$

(A4) For $i = 1, 2, \dots, m$, the maps $r_i : [0, T] \rightarrow [-\tau, T]$ are continuous satisfying the delay property $r_i(t) \leq t$ for $t \in [0, T]$.

Further, if A is monotone, $1 + \alpha A$ is invertible for $\alpha > 0$ and the inverse operator $(1 + \alpha A)^{-1}$ is Lipschitz continuous, that is,

$$\|(1 + \alpha A)^{-1}u - (1 + \alpha A)^{-1}v\| \leq \|u - v\|, \quad u, v \in D((1 + \alpha A)^{-1}). \quad (2.4)$$

Also, the inverse of an invertible monotonic operator in Hilbert space H is monotonic, but it might not be true in the general case.

For an m -monotonic operator A , we introduce the following sequences of operator ($n = 1, 2, \dots$):

$$\begin{aligned} J_n &= (1 + n^{-1}A)^{-1}, \\ A_n &= AJ_n = n(I - J_n), \end{aligned} \quad (2.5)$$

where AJ_n denotes the composition of the two maps A and J_n . The J_n and A_n are defined everywhere on H . We recall simple lemmas that we will have occasions to use. More details may be found in [5].

LEMMA 2.1. *Let A be m -monotonic. J_n and A_n are uniformly Lipschitz continuous, with*

$$\begin{aligned} \|J_n u - J_n v\| &\leq \|u - v\|, \\ \|A_n u - A_n v\| &\leq 2n\|u - v\|. \end{aligned} \quad (2.6)$$

LEMMA 2.2. *Let A be m -monotonic. Then A_n are also monotonic. Furthermore,*

$$\|A_n u\| \leq \|Au\|, \quad \text{for } u \in D(A). \quad (2.7)$$

LEMMA 2.3. *Let A be m -monotonic in H .*

- (a) *If $X^n \in D(A)$, $n = 1, 2, \dots$, $X^n \rightarrow u \in H$, and if $\|AX^n\|$ are bounded, then $u \in D(A)$ and $AX_n \rightarrow Au$.*
- (b) *If $x_n \in H$, $n = 1, 2, \dots$, $x_n \rightarrow u \in H$, and if $\|A_n x_n\|$ are bounded, then $u \in D(A)$ and $A_n x_n \rightarrow Au$.*

PROOF. (a) The monotonicity condition gives

$$(Av - AX^n, v - X^n) \geq 0, \quad (2.8)$$

for any $v \in D(A)$. Since H is reflexive and the $\|AX^n\|$ are bounded, there is a subsequence $\{X^{n'}\}$ of $\{X^n\}$ such that $AX^{n'} \rightarrow w \in H$. Since $v - X^{n'} \rightarrow v - u$, we obtain from (2.8) the inequality $(Av - w, v - u) \geq 0$. For fixed $z \in H$, $t > 0$, put $v_t = A_1^{-1}(u + w + tz)$, where $A_1 = (I + A)$. It follows that $Av_t + v_t - u - w = tz$, hence $t(z, v_t - u) \geq 0$ so $(z, v_t - u) \geq 0$. Letting $t \rightarrow 0$, $v_t \rightarrow A_1^{-1}(u + w)$, hence $(z, A_1^{-1}(u + w) - u) \geq 0$, so $u = A_1^{-1}(u + w)$, $u \in D(A)$, and $Au = w$. It shows that $AX^n \rightarrow w = Au$.

(b) Set $X^n = J_n x_n \in D(A)$. Then $AX^n = (AJ_n x_n) = A_n x_n$ and $\|AX^n\|$ are bounded. Also, $x_n - X^n = (I - J_n)x_n = n^{-1}A_n x_n \rightarrow 0$, so that $X^n \rightarrow u$. Thus, by the result of (a), we get $u \in D(A)$ and $A_n x_n = AX^n \rightarrow Au$. $\square$

Now, we state the following main result of this paper which will be proved in the next section after proving some a priori estimates.

THEOREM 2.4. *Suppose that the conditions (A1), (A2), (A3), (A4) are satisfied. Then (1.1) has a strong solution $u \in \mathcal{C}_T$ either on $[-\tau, T]$ or on the maximal interval of existence $[-\tau, t_{\max})$, $0 < t_{\max} \leq T$, and in the later case, either $\lim_{t \rightarrow t_{\max}^-} \|u(t)\| = \infty$ or $u(t)$ goes to the boundary of $D(A)$ as $t \rightarrow t_{\max}^-$. Moreover, u is Lipschitz continuous on every compact subinterval of existence.*

3. Discretization scheme and a priori estimates. In this section, we establish the existence and uniqueness of a strong solution to (1.4) for any given $\chi \in \mathcal{C}_T$ with $\chi(0) \in D(A)$. For the application of the method of lines to (1.4), we proceed as follows. Let $R_0 := \sup_{t \in [-\tau, T]} \|\chi(t) - \chi(0)\|$. For any $0 < R \leq R_0$, we choose t_0 such that

$$0 < t_0 \leq T, \quad t_0 [\|A\chi(0)\| + 3L_f(R_0)(T + (m+1)R_0) + \|f(0, \chi(0), \chi(0), \dots, \chi(0))\|] \leq R. \quad (3.1)$$

For $n \in \mathbb{N}$, let $h_n = t_0/n$. We set $u_0^n = \chi(0)$ for all $n \in \mathbb{N}$ and define each of $\{u_j^n\}_{j=1}^n$ as the unique solution of the equation

$$\frac{u - u_{j-1}^n}{h_n} + Au = f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n)), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))), \quad (3.2)$$

where $\tilde{u}_0^n(t) = \chi(t)$ for $t \in [-\tau, 0]$, $\tilde{u}_0^n(t) = \chi(0)$ for $t \in [0, t_0]$ and for $2 \leq j \leq n$,

$$\tilde{u}_{j-1}^n(t) = \begin{cases} \chi(t), & t \in [-\tau, 0], \\ u_{i-1}^n + \frac{1}{h_n}(t - t_{i-1}^n)(u_i^n - u_{i-1}^n), & t \in [t_{i-1}^n, t_i^n], \quad i = 1, 2, \dots, j-1, \\ u_{j-1}^n, & t \in [t_{j-1}^n, t_0]. \end{cases} \quad (3.3)$$

The existence of a unique $u_j^n \in D(A)$ satisfying (3.2) is a consequence of the m -monotonicity of A . Using (A2), we first prove that the points $\{u_j^n\}_{j=0}^n$ lie in a ball with its radius independent of the discretization parameters j , h_n , and n . We then prove a priori estimates on the difference quotients $\{u_j^n - u_{j-1}^n/h_n\}$ using (A2). We define the sequence $\{U^n\} \subset \mathcal{C}_T$ of polygonal functions

$$U^n(t) = \begin{cases} \chi(t), & t \in [-\tau, 0], \\ u_{j-1}^n + \frac{1}{h_n}(t - t_{j-1}^n)(u_j^n - u_{j-1}^n), & t \in (t_{j-1}^n, t_j^n], \end{cases} \quad (3.4)$$

and prove the convergence of $\{U^n\}$ to a unique strong solution u of (1.4) in $\mathcal{C}_T$ as $n \rightarrow \infty$.

Now, we first show that $\{u_j^n\}_{j=0}^n$ lie in a ball in H of radius independent of j , h_n , and n .

LEMMA 3.1. For $n \in \mathbb{N}$, $j = 1, 2, \dots, n$,

$$\|u_j^n - \chi(0)\| \leq R. \quad (3.5)$$

PROOF. From (3.2) for $j = 1$, we have

$$\frac{u_1^n - u_0^n}{h_n} + Au_1^n = f(t_1^n, u_0^n, \tilde{u}_0^n(r_1(t_1^n)), \dots, \tilde{u}_0^n(r_m(t_1^n))). \quad (3.6)$$

Subtracting Au_0^n on both sides, we get

$$\frac{u_1^n - u_0^n}{h_n} + Au_1^n - Au_0^n = -Au_0^n + f(t_1^n, u_0^n, \tilde{u}_0^n(r_1(t_1^n)), \dots, \tilde{u}_0^n(r_m(t_1^n))). \quad (3.7)$$

Taking the inner product with $u_1^n - u_0^n$, we get

$$\begin{aligned} & \left(\frac{u_1^n - u_0^n}{h_n}, u_1^n - u_0^n \right) + (Au_1^n - Au_0^n, u_1^n - u_0^n) \\ & = -(Au_0^n, u_1^n - u_0^n) + (f(t_1^n, u_0^n, \tilde{u}_0^n(r_1(t_1^n))), \dots, \tilde{u}_0^n(r_m(t_1^n))), u_1^n - u_0^n). \end{aligned} \quad (3.8)$$

By the monotonicity of A , we have

$$\|u_1^n - u_0^n\| \leq h_n [\|Au_0^n\| + \|f(t_1^n, u_0^n, \tilde{u}_0^n(r_1(t_1^n))), \dots, \tilde{u}_0^n(r_m(t_1^n)))\|]. \quad (3.9)$$

Using (A3), we get

$$\|u_1^n - \chi(0)\| \leq h_n [\|A\chi(0)\| + 3L_f(R_0)(T + (m+1)R_0) + \|f(0, \chi(0), \dots, \chi(0))\|] \leq R. \quad (3.10)$$

Assume that $\|u_i^n - \chi(0)\| \leq R$ for $i = 1, 2, \dots, j-1$.

Now, for $2 \leq j \leq n$,

$$\frac{u_j^n - u_{j-1}^n}{h_n} + Au_j^n = f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n))), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))). \quad (3.11)$$

Subtracting $A\chi(0)$ on both sides, we get

$$\frac{u_j^n - u_{j-1}^n}{h_n} + Au_j^n - A\chi(0) = -A\chi(0) + f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n))), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))). \quad (3.12)$$

Taking the inner product with $u_j^n - \chi(0)$, we get

$$\begin{aligned} & \frac{1}{h_n} (u_j^n - u_{j-1}^n, u_j^n - \chi(0)) + (Au_j^n - A\chi(0), u_j^n - \chi(0)) \\ & = -(A\chi(0), u_j^n - \chi(0)) + (f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n))), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))), u_j^n - \chi(0)). \end{aligned} \quad (3.13)$$

By the monotonicity of A , we have

$$\begin{aligned} \|u_j^n - \chi(0)\| & \leq \|u_{j-1}^n - \chi(0)\| \\ & + h_n [\|A\chi(0)\| + \|f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n))), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n)))\|]. \end{aligned} \quad (3.14)$$

Using (A3), we get

$$\begin{aligned} \|u_j^n - \chi(0)\| & \leq \|u_{j-1}^n - \chi(0)\| \\ & + h_n [\|A\chi(0)\| + 3L_f(R_0)(T + (m+1)R_0) + \|f(0, \chi(0), \chi(0), \dots, \chi(0))\|]. \end{aligned} \quad (3.15)$$

Repeating the above inequality, we obtain

$$\begin{aligned} \|u_j^n - \chi(0)\| &\leq jh_n[\|A\chi(0)\| + 3L_f(R_0)(T + (m+1)R_0) \\ &\quad + \|f(0, \chi(0), \chi(0), \dots, \chi(0))\|] \leq R, \end{aligned} \quad (3.16)$$

as $jh_n \leq t_0$ for $0 \leq j \leq n$. This completes the proof of the lemma. $\square$

Now, we establish a priori estimates for the difference quotients $\{(u_j^n - u_{j-1}^n)/h_n\}$.

LEMMA 3.2. *There exists a positive constant K independent of the discretization parameters n, j , and h_n such that*

$$\left\| \frac{u_j^n - u_{j-1}^n}{h_n} \right\| \leq K, \quad j = 1, 2, \dots, n, \quad n = 1, 2, \dots \quad (3.17)$$

PROOF. In this proof and subsequently, K will represent a generic constant independent of j, h_n , and n . Subtracting $Au_0^n = A\chi(0)$ from both sides in (3.2), we get

$$\frac{u_1^n - u_0^n}{h_n} + Au_1^n - Au_0^n = -Au_0^n + f(t_1^n, u_0^n, \tilde{u}_0^n(r_1(t_1^n)), \dots, \tilde{u}_0^n(r_m(t_1^n))). \quad (3.18)$$

Taking the inner product with $(u_1^n - u_0^n)$, we get

$$\begin{aligned} &\left(\frac{u_1^n - u_0^n}{h_n}, u_1^n - u_0^n \right) + (Au_1^n - Au_0^n, u_1^n - u_0^n) \\ &= -(Au_0^n, u_1^n - u_0^n) + (f(t_1^n, u_0^n, \tilde{u}_0^n(r_1(t_1^n)), \dots, \tilde{u}_0^n(r_m(t_1^n))), u_1^n - u_0^n). \end{aligned} \quad (3.19)$$

Using monotonicity of A , we get

$$\left\| \frac{u_1^n - u_0^n}{h_n} \right\| \leq \|A\chi(0)\| + \|f(t_1^n, u_0^n, \tilde{u}_0^n(r_1(t_1^n)), \dots, \tilde{u}_0^n(r_m(t_1^n)))\|. \quad (3.20)$$

Using (A3), we have

$$\begin{aligned} \left\| \frac{u_1^n - u_0^n}{h_n} \right\| &\leq \|A\chi(0)\| + \|f(0, \chi(0), \chi(0), \dots, \chi(0))\| + 3L_f(R_0)(T + (m+1)R_0) \\ &\leq K. \end{aligned} \quad (3.21)$$

Now, for $2 \leq j \leq n$, taking the inner product with $(u_j^n - u_{j-1}^n)$ in (3.2), we get

$$\begin{aligned} &\left(\frac{u_j^n - u_{j-1}^n}{h_n}, u_j^n - u_{j-1}^n \right) + (Au_j^n, u_j^n - u_{j-1}^n) \\ &= (f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n)), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))), u_j^n - u_{j-1}^n). \end{aligned} \quad (3.22)$$

Also,

$$\begin{aligned} &\left(\frac{u_j^n - u_{j-1}^n}{h_n} - \frac{u_{j-1}^n - u_{j-2}^n}{h_n}, u_j^n - u_{j-1}^n \right) + (Au_j^n - Au_{j-1}^n, u_j^n - u_{j-1}^n) \\ &= (f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n)), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))) \\ &\quad - f(t_{j-1}^n, u_{j-2}^n, \tilde{u}_{j-2}^n(r_1(t_{j-1}^n)), \dots, \tilde{u}_{j-2}^n(r_m(t_{j-1}^n))), u_j^n - u_{j-1}^n). \end{aligned} \quad (3.23)$$

Using the monotonicity of A , we get

$$\begin{aligned} \left\| \frac{u_j^n - u_{j-1}^n}{h_n} \right\| &\leq \left\| \frac{u_{j-1}^n - u_{j-2}^n}{h_n} \right\| \\ &\quad + \|f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n)), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))) \\ &\quad - f(t_{j-1}^n, u_{j-2}^n, \tilde{u}_{j-2}^n(r_1(t_{j-1}^n)), \dots, \tilde{u}_{j-2}^n(r_m(t_{j-1}^n)))\|. \end{aligned} \quad (3.24)$$

Now,

$$\begin{aligned} &\|f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n)), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))) \\ &\quad - f(t_{j-1}^n, u_{j-2}^n, \tilde{u}_{j-2}^n(r_1(t_{j-1}^n)), \dots, \tilde{u}_{j-2}^n(r_m(t_{j-1}^n)))\| \\ &\leq C \left[h_n + (m+1)h_n \left\| \frac{u_j^n - u_{j-1}^n}{h_n} \right\| \right]. \end{aligned} \quad (3.25)$$

From the above inequality, we get

$$\begin{aligned} \max_{\{1 \leq k \leq j\}} \left\| \frac{u_k^n - u_{k-1}^n}{h_n} \right\| &\leq (1 + Ch_n) \max_{\{1 \leq k \leq j-1\}} \left\| \frac{u_k^n - u_{k-1}^n}{h_n} \right\| + Ch_n \\ &\leq (1 + Ch_n) \left[1 + \max_{\{1 \leq k \leq j-1\}} \left\| \frac{u_k^n - u_{k-1}^n}{h_n} \right\| \right], \end{aligned} \quad (3.26)$$

where C is a positive constant independent of j , h_n , and n . Repeating the above inequality, we get

$$\max_{\{1 \leq k \leq j\}} \left\| \frac{u_j^n - u_{j-1}^n}{h_n} \right\| \leq (1 + Ch_n)^j D \leq De^{TC} \leq K. \quad (3.27)$$

This completes the proof of the lemma. $\square$

We introduce another sequence $\{X^n\}$ of step functions from $[0, T]$ into H by

$$X^n(t) = \begin{cases} \chi(0), & t = 0, \\ u_j^n, & t \in (t_{j-1}^n, t_j^n]. \end{cases} \quad (3.28)$$

REMARK 3.3. From [Lemma 3.2](#), it follows that the functions U^n and $\tilde{u}_r^n$, $0 \leq r \leq n-1$, are Lipschitz continuous on $[0, t_0]$ with a uniform Lipschitz constant K . The sequence $U^n(t) - X^n(t) \rightarrow 0$ in H as $n \rightarrow \infty$ uniformly on $[0, t_0]$. Furthermore, $X^n(t) \in D(A)$ for $t \in [0, t_0]$ and the sequences $\{U^n(t)\}$ and $\{X^n(t)\}$ are bounded in H , uniformly in $n \in \mathbb{N}$ and $t \in [0, t_0]$. The sequence $\{AX^n(t)\}$ is bounded uniformly in $n \in \mathbb{N}$ and $t \in [0, t_0]$.

For notational convenience, let

$$f^n(t) = f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n)), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))), \quad t \in (t_{j-1}^n, t_j^n], \quad 1 \leq j \leq n. \quad (3.29)$$

Then (3.2) may be rewritten as

$$\frac{d^-}{dt}U^n(t) + AX^n(t) = f^n(t), \quad t \in (0, t_0], \quad (3.30)$$

where d^-/dt denotes the left derivative in $(0, t_0]$. Also, for $t \in (0, t_0]$, we have

$$\int_0^t AX^n(s)ds = \chi(0) - U^n(t) + \int_0^t f^n(s)ds. \quad (3.31)$$

LEMMA 3.4. *There exists $u \in \mathcal{C}_{t_0}$ such that $U^n \rightarrow u$ in $\mathcal{C}_{t_0}$ as $n \rightarrow \infty$. Moreover, u is Lipschitz continuous on $[0, t_0]$.*

PROOF. From (3.30) for $t \in (0, t_0]$, and taking the inner product with $(X^n(t) - X^k(t))$, we have

$$\begin{aligned} & \left(\frac{d^-}{dt}U^n(t) - \frac{d^-}{dt}U^k(t), X^n(t) - X^k(t) \right) \\ & \quad + (AX^n(t) - AX^k(t), X^n(t) - X^k(t)) \\ & = (f^n(t) - f^k(t), X^n(t) - X^k(t)). \end{aligned} \quad (3.32)$$

Using monotonicity of A , we have

$$\left(\frac{d^-}{dt}(U^n(t) - U^k(t)), X^n(t) - X^k(t) \right) \leq (f^n(t) - f^k(t), X^n(t) - X^k(t)). \quad (3.33)$$

From the above inequality and the fact that

$$2 \left(\frac{d^-}{dt}U^n(t) - \frac{d^-}{dt}U^k(t), U^n(t) - U^k(t) \right) = \frac{d^-}{dt} \|U^n(t) - U^k(t)\|^2, \quad (3.34)$$

we get

$$\begin{aligned} & \frac{1}{2} \frac{d^-}{dt} \|U^n(t) - U^k(t)\|^2 \\ & \leq \left(\frac{d^-}{dt}(U^n(t) - U^k(t)) + f^n(t) - f^k(t) - f^n(t) + f^k(t), \right. \\ & \quad \left. U^n(t) - U^k(t) - X^n(t) + X^k(t) \right) + (f^n(t) - f^k(t), X^n(t) - X^k(t)) \\ & = \left(\frac{d^-}{dt}(U^n(t) - U^k(t)) - f^n(t) + f^k(t), U^n(t) - U^k(t) - X^n(t) + X^k(t) \right) \\ & \quad + (f^n(t) - f^k(t), U^n(t) - U^k(t)). \end{aligned} \quad (3.35)$$

Now,

$$\begin{aligned} \|f^n(t) - f^k(t)\| &= \|f(t_j^n, u_{j-1}^n, \tilde{u}_{j-1}^n(r_1(t_j^n)), \dots, \tilde{u}_{j-1}^n(r_m(t_j^n))) \\ & \quad - f(t_l^k, u_{l-1}^k, \tilde{u}_{l-1}^k(r_1(t_l^k)), \dots, \tilde{u}_{l-1}^k(r_m(t_l^k)))\|. \end{aligned} \quad (3.36)$$

Using (A3), we get

$$\|f^n(t) - f^k(t)\| \leq \epsilon_{nk}(t) + K \|U^n - U^k\|_t, \quad (3.37)$$

where

$$\begin{aligned} \epsilon_{nk}(t) = K & \left[h_n + h_k + |t_j^n - t_l^k| + \|X^n(t - h_n) - U^n(t)\| \right. \\ & + \|X^k(t - h_k) - U^k(t)\| \\ & \left. + \sum_{i=1}^m (|r_i(t) - r_i(t_j^n)| + |r_i(t) - r_i(t_l^k)|) \right], \end{aligned} \quad (3.38)$$

for $t \in (t_{j-1}^n, t_j^n]$ and $t \in (t_{l-1}^k, t_l^k]$, $1 \leq j \leq n$, $1 \leq l \leq k$. Therefore $\epsilon_{nk}(t) \rightarrow 0$ as $n, k \rightarrow \infty$ uniformly on $[0, t_0]$. This implies that for a.e. $t \in [0, t_0]$,

$$\begin{aligned} & \frac{d^-}{dt} \|U^n(t) - U^k(t)\|^2 \\ & \leq 2 \left[\left\| \frac{d^-}{dt} (U^n(t) - U^k(t) - f^n(t) - f^k(t)) \right\| \right. \\ & \quad \times \|U^n(t) - U^k(t) - X^n(t) + X^k(t)\| \\ & \quad \left. + \|f^n(t) - f^k(t)\| \|U^n(t) - U^k(t)\| \right] \\ & \leq K [\epsilon_{nk} + \|U^n - U^k\|_t^2], \end{aligned} \quad (3.39)$$

where ϵ_{nk} is a sequence of numbers such that $\epsilon_{nk} \rightarrow 0$ as $n, k \rightarrow \infty$. Integrating the above inequality over $(0, s)$, $0 < s \leq t \leq t_0$, and using the fact that $U^n = \phi$ on $[-\tau, 0]$ for all n , we get

$$\|U^n(s) - U^k(s)\|^2 \leq K \left[T\epsilon_{nk} + \int_0^t \|U^n - U^k\|_s^2 ds \right]. \quad (3.40)$$

Taking the supremum over $[0, t]$, we get

$$\sup_{s \in [0, t]} \|U^n(s) - U^k(s)\|^2 \leq K \left[T\epsilon_{nk} + \int_0^t \|U^n - U^k\|_s^2 ds \right]. \quad (3.41)$$

Thus,

$$\|U^n - U^k\|_t^2 \leq K \left[T\epsilon_{nk} + \int_0^t \|U^n - U^k\|_s^2 ds \right]. \quad (3.42)$$

Applying Gronwall's inequality, we conclude that there exists $u \in \mathcal{C}_{t_0}$ such that $U^n \rightarrow u$ in $\mathcal{C}_{t_0}$. Clearly, $u = \phi$ on $[-\tau, 0]$ and, from [Remark 3.3](#), it follows that u is Lipschitz continuous on $[0, t_0]$. This completes the proof of the lemma. $\square$

PROOF OF THEOREM 2.4. We first prove the existence on $[-\tau, t_0]$ and then prove the unique continuation of the solution on $[-\tau, T]$. From [Lemma 2.3\(a\)](#), we know that $u(t) \in D(A)$ for $t \in [0, t_0]$, $AX^n(t) \rightharpoonup Au(t)$ on $[0, t_0]$. Here, $\rightharpoonup$ denotes the weak convergence in H . Also, we may show that $Au(t)$ is weakly continuous on $[0, t_0]$ as follows.

Let $t_i \rightarrow t$; $t_i, t \in [0, t_0]$, then $u(t_i) \rightarrow u(t)$ and, by boundedness of $A(u(t_i))$ and reflexivity of H , there exists a sequence $\{u(t_j)\}$ of $\{u(t_i)\}$ such that $Au(t_j) \rightharpoonup w(t)$. Then it follows from [Lemma 2.3\(a\)](#) that $Au(t_i) \rightharpoonup Au(t)$.

For every $x^* \in X^*$ and $t \in (0, t_0]$, we have

$$\int_0^t \langle AX^n(s), x^* \rangle ds = \langle \chi(0), x^* \rangle - \langle U^n(t), x^* \rangle + \int_0^t \langle f^n(s), x^* \rangle ds. \quad (3.43)$$

Using [Lemma 3.4](#) and the bounded convergence theorem, we obtain as $n \rightarrow \infty$,

$$\begin{aligned} \int_0^t \langle Au(s), x^* \rangle ds &= \langle \chi(0), x^* \rangle - \langle u(t), x^* \rangle \\ &\quad + \int_0^t \langle f(s, u(s), u(r_1(s)), \dots, u(r_m(s))), x^* \rangle ds. \end{aligned} \quad (3.44)$$

If $Au(t)$ is Bochner integrable (cf. [\[2\]](#)) on $[0, t_0]$, from [\(3.44\)](#), we get

$$\frac{d}{dt} u(t) + Au(t) = f(t, u(t), u(r_1(t)), \dots, u(r_m(t))), \quad \text{a.e. } t \in [0, t_0]. \quad (3.45)$$

To show that $Au(t)$ is Bochner integrable, we will use [Lemmas 2.1](#) and [2.2](#) which state that operators J_n and A_n are Lipschitz continuous. Also,

$$\|A_n u\| \leq \|Au\|, \quad \text{for } u \in D(A). \quad (3.46)$$

Now,

$$\begin{aligned} \|A_n U^n(t)\| &\leq \|A_n U^n(t) - A_n X^n(t)\| + \|A_n X^n(t)\| \\ &\leq 2n \|U^n(t) - X^n(t)\| + \|AX^n(t)\| \\ &\leq K, \end{aligned} \quad (3.47)$$

where K is a constant.

Then [Lemma 2.3\(b\)](#) implies that $A_n U^n(t) \rightharpoonup Au(t)$.

Let H_0 be the smallest closed linear subspace of H containing all the values of the $A_n U^n(t)$ for $t \in [0, t_0]$ and $n = 1, 2, \dots$. Since $A_n U^n(t)$ are continuous, H_0 is separable. Since $A_n U^n(t) \rightharpoonup Au(t)$ and H_0 is weakly closed, $Au(t) \in H_0$ too. Thus $Au(t)$ is separably valued. Since it is weakly continuous, it is strongly measurable and, being bounded, it is Bochner integrable (see [\[12\]](#)).

Clearly, u is Lipschitz continuous on $[0, t_0]$ and $u(t) \in D(A)$ for $t \in [0, t_0]$. Now we prove the uniqueness of a function $u \in \mathcal{C}_{t_0}$ which is differentiable a.e. on $[0, t_0]$ with $u(t) \in D(A)$ a.e. on $[0, t_0]$ and $u = \phi$ on $[-\tau, 0]$ satisfying [\(3.45\)](#). Let $u_1, u_2 \in \mathcal{C}_{t_0}$ be two such functions. Let

$$R = \max \{ \|u_1\|_{t_0}, \|u_2\|_{t_0} \}. \quad (3.48)$$

Then for $u = u_1 - u_2$, we have

$$\begin{aligned} &\left(\frac{du}{dt}(t), u(t) \right) + (Au_1(t) - Au_2(t), u(t)) \\ &= (f(t, u_1(t), u_1(r_1(t)), \dots, u_1(r_m(t))) \\ &\quad - f(t, u_2(t), u_2(r_1(t)), \dots, u_2(r_m(t))), u(t)). \end{aligned} \quad (3.49)$$

By the monotonicity of A ,

$$\begin{aligned} \frac{d}{dt} \|u(t)\|^2 &\leq 2 \|f(t, u_1(t), u_1(r_1(t)), \dots, u_1(r_m(t))) \\ &\quad - f(t, u_2(t), u_2(r_1(t)), \dots, u_2(r_m(t)))\| \|u(t)\|. \end{aligned} \quad (3.50)$$

It implies that

$$\frac{d}{dt} \|u(t)\|^2 \leq C(R) \|u\|_t^2, \quad \text{a.e. } t \in [0, t_0], \quad (3.51)$$

where $C : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a nondecreasing function. Integrating over $(0, s)$ for $0 < s \leq t \leq t_0$, and using the fact that $u \equiv 0$ on $[-\tau, 0]$, we get

$$\|u(s)\|^2 \leq C(R) \int_0^t \|u\|_s^2 ds. \quad (3.52)$$

Taking the supremum over $[0, t]$, we get

$$\sup_{s \in [0, t]} \|u(s)\|^2 \leq C(R) \int_0^t \|u\|_s^2 ds. \quad (3.53)$$

Thus,

$$\|u\|_t^2 \leq C(R) \int_0^t \|u\|_s^2 ds. \quad (3.54)$$

Application of Gronwall's inequality implies that $u \equiv 0$ on $[-\tau, t_0]$.

Now, we prove the continuation of the solution u on $[-\tau, T]$. Suppose $t_0 < T$ and consider the problem

$$\begin{aligned} w'(t) + Aw(t) &= \tilde{f}(t, w(t), w(\tilde{r}_1(t)), w(\tilde{r}_2(t)), \dots, w(\tilde{r}_m(t))), \quad 0 < t \leq T - t_0, \\ w &= \tilde{\chi}, \quad \text{on } [-\tau - t_0, 0], \end{aligned} \quad (3.55)$$

where

$$\begin{aligned} \tilde{f}(t, u_1, u_2, \dots, u_{m+1}) &= f(t + t_0, u_1, u_2, \dots, u_{m+1}), \quad 0 \leq t \leq T - t_0, \\ \tilde{\chi}(t) &= \begin{cases} \chi(t + t_0), & t \in [-\tau - t_0, -t_0], \\ u(t + t_0), & t \in [-t_0, 0], \end{cases} \\ \tilde{r}_i(t) &= r_i(t + t_0) - t_0, \quad t \in [0, T - t_0], \quad i = 1, 2, \dots, m. \end{aligned} \quad (3.56)$$

Since $\tilde{\chi}(0) = u(t_0) \in D(A)$, $\tilde{f}$ satisfies (A3), and $\tilde{r}_i$, $i = 1, 2, \dots, m$, satisfy (A4) on $[0, T - t_0]$, we may proceed as before and prove the existence of a unique $w \in C([-\tau - t_0, t_1]; X)$, $0 < t_1 \leq T - t_0$, such that w is Lipschitz continuous on $[0, t_1]$, $w(t) \in D(A)$ for $t \in [0, t_1]$, and w satisfies

$$\begin{aligned} w'(t) + Aw(t) &= \tilde{f}(t, w(t), w(\tilde{r}_1(t)), w(\tilde{r}_2(t)), \dots, w(\tilde{r}_m(t))), \quad \text{a.e. } t \in [0, t_1], \\ w &= \tilde{\chi}, \quad \text{on } [-\tau - t_0, 0]. \end{aligned} \quad (3.57)$$

Then the function

$$\bar{u}(t) = \begin{cases} u(t), & t \in [-\tau, t_0], \\ w(t - t_0), & t \in [t_0, t_0 + t_1], \end{cases} \quad (3.58)$$

is Lipschitz continuous on $[0, t_0 + t_1]$, $\bar{u}(t) \in D(A)$ for $t \in [0, t_0 + t_1]$, and satisfies (1.4) a.e. on $[0, t_1]$. Continuing this way, we may prove the existence either on the whole interval $[-\tau, T]$ or on the maximal interval of existence $[-\tau, t_{\max})$, $0 < t_{\max} \leq T$. In case $\lim_{t \rightarrow t_{\max}^-} \|u(t)\| < \infty$, then as $u(t) \in D(A)$ for $t \in [0, t_{\max})$, we have that $\lim_{t \rightarrow t_{\max}^-} u(t)$ is in the closure of $D(A)$ in H , and if it is in $D(A)$, then proceeding as before, we may extend $u(t)$ beyond $t_{\max}$ contradicting the definition of the maximal interval of existence.

Now, let u_χ be the strong solution of (1.4) corresponding to $\chi \in \mathcal{C}_{\tilde{T}}$ satisfying $h(\chi|_{[-\tau, 0]}) = \phi_0$ ($\tilde{T}$ is either equal to T or $\tilde{T} < t_{\max}$). If there are $\chi^1, \chi^2 \in \mathcal{C}_{\tilde{T}}$ such that $h(\chi^1|_{[-\tau, 0]}) = h(\chi^2|_{[-\tau, 0]}) = \phi_0$ and $\chi^1 \neq \chi^2$ on $[-\tau, 0]$, then clearly u_{χ^1} and u_{χ^2} , satisfying (1.4), are different. This completes the proof of Theorem 2.4. $\square$

4. Applications. Theorem 2.4 may be applied to get the existence and uniqueness results for (1.1) in the case when the operator A , with the domain $D(A) = H^{2m}(\Omega) \cap H_0^m(\Omega)$ into $H := L^2(\Omega)$, is associated with the nonlinear partial differential operator

$$Au = \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha A_\alpha(x, u(x), Du, \dots, D^\alpha u), \quad (4.1)$$

in a bounded Ω in $\mathbb{R}^n$ with sufficiently smooth boundary $\partial\Omega$, where $A_\alpha(x, \xi)$ are real functions defined on $\Omega \times \mathbb{R}^N$ for some $N \in \mathbb{N}$ and satisfying the following conditions.

- (I) A_α are measurable in x and continuous in ξ . There exist $p = 2$, $g \in L^2(\Omega)$, and a positive constant C such that

$$|A_\alpha(x, \xi)| \leq C(|\xi| + g(x)), \quad \text{a.e. } x \in \Omega, \quad (4.2)$$

where $\xi = (\xi_\alpha; |\alpha| \leq m)$.

- (II) For any $(\xi, \eta) \in \mathbb{R}^N \times \mathbb{R}^N$ and for almost every $x \in \Omega$, the following inequality holds:

$$\sum_{|\alpha| \leq m} (A_\alpha(x, \xi) - A_\alpha(x, \eta))(\xi_\alpha - \eta_\alpha) \geq 0. \quad (4.3)$$

In (1.1), we may take f as the function $f : [0, T] \times (L^2(\Omega))^{m+1} \rightarrow L^2(\Omega)$, given by

$$f(t, u_1, u_2, \dots, u_{m+1}) = f_0(t) + a(t) \sum_{i=1}^{m+1} \|u_i\|_{L^2(\Omega)} u_i, \quad (4.4)$$

where $f_0 : [0, T] \rightarrow L^2(\Omega)$ and $a : [0, T] \rightarrow \mathbb{R}$ are Lipschitz continuous functions on $[0, T]$ and $\|\cdot\|_{L^2(\Omega)}$ denotes the norm in $L^2(\Omega)$. Let $\phi_0 \in C([-\tau, 0]; L^2(\Omega))$ be such that $\phi_0(0) \in D(A)$. For the functions r_i , $i = 1, 2, \dots, n$, and h , we may have any of the following.

- (b1) For $i = 1, 2, \dots, m$, let $r_i(t) = t - \tau_i$, $t \in [0, T]$.
 (b2) For $i = 1, 2, \dots, m$, let $r_i(t) = k_i t$, $t \in [0, T]$, $0 < k_i \leq 1$.
 (b3) Let $T = 1$ and $N_0 \in \mathbb{N}$. For $i = 1, 2, \dots, m$, for $t \in [0, T]$, let $r_i(t) = k_i t^{N_0}$, $0 < k_i \leq 1$.
 (h1) For $u \in C([- \tau, 0]; L^2(\Omega))$, let

$$g(u) = \int_{-\tau}^0 k(s)u(s)ds, \quad \phi_0(t) \equiv u_0, \quad t \in [-\tau, 0], \quad u_0 \in D(A), \quad (4.5)$$

where $\kappa = \int_{-\tau}^0 k(s)ds \neq 0$. Let $h(u)(t) \equiv g(u)$ for $t \in [-\tau, 0]$. In this case, we may take $\chi(t) \equiv (1/\kappa)u_0$ on $[-\tau, T]$.

- (h2) For $u \in C([- \tau, 0]; L^2(\Omega))$, $-\tau \leq a_1 < a_2 < \dots < a_r \leq 0$, and $c_i > 0$, $i = 1, 2, \dots, r$, with $C := \sum_{i=1}^r c_i \neq 0$, let

$$g(u) = \sum_{i=1}^r c_i u(a_i), \quad (4.6)$$

with h and ϕ_0 as in (h1). In this case, we may take $\chi(t) \equiv (1/C)u_0$ for $t \in [-\tau, T]$.

- (h3) With u , a_i , and c_i as in (h2) and $\epsilon_i > 0$ for $i = 1, 2, \dots, r$, let

$$g(u) = \sum_i \frac{c_i}{\epsilon_i} \int_{a_i - \epsilon_i}^{a_i} u(s)ds, \quad (4.7)$$

with h , ϕ_0 , and u_0 as in (h1). In this case, we may take $\chi(t)$ as in (h2).

ACKNOWLEDGMENTS. The authors would like to thank the National Board for Higher Mathematics for providing the financial support to carry out this work under its research project no. NBHM/2001/R&D-II.

REFERENCES

- [1] D. Bahuguna, *Quasilinear integrodifferential equations in Banach spaces*, Nonlinear Anal. **24** (1995), no. 2, 175–183.
- [2] D. Bahuguna and V. Raghavendra, *Application of Rothe's method to nonlinear Schrödinger type equations*, Appl. Anal. **31** (1988), no. 1-2, 149–160.
- [3] ———, *Application of Rothe's method to nonlinear integro-differential equations in Hilbert spaces*, Nonlinear Anal. **23** (1994), no. 1, 75–81.
- [4] K. Balachandran and M. Chandrasekaran, *Existence of solutions of a delay differential equation with nonlocal condition*, Indian J. Pure Appl. Math. **27** (1996), no. 5, 443–449.
- [5] V. Barbu, *Nonlinear Semigroups and Differential Equations in Banach Spaces*, Noordhoff International Publishing, Leiden, 1976.
- [6] L. Byszewski, *Theorems about the existence and uniqueness of solutions of a semilinear evolution nonlocal Cauchy problem*, J. Math. Anal. Appl. **162** (1991), no. 2, 494–505.
- [7] L. Byszewski and V. Lakshmikantham, *Theorem about the existence and uniqueness of a solution of a nonlocal abstract Cauchy problem in a Banach space*, Appl. Anal. **40** (1991), no. 1, 11–19.
- [8] A. G. Kartsatos, *On the construction of methods of lines for functional evolutions in general Banach spaces*, Nonlinear Anal. **25** (1995), no. 12, 1321–1331.
- [9] A. G. Kartsatos and M. E. Parrott, *A method of lines for a nonlinear abstract functional evolution equation*, Trans. Amer. Math. Soc. **286** (1984), no. 1, 73–89.
- [10] ———, *A method of lines for a nonlinear abstract functional evolution equation*, Trans. Amer. Math. Soc. **286** (1984), no. 1, 73–89.

- [11] Y. P. Lin and J. H. Liu, *Semilinear integrodifferential equations with nonlocal Cauchy problem*, Nonlinear Anal. **26** (1996), no. 5, 1023–1033.
- [12] K. Yosida, *Functional Analysis*, Die Grundlehren der Mathematischen Wissenschaften, Band 123, Springer-Verlag, Berlin, 1965.

S. Agarwal: Department of Mathematics, Indian Institute of Technology, Kanpur 208016, India
E-mail address: shrutiag@iitk.ac.in

D. Bahuguna: Department of Mathematics, Indian Institute of Technology, Kanpur 208016, India
E-mail address: dhiren@iitk.ac.in