

DIMENSIONS OF SUBSETS OF CANTOR-TYPE SETS

FANG-XIONG ZHEN

Received 15 March 2006; Accepted 9 July 2006

We define the Cantor-type set E first, and then the Besicovitch subset $B_{\mathbf{p}}$ of E . We mainly show the dimensions of subsets of Cantor-type set E in compatible case and incompatible case.

Copyright © 2006 Hindawi Publishing Corporation. All rights reserved.

1. Introduction

Let $I = [0, 1]$ be the unit interval on the real line and $m > 1$ be an integer. Let $J = \{0, 1, \dots, m - 1\}$. For every point $x \in I$, there is a unique base- m representation $x = \sum_{k=1}^{\infty} j_k m^{-k}$ with $j_k \in J$ except for countable many points. Since countable sets do not interfere with our work, we neglect them here.

For each $j \in J$, $x \in [0, 1]$, and $n \in \mathbb{N}$, let $\tau_j(x, n) = \#\{k : i_k = j, 1 \leq k \leq n\}$, then the limit $\tau_j(x) = \lim_{n \rightarrow \infty} (1/n)\tau_j(x, n)$ is called the frequency of number j in the base- m representation of x . Here and in the following context, the notation “ $\#A$ ” denotes the number of elements in set A .

A classical result of Borel [3] says that for Lebesgue almost every $x \in [0, 1]$, we have $\tau_j(x) = 1/m$. As for another problem, for a given probability vector $\mathbf{p} = (p_0, p_1, \dots, p_{m-1})$ such that $\sum_{j \in J} p_j = 1$, consider the set

$$\Lambda_m(p_0, p_1, \dots, p_{m-1}) = \{x \in [0, 1] : \tau_j(x) = p_j, \text{ for } j \in J\}. \quad (1.1)$$

That is, the set $\Lambda_m(p_0, p_1, \dots, p_{m-1})$ is composed of the number in $[0, 1]$ having a ratio p_j of digits equal to j in its base- m representation for each j . A precursor theory is due to Besicovitch, when he showed in [1] that if $p \in (0, 1/2)$, then

$$\dim \Lambda_2(p, 1 - p) = \frac{p \log p + (1 - p) \log(1 - p)}{-\log 2}. \quad (1.2)$$

2 Dimensions of subsets of cantor-type sets

Eggleston generalized this formula for $m > 2$. He showed in [4] that

$$\dim \Lambda_m(p_0, p_1, \dots, p_{m-1}) = -\frac{\sum_{j \in J} p_j \log p_j}{\log m}. \quad (1.3)$$

From then on, $\Lambda_m(p_0, p_1, \dots, p_{m-1})$ is called Besicovitch set. And later, Billingsley furthered their work and the related information can be found in his book [2].

The research developed in this paper is motivated by Morán and Rey's work, and some of the proofs are edified by the methods mentioned in their article [7]. Let $M = \{1, 2, \dots, m\}$ and for any $j \in M$, $\omega = i_1 i_2 \cdots i_n \cdots \in M^\infty$, $n \in \mathbb{N}$, we define

$$\delta_j(\omega, n) = \#\{k : i_k = j, 1 \leq k \leq n\}. \quad (1.4)$$

Whenever there exists the limit

$$\delta_j(\omega) = \lim_{n \rightarrow \infty} \frac{1}{n} \delta_j(\omega, n), \quad (1.5)$$

it is called the frequency of number j in infinite length word ω . Then for a given probability vector $\mathbf{p} = (p_1, p_2, \dots, p_m)$, we define the Besicovitch set $B_{\mathbf{p}}$ to be the subset of Cantor-type set E (see Section 2) given by

$$B_{\mathbf{p}} = \{\varphi(\omega) : \delta_j(\omega) = p_j, \omega \in M^\infty, \text{ for any } j \in M\}. \quad (1.6)$$

Here φ is a bijective between M^∞ and the Cantor-type set E and M^∞ is the set of all infinite length sequences consisted by M . A remark from [7] says that $B_{\mathbf{p}}$ is a Borel set. In this paper, we give the Hausdorff dimension of $B_{\mathbf{p}}$ for any given probability vector $\mathbf{p} = (p_1, p_2, \dots, p_m)$ with $p_j > 0$ for $j \in M$ as the following theorem.

THEOREM 1.1. *Denote*

$$s = \frac{\sum_{j \in M} p_j \log p_j}{\sum_{j \in M} p_j \log r_j}, \quad (1.7)$$

$\dim B_{\mathbf{p}} = s$, and $s \leq \alpha$, where α is uniquely decided by $\sum_{j \in M} r_j^\alpha = 1$. The equality is attained when \mathbf{p} coincides with $(r_1^\alpha, r_2^\alpha, \dots, r_m^\alpha)$.

Furthermore, we will show that there exists a subset B of Besicovitch set $B_{\mathbf{p}}$ such that B has full $\mu_{\mathbf{p}}$ -measure (see Section 2) but zero Hausdorff measure when $\mathbf{p} \neq (r_1^\alpha, r_2^\alpha, \dots, r_m^\alpha)$, which is described as follows.

THEOREM 1.2. *There exists a Borel subset B of $B_{\mathbf{p}}$ such that $\mu_{\mathbf{p}}(B) = 1$ and $\mathcal{H}^s(B) = 0$ in the incompatible case, that is, $\mathbf{p} \neq (r_1^\alpha, r_2^\alpha, \dots, r_m^\alpha)$.*

As a corollary of Theorem 1.2, we can easily have that $\mu_{\mathbf{p}}$, \mathcal{H}^α , and \mathcal{P}^α are equivalent on E in the compatible case, that is, $\mathbf{p} = (r_1^\alpha, r_2^\alpha, \dots, r_m^\alpha)$ (see Corollary 3.7). Furthermore, we can get another byproduct (see Corollary 3.8) that the set $E \setminus B_{\mathbf{p}}$ has zero \mathcal{H}^α and \mathcal{P}^α -measure when \mathbf{p} is substituted by $(r_1^\alpha, r_2^\alpha, \dots, r_m^\alpha)$.

If the gap condition (see Section 2) holds, we will see that not only $B_{\mathbf{p}}$ and the Cantor-type set E have the same Hausdorff measure and packing measure, but also both of them

are α -sets in the case of $\mathbf{p} = (r_1^\alpha, r_2^\alpha, \dots, r_m^\alpha)$ (see Corollary 3.11). This can be regarded as a corollary from Corollary 3.7. When it comes to the incompatible case, the Hausdorff measure of $B_{\mathbf{p}}$ is infinity (Proposition 3.12), and so is the packing measure.

From Corollary 3.8, we see that the complementary of $B_{\mathbf{p}}$ with respect to E has zero \mathcal{H}^α -measure in the compatible case, which implies that it is rather small from the viewpoint of measure theory.

The following section will give the basic definitions and notations of the paper, and the proofs of the above results are displayed in Section 3.

2. Definitions and notations

Let $I = [0, 1]$ be the unit interval and $M = \{1, 2, \dots, m\}$. Let Ω_n be the set of all sequences of length n with each sequence consisting of letters in M and let $\Omega = \bigcup_{n=1}^{\infty} \Omega_n$. Suppose that a sequence of closed subsets $\{I_\omega : \omega \in \Omega_n\}$ of I satisfies

- (1) $I_{\omega*i} \subset I_\omega$ for any $\omega \in \Omega$ and $i \in M$,
- (2) $I_{\omega*i} \cap I_{\omega*j} = \emptyset$ for any $\omega \in \Omega$ and $i, j \in M$ with $i \neq j$,
- (3) $|I_{\omega*i}|/|I_\omega| = r_i$ with $r_i \in (0, 1)$ for any $\omega \in \Omega$ and $i \in M$.

Then the set $E = \bigcap_{n=1}^{\infty} \bigcup_{\omega \in \Omega_n} I_\omega$ is called a Cantor-type set.

Notice that $\bigcap_{n=1}^{\infty} I_{i_1 i_2 \dots i_n}$ consists of a single point which we will denote by $\varphi(\omega)$ since $\omega \in \Omega_n$, and we write $I_{i_1 i_2 \dots i_n}$ for I_ω sometimes. We can easily see that $\varphi : M^\infty \rightarrow E$ is 1-1 and onto.

We denote by $\nu_{\mathbf{p}}$ the probability measure on M^∞ defined by the infinite product $\mathbf{p} \times \mathbf{p} \times \mathbf{p} \times \dots$. Let $\mu_{\mathbf{p}}$ be the projected measure supported by E , that is, $\mu_{\mathbf{p}} = \nu_{\mathbf{p}} \circ \varphi^{-1}$, then for any n -level basic interval $I_\omega (\omega = i_1 i_2 \dots i_n \in \Omega_n)$, we have $\mu_{\mathbf{p}}(I_\omega) = \prod_{j=1}^n p_{i_j}$. By the *strong law of large numbers*, we see that $\mu_{\mathbf{p}}(B_{\mathbf{p}}) = 1$ and since $B_{\mathbf{p}} \subset E$, thus $\mu_{\mathbf{p}}(B_{\mathbf{p}}) = \mu_{\mathbf{p}}(E) = 1$. Such a measure $\mu_{\mathbf{p}}$ is called Bernoulli measure.

For the use in next section, we give some notations here.

Let $\mathcal{F} = \{I_\omega : \omega \in \Omega\}$, we define

$$\begin{aligned} \mathcal{H}_{\delta, \mathcal{F}}^s(E) &= \inf \left\{ \sum |I_\omega^j|^s : \{I_\omega^j\} \subset \mathcal{F}, |I_\omega^j| \leq \delta, E \subset \bigcup_j I_\omega^j \right\}, \\ \mathcal{H}_{\mathcal{F}}^s(E) &= \lim_{\delta \rightarrow 0} \mathcal{H}_{\delta, \mathcal{F}}^s(E). \end{aligned} \quad (2.1)$$

Similar to the definition of Hausdorff dimension, we define

$$\begin{aligned} \dim_{\mathcal{F}} E &= \sup \{s \geq 0 : \mathcal{H}_{\mathcal{F}}^s(E) = \infty\} \\ &= \inf \{s \geq 0 : \mathcal{H}_{\mathcal{F}}^s(E) = 0\}. \end{aligned} \quad (2.2)$$

Now, we would like to define $\mu_{\mathbf{p}}^s$ -measure as follows:

$$\begin{aligned} \mu_{\mathbf{p}}^s(E, \delta) &= \inf \left\{ \sum \mu_{\mathbf{p}}(U_i)^s : \mu_{\mathbf{p}}(U_i) \leq \delta, E \subset \bigcup_i U_i \right\}, \\ \mu_{\mathbf{p}}^s(E) &= \lim_{\delta \rightarrow 0} \mu_{\mathbf{p}}^s(E, \delta) \end{aligned} \quad (2.3)$$

4 Dimensions of subsets of cantor-type sets

then the $\mu_{\mathbf{p}}$ -dimension of E is given by

$$\begin{aligned} \dim_{\mu_{\mathbf{p}}} E &= \sup \{s \geq 0 : \mu_{\mathbf{p}}^s(E) = \infty\} \\ &= \inf \{s \geq 0 : \mu_{\mathbf{p}}^s(E) = 0\}. \end{aligned} \quad (2.4)$$

We see that if $\mu_{\mathbf{p}}(E) > 0$, then $\dim_{\mu_{\mathbf{p}}} E = 1$.

3. Proofs

Now we set to prove the results displayed in Section 1. First, let us look at some lemmas.

LEMMA 3.1 [5]. *Suppose that $\mathbf{p} = (p_1, p_2, \dots, p_m)$ is a probability vector with $\sum_{j \in M} p_j = 1$ and $p_j > 0$. For any $q_1, \dots, q_m \in \mathbb{R}$,*

$$\sum_{i=1}^m p_i (-\log p_i + q_i) \leq \log \sum_{i=1}^m e^{q_i}. \quad (3.1)$$

The equality is attained if and only if $p_i = e^{q_i} (\sum_{i=1}^m e^{q_i})^{-1}$ for $i \in M$.

LEMMA 3.2. *For any subset $F \subset E$, $\alpha \geq 0$, there exists a constant $c > 0$ such that $c\mathcal{H}_{\mathbb{F}}^{\alpha}(F) \leq \mathcal{H}^{\alpha} \leq \mathcal{H}_{\mathbb{F}}^{\alpha}(F)$.*

Proof. Denote $r = \min_{1 \leq i \leq m} \{r_i\}$, $A_r^{(n)} = \{I_{\omega} : r^n < |I_{\omega}| \leq r^{n-1}\}$. For any δ -covering $\{U_i\}$ of F , for any U_i , there exists n_i such that $r^{n_i} < |U_i| \leq r^{n_i-1}$. For any $x \in U_i$, $F \cap U_i \subset \bigcup_{I_{\omega} \in A_r^{n_i}, I_{\omega} \cap U_i \neq \emptyset} I_{\omega} \subset B(x, 2r^{n_i-1})$, then we have $\#\{I_{\omega} : I_{\omega} \in A_r^{n_i}, I_{\omega} \cap U_i \neq \emptyset\} \leq 4r^{n_i-1}/r^{n_i} = 4/r$, so for this U_i , there exists a sequence $\{I_{ij}^{n_i}\}_{i=1,2,\dots}^{l_i} \subset A_r^{n_i}$ such that $F \cap U_i \subset \bigcup_{j=1}^{l_i} I_{ij}^{(n_i)}$, $l_i \leq 4/r$. Since $|U_i| \geq r^{n_i} \geq r |I_{ij}^{(n_i)}|$ for $j = 1, 2, \dots, l_i$, then $l_i |U_i|^{\alpha} \geq \sum_{j=1}^{l_i} r^{\alpha} |I_{ij}^{(n_i)}|^{\alpha}$, we further get $(4/r) \sum_{i=1}^{\infty} |U_i|^{\alpha} \geq r^{\alpha} \sum_{i=1}^{\infty} \sum_{j=1}^{l_i} |I_{ij}^{(n_i)}|^{\alpha}$, thus we get $\mathcal{H}^{\alpha}(F) \geq (r^{\alpha+1}/4) \mathcal{H}_{\mathbb{F}}^{\alpha}(F)$. Letting $c = r^{\alpha+1}/4$, we have $c\mathcal{H}_{\mathbb{F}}^{\alpha}(F) \leq \mathcal{H}^{\alpha} \leq \mathcal{H}_{\mathbb{F}}^{\alpha}(F)$. \square

To prove Theorem 1.1, we need another lemma.

LEMMA 3.3. *If*

$$E \subset \left\{ \varphi(\omega) : \lim_{|\omega| \rightarrow \infty} \frac{\log \mu_{\mathbf{p}}(I_{\omega})}{\log |I_{\omega}|} = \delta \right\}, \quad (3.2)$$

then $\dim E = \delta \dim_{\mu_{\mathbf{p}}} E$.

Proof. With Lemma 3.2, we can get the same dimension result if we use the basic intervals to cover E , so we first prove that if

$$E \subset \left\{ \varphi(\omega) : \liminf_{|\omega| \rightarrow \infty} \frac{\log \mu_{\mathbf{p}}(I_{\omega})}{\log |I_{\omega}|} \geq \delta \right\}, \quad (3.3)$$

then $\dim E \geq \delta \dim_{\mu_{\mathbf{p}}} E$.

If $\varphi(\omega) \in E$, then for any $\varepsilon > 0$, there exists $N > 0$ such that for any $n > N$, we have $\mu_{\mathbf{p}}(I_{\omega}) \leq |I_{\omega}|^{\delta-\varepsilon}$. Thus there are only many finite ω such that I_{ω} does not satisfy the above

inequality. Let

$$E_k = \left\{ \varphi(\omega) \in E : |I_\omega| > \frac{1}{k} \text{ or } \mu_{\mathbf{p}}(I_\omega) \leq |I_\omega|^{\delta-\varepsilon} \right\}. \quad (3.4)$$

We see that E_k is increasing. On the other hand, since we have $\mu_{\mathbf{p}}(I_\omega) > 0$ for any $\omega \in \Omega$, then there exists k such that $|I_\omega| > 1/k$ for any given $\omega \in \Omega$, which implies that $\varphi(\omega) \in E_k$. Thus we have $E = \bigcup_k E_k$.

For any $(1/k >)\rho > 0$, $\varepsilon_1 > 0$, $\varepsilon_2 > 0$, there exists a sequence of intervals $\{I_\omega^j\} \subset \mathcal{F}$ such that $|I_\omega^j| \leq \rho$, $I_\omega^j \cap E_k \neq \emptyset$, and $E_k \subset \bigcup_j I_\omega^j$. Since $\dim_{\mathcal{F}} E = \dim E$ and $E_k \subset E$, we have $\mathcal{H}_\rho^{\dim E + \varepsilon_2}(E_k) < \varepsilon_1$, that is, $\sum |I_\omega^j|^{\dim E + \varepsilon_2} < \varepsilon_1$, which follows $\sum (\mu_{\mathbf{p}}(I_\omega^j))^{\dim E + \varepsilon_2 / (\delta - \varepsilon)} < \varepsilon_1$. Thus we have $\dim_{\mu_{\mathbf{p}}} E_k \leq (\dim E + \varepsilon_2) / (\delta - \varepsilon)$. Since ε and ε_2 are arbitrary small, we have $\dim E \geq \delta \dim_{\mu_{\mathbf{p}}} E_k$, which further implies $\dim E \geq \delta \dim_{\mu_{\mathbf{p}}} E$.

With the same method we can prove that if

$$E \subset \left\{ \varphi(\omega) : \limsup_{|\omega| \rightarrow \infty} \frac{\log \mu_{\mathbf{p}}(I_\omega)}{\log |I_\omega|} \leq \delta \right\}, \quad (3.5)$$

then $\dim E \leq \delta \dim_{\mu_{\mathbf{p}}} E$.

With the above arguments, we complete the proof. \square

Proof of Theorem 1.1. For any $\omega \in \Omega$,

$$\begin{aligned} \lim_{|\omega| \rightarrow \infty} \frac{\log \mu_{\mathbf{p}}(I_\omega)}{\log |I_\omega|} &= \lim_{|\omega| \rightarrow \infty} \frac{\sum_{j=1}^n \log p_{i_j}}{\sum_{j=1}^n \log r_{i_j}} \\ &= \lim_{|\omega| \rightarrow \infty} \frac{\sum_{j \in M} \delta_j(\omega, n) \log p_j}{\sum_{j \in M} \delta_j(\omega, n) \log r_j} \\ &= \frac{\sum_{j \in M} p_j \log p_j}{\sum_{j \in M} p_j \log r_j} = s. \end{aligned} \quad (3.6)$$

Since $\mu_{\mathbf{p}}(B_{\mathbf{p}}) = 1$, by Lemma 3.3 we have $\dim B_{\mathbf{p}} = s$.

Put $q_i = \log r_i^\alpha$ in Lemma 3.1, we get $s \leq \alpha$. Furthermore, if we put

$$p_i = e^{\log r_i^\alpha} \left(\sum_{i=1}^m e^{\log r_i^\alpha} \right)^{-1} = r_i^\alpha, \quad (3.7)$$

we have $s = \alpha$. \square

LEMMA 3.4 (law of iterated logarithm (Hartman-Winter)). *Suppose that $\{X_n : n \geq 1\}$ is a sequence of independent, identically distributed random variables (i.i.d.r.v.) satisfying $\mathbf{E}(X_n) = 0$ and $\mathbf{E}(X_n^2) = \sigma^2 \in (0, +\infty)$, then*

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{\sum_{i=1}^n X_i}{\sqrt{2n\sigma^2 \log \log n\sigma^2}} &= 1, \quad p \text{ a.e.}, \\ \liminf_{n \rightarrow \infty} \frac{\sum_{i=1}^n X_i}{\sqrt{2n\sigma^2 \log \log n\sigma^2}} &= -1 \quad p \text{ a.e.} \end{aligned} \quad (3.8)$$

6 Dimensions of subsets of cantor-type sets

Remark 3.5. We suppose that \mathbf{E} denotes the mathematical expectation and \mathbf{V} denotes the mean variance here and in the following context.

Proof of Theorem 1.2. Put $p = \mu_{\mathbf{p}} \circ \varphi$, $X_i = \log p_i - \log r_i$, $i \in M$, then $\{X_i\}$ is a sequence of i.i.d.r.v. with respect to p . We see that $\mathbf{E}(X_i) = \sum_{i=1}^m p_i (\log p_i - \log r_i) = 0$. Suppose that $\mathbf{p} \neq (r_1^\alpha, \dots, r_m^\alpha)$, then

$$0 < \mathbf{E}(X_i^2) = \sum_{i=1}^m p_i (\log p_i - \log r_i)^2 =: \sigma^2 < \infty. \quad (3.9)$$

Set

$$B' = \left\{ \varphi(\omega) : \limsup_{n \rightarrow \infty} \frac{\sum_{i=1}^n X_i}{\sqrt{2n\sigma^2 \log \log n\sigma^2}} = 1 \right\}, \quad (3.10)$$

then $\mu_{\mathbf{p}}(B') = 1$ by Lemma 3.4. Since $\mu_{\mathbf{p}}(B_{\mathbf{p}}) = 1$. Let $B = B' \cap B_{\mathbf{p}}$, we get $\mu_{\mathbf{p}}(B) = 1$.

With the above arguments, we see that $\sum_{j=1}^{\infty} X_{i_j} = \infty$ a.e., that is,

$$\log \frac{\prod_{j=1}^n p_{i_j}}{\prod_{j=1}^n r_{i_j}^s} \rightarrow \infty, \quad \text{as } n \rightarrow \infty, \quad (3.11)$$

from which we also have

$$\frac{\mu_{\mathbf{p}}(I_\omega)}{|I_\omega|^s} \rightarrow \infty, \quad \text{as } |\omega| \rightarrow \infty. \quad (3.12)$$

Let $\mathcal{F}_n^k = \{I_\omega : \mu_{\mathbf{p}}(I_\omega)/|I_\omega|^s > k, \text{ for } \omega \in \Omega_n\}$, $\mathcal{F}^k = \bigcup_{n=1}^{\infty} \mathcal{F}_n^k$. For any $\delta > 0$, $k > 0$, choose $\{I_j^k\} \subset \mathcal{F}^k$ such that $B \subset \bigcup_j I_j^k$, $|I_j^k| \leq \delta$, and $I_i^k \cap I_j^k = \emptyset (i \neq j)$. By the definition of \mathcal{F}^k , we have

$$\frac{\mu_{\mathbf{p}}(I_j^k)}{|I_j^k|^s} > k, \quad \forall j. \quad (3.13)$$

Thus

$$\mathcal{H}_\delta^s(B) \leq \sum |I_j^k|^s \leq \frac{1}{k} \sum \mu_{\mathbf{p}}(I_j^k) \leq \frac{1}{k}. \quad (3.14)$$

Letting $k \rightarrow \infty$, we have $\mathcal{H}_\delta^s(B) = 0$, from which it follows that $\mathcal{H}^s(B) = 0$. \square

LEMMA 3.6 [6]. *Suppose that μ is a finite Borel measure such that*

$$0 < \limsup_{|\omega| \rightarrow \infty} \frac{\mu(I_\omega)}{|I_\omega|^\alpha} < \infty, \quad 0 < \liminf_{|\omega| \rightarrow \infty} \frac{\mu(I_\omega)}{|I_\omega|^\alpha} < \infty, \quad \text{for any } \omega \in \Omega, \quad (3.15)$$

then $\mu \sim \mathcal{H}^\alpha \sim \mathcal{P}^\alpha$ on E .

COROLLARY 3.7. $\mu_{\mathbf{p}} \sim \mathcal{H}^\alpha \sim \mathcal{P}^\alpha$ on E in case of $\mathbf{p} = (r_1^\alpha, r_2^\alpha, \dots, r_m^\alpha)$.

Proof. Since $\mu_{\mathbf{p}}(I_{\omega}) = \prod_{j=1}^n p_{i_j}$, for $\omega = i_1 i_2 \cdots i_n \in \Omega_n$, then

$$\frac{\mu_{\mathbf{p}}(I_{\omega})}{|I_{\omega}|^{\alpha}} = \frac{\prod_{j=1}^n p_{i_j}}{\prod_{j=1}^n r_{i_j}^{\alpha}} = 1, \quad \text{when } \mathbf{p} = (r_1^{\alpha}, r_2^{\alpha}, \dots, r_m^{\alpha}). \quad (3.16)$$

So by Lemma 3.6, we have $\mu_{\mathbf{p}} \sim \mathcal{H}^{\alpha} \sim \mathcal{P}^{\alpha}$ on E in case of $\mathbf{p} = (r_1^{\alpha}, r_2^{\alpha}, \dots, r_m^{\alpha})$. \square

COROLLARY 3.8. $\mathcal{H}^{\alpha}(E \setminus B_{\mathbf{p}}) = \mathcal{P}^{\alpha}(E \setminus B_{\mathbf{p}}) = 0$, $\mathcal{H}^{\alpha}(E) = \mathcal{H}^{\alpha}(B_{\mathbf{p}})$, and $\mathcal{P}^{\alpha}(E) = \mathcal{P}^{\alpha}(B_{\mathbf{p}})$ in case of $\mathbf{p} = (r_1^{\alpha}, r_2^{\alpha}, \dots, r_m^{\alpha})$.

Proof. By the definition of equivalence, $\mu_{\mathbf{p}} \sim \mathcal{H}^{\alpha}$ on $E \Leftrightarrow$ for any $F \subset E$, $\mu_{\mathbf{p}}(F) = 0 \Leftrightarrow \mathcal{H}^{\alpha}(F) = 0$. It is clear that $B_{\mathbf{p}}$ is the subset of E with $\mu_{\mathbf{p}}(E) = \mu_{\mathbf{p}}(B_{\mathbf{p}}) = 1$, then we get immediately $\mu_{\mathbf{p}}(E \setminus B_{\mathbf{p}}) = 0$, which further implies $\mathcal{H}^{\alpha}(E \setminus B_{\mathbf{p}}) = 0$ by Corollary 3.7. Furthermore, $\mathcal{H}^{\alpha}(B_{\mathbf{p}}) \leq \mathcal{H}^{\alpha}(E) \leq \mathcal{H}^{\alpha}(B_{\mathbf{p}}) + \mathcal{H}^{\alpha}(E \setminus B_{\mathbf{p}}) = \mathcal{H}^{\alpha}(B_{\mathbf{p}})$, we have $\mathcal{H}^{\alpha}(E) = \mathcal{H}^{\alpha}(B_{\mathbf{p}})$.

Similar arguments will prove that $\mathcal{P}^{\alpha}(E \setminus B_{\mathbf{p}}) = 0$ and $\mathcal{P}^{\alpha}(E) = \mathcal{P}^{\alpha}(B_{\mathbf{p}})$. \square

For the simplicity of proof of the following results, we would like to make a hypothesis here.

We say that the Cantor-type set E satisfies the gap condition if there exists a constant $\delta > 0$ such that

$$\text{dist}(I_{i_1 i_2 \cdots i_n}, I_{i_1 i_2 \cdots i'_n}) \geq \delta \max\{|I_{i_1 i_2 \cdots i_n}|, |I_{i_1 i_2 \cdots i'_n}|\} \quad (3.17)$$

for any $n \in \mathbb{N}$ and $i_n \neq i'_n$.

LEMMA 3.9 [6]. *If the gap condition holds, then for any $\omega \in \Omega$ and finite Borel measure μ , the following inequalities hold:*

$$\begin{aligned} (a\delta)^{\alpha} \liminf_{|\omega| \rightarrow \infty} \frac{|I_{\omega}|^{\alpha}}{\mu(I_{\omega})} &\leq \liminf_{r \rightarrow 0} \frac{r^{\alpha}}{\mu(B(\varphi(\omega), r))} \leq a^{-\alpha} \liminf_{|\omega| \rightarrow \infty} \frac{|I_{\omega}|^{\alpha}}{\mu(I_{\omega})}, \\ (a\delta)^{\alpha} \limsup_{|\omega| \rightarrow \infty} \frac{|I_{\omega}|^{\alpha}}{\mu(I_{\omega})} &\leq \limsup_{r \rightarrow 0} \frac{r^{\alpha}}{\mu(B(\varphi(\omega), r))} \leq a^{-\alpha} \limsup_{|\omega| \rightarrow \infty} \frac{|I_{\omega}|^{\alpha}}{\mu(I_{\omega})}, \end{aligned} \quad (3.18)$$

where $a = \min_{i \in M} \{r_i\}$.

LEMMA 3.10 [8]. *For finite Borel measure μ , there exists constants c_1, c_2, c_3, c_4 such that*

$$\begin{aligned} c_1 \mu(E) \inf_{x \in E} \liminf_{r \rightarrow 0} \frac{r^{\alpha}}{\mu(B(x, r))} &\leq \mathcal{H}^{\alpha}(E) \leq c_2 \mu(\mathbb{R}) \sup_{x \in E} \liminf_{r \rightarrow 0} \frac{r^{\alpha}}{\mu(B(x, r))}, \\ c_3 \mu(E) \inf_{x \in E} \limsup_{r \rightarrow 0} \frac{r^{\alpha}}{\mu(B(x, r))} &\leq \mathcal{P}^{\alpha}(E) \leq c_4 \mu(\mathbb{R}) \sup_{x \in E} \limsup_{r \rightarrow 0} \frac{r^{\alpha}}{\mu(B(x, r))}. \end{aligned} \quad (3.19)$$

COROLLARY 3.11. *If E is the Cantor-type set satisfying gap condition and $B_{\mathbf{p}}$ is the Besicovitch set defined in Section 1, then $0 < \mathcal{H}^{\alpha}(E) = \mathcal{H}^{\alpha}(B_{\mathbf{p}}) < \infty$ and $0 < \mathcal{P}^{\alpha}(E) = \mathcal{P}^{\alpha}(B_{\mathbf{p}}) < \infty$, when $\mathbf{p} = (r_1^{\alpha}, r_2^{\alpha}, \dots, r_m^{\alpha})$.*

8 Dimensions of subsets of cantor-type sets

Proof. From Lemma 3.10, we see that the density of balls is comparable with that of intervals. So we have

$$c'_1 \mu(E) \inf_{x \in E} \liminf_{|\omega| \rightarrow \infty} \frac{|I_\omega|^\alpha}{\mu(I_\omega)} \leq \mathcal{H}^\alpha(E) \leq c'_2 \mu(\mathbb{R}) \sup_{x \in E} \liminf_{|\omega| \rightarrow \infty} \frac{|I_\omega|^\alpha}{\mu(I_\omega)} \quad (3.20)$$

together with Lemmas 3.9 and 3.10.

Since

$$\frac{\mu_{\mathbf{p}}(I_\omega)}{|I_\omega|^\alpha} = 1, \quad \text{in case of } \mathbf{p} = (r_1^\alpha, r_2^\alpha, \dots, r_m^\alpha), \quad (3.21)$$

so we get $c'_1 \leq \mathcal{H}^\alpha(E) \leq c'_2$ by substituting μ with $\mu_{\mathbf{p}}$ in (3.20), which implies that $0 < \mathcal{H}^\alpha(E) < \infty$. Since we have proved in Corollary 3.8 that $\mathcal{H}^\alpha(E) = \mathcal{H}^\alpha(B_{\mathbf{p}})$, we have $0 < \mathcal{H}^\alpha(E) = \mathcal{H}^\alpha(B_{\mathbf{p}}) < \infty$.

Analogously, we can prove $0 < \mathcal{P}^\alpha(E) = \mathcal{P}^\alpha(B_{\mathbf{p}}) < \infty$. \square

Remark 3.12. Corollary 3.11 reinforces the assertion in Theorem 1.1.

Acknowledgment

The author expresses his deep appreciation to Dr. Wu Ya-hao for his helpful suggestions.

References

- [1] A. S. Besicovitch, *On the sum of digits of real numbers represented in the dyadic system*, *Mathematische Annalen* **110** (1935), no. 1, 321–330.
- [2] P. Billingsley, *Ergodic Theory and Information*, John Wiley & Sons, New York, 1965.
- [3] E. Borel, *Sur les probabilités dénombrables et leurs applications arithmétiques*, *Rendiconti del Circolo Matematico di Palermo* **26** (1909), 247–271.
- [4] H. G. Eggleston, *The fractional dimension of a set defined by decimal properties*, *The Quarterly Journal of Mathematics. Oxford. Second Series* **20** (1949), 31–36.
- [5] K. Falconer, *Techniques in Fractal Geometry*, John Wiley & Sons, Chichester, 1997.
- [6] S. Ikeda and M. Nakamura, *Dimensions of measures on perturbed Cantor sets*, *Topology and Its Applications* **122** (2002), no. 1-2, 223–236.
- [7] M. Morán and J.-M. Rey, *Singularity of self-similar measures with respect to Hausdorff measures*, *Transactions of the American Mathematical Society* **350** (1998), no. 6, 2297–2310.
- [8] S. J. Taylor and C. Tricot, *Packing measure, and its evaluation for a Brownian path*, *Transactions of the American Mathematical Society* **288** (1985), no. 2, 679–699.

Fang-Xiong Zhen: Department of Mathematics, Xianning College, Xianning, Hubei 437005, China
E-mail address: cookie4993@sina.com



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

