

WEAK GROTHENDIECK'S THEOREM

LAHCÈNE MEZRAG

Received 14 June 2005; Revised 9 March 2006; Accepted 20 June 2006

Let $E_n \subset L_1^{2n}$ be the n -dimensional subspace which appeared in Kašin's theorem such that $L_1^{2n} = E_n \oplus E_n^\perp$ and the L_1^{2n} and L_2^{2n} norms are universally equivalent on both E_n and E_n^\perp . In this paper, we introduce and study some properties concerning extension and weak Grothendieck's theorem (WGT). We show that the Schatten space S_p for all $0 < p \leq \infty$ does not verify the theorem of extension. We prove also that S_p fails GT for all $1 \leq p \leq \infty$ and consequently by one result of Maurey does not satisfy WGT for $1 \leq p \leq 2$. We conclude by giving a characterization for spaces verifying WGT.

Copyright © 2006 Hindawi Publishing Corporation. All rights reserved.

1. Introduction

This work was inspired by the celebrated theorem of Kašin [5]. We use his decomposition cited in the abstract and which states that L_1^{2n} (this space is of dimension $2n$ and which will be defined in the sequel) can be decomposed into two orthogonal n -dimensional subspaces "respecting" the inner product induced by the norm of L_2^{2n} and on each the norms of L_1^{2n} and L_2^{2n} are universally equivalent on these subspaces. It is interesting to observe that the constants of equivalence are independent of n . Recently this was investigated by Anderson [1] and Schechtman [15]. We will say that a Banach space X verifies weak Grothendieck's theorem if $\pi_2(X, l_2) = B(X, l_2)$. Let $\{\varphi_i\}_{1 \leq i \leq n}$ be a sequence of orthogonal random variables in L_2^{2n} , which generates E_n . Consider $0 < p \leq \infty$. Let $u : E_n \rightarrow S_p^n$ be a linear operator and let \tilde{u} be any extension of u . In this paper we show that $\|\tilde{u}\| \geq C\sqrt{n}$, where C is an absolute constant. We prove that S_p fails extension theorem for all $1 \leq p \leq \infty$. We also show that S_p does not verify GT for $1 \leq p \leq \infty$ and consequently fails WGT for all $1 \leq p \leq 2$ by using one result of Maurey. We end this work by giving a characterization for operators satisfying WGT.

We start the first section by recalling some necessary notations and definitions such as the definition of cotype q -Kašin as studied in [9] and which is inspired by the Kašin decomposition. We introduce also the property of weak Grothendieck's theorem.

In section two, we recall the Schatten spaces S_p which are the noncommutative analogues of the l_p -spaces and we give some properties concerning these spaces. After this,

2 Weak Grothendieck's theorem

we show that the space S_p fails the property of extension for all $0 < p \leq \infty$ and GT for all $1 \leq p \leq \infty$. We deduce that the space S_p does not verify WGT for all p , $1 \leq p \leq 2$. We do not know if S_p is of cotype 2-Kašin for $1 \leq p \leq 2$ like the classical cotype. We know that the Schatten space S_p is of cotype 2 for $1 \leq p \leq 2$ as the usual l_p -spaces; see [16]. By another method which is not adjustable to our case we have proved in [10] that $L_p([0, 1], dx)$ and l_p for $0 < p < 1$ fail the extension property.

In Section 4, we characterize the spaces which satisfy weak Grothendieck's theorem.

2. Notation and preliminaries

Let $0 < p \leq +\infty$. We denote by L_p^n the space \mathbb{R}^n (or \mathbb{C}^n) equipped with the norm (and only a p -norm if $0 < p < 1$)

$$\|(a_i)\|_{L_p^n} = \left(\frac{1}{n} \sum_{i=1}^n |a_i|^p \right)^{1/p}, \quad (2.1)$$

and if $p = \infty$, we take $\max |a_i|$.

Recall that a p -norm on a vector space X is a functional

$$\begin{aligned} \|\cdot\| : X &\longrightarrow \mathbb{R}_+, \\ x &\longmapsto \|x\| \end{aligned} \quad (2.2)$$

such that

$$\begin{aligned} \|x\| &= 0 \iff x = 0, \\ \|\lambda x\| &= |\lambda| \|x\| \quad \forall \lambda \text{ in } \mathbb{C}, \\ \|x + y\| &\leq (\|x\|^p + \|y\|^p)^{1/p} \quad \forall x, y \text{ in } X, \end{aligned} \quad (2.3)$$

X is called a p -normed space if its topology can be defined by a p -norm.

L_p^n is isometric to $L_p^n(\Omega_n, \mathcal{P}(\Omega_n), \mu_n)$ where Ω_n is the set $\{1, 2, \dots, n\}$, $\mathcal{P}(\Omega_n)$ the σ -algebra of all subsets $A \subset \Omega_n$ and μ_n the uniform probability on Ω_n (i.e., $\mu_n(i) = 1/n$ for all i in Ω_n). Hence each element in L_p^n can be considered as a random variable which we denote in the sequel by φ and we have for $0 < p \leq q \leq \infty$,

$$\|\varphi\|_{L_p^n} \leq \|\varphi\|_{L_q^n} \leq n^{1/p-1/q} \|\varphi\|_{L_p^n}. \quad (2.4)$$

Moreover, we will denote by $l_p^n(X)$ for any Banach space X (resp., $L_p^n(X)$), the space X^n equipped with the norm if $1 \leq p \leq +\infty$ and the p -norm if $0 < p < 1$:

$$\begin{aligned} \|(x_i)\|_{l_p^n(X)} &= \left(\sum_{i=1}^n \|x_i\|_X^p \right)^{1/p}, \\ \left(\text{resp., } \|(x_i)\|_{L_p^n(X)} &= \left(\frac{1}{n} \sum_{i=1}^n \|x_i\|_X^p \right)^{1/p} \right) \end{aligned} \quad (2.5)$$

for all $(x_i)_{1 \leq i \leq n} \subset X$. If $p = \infty$, the sums should be replaced by sup.

We will use the following decomposition due to B. S. Kašin (see also [13] and recently [1, 15]), which is the principal inspiration of our idea.

THEOREM 2.1 [5]. *Consider p in $\{1, 2\}$ and n in \mathbb{N} . There are three constants A_p , B_p , and C (C independent of p and n) and a sequence $(\varphi_i)_{1 \leq i \leq n}$ of orthogonal random variables in L_2^{2n} such that for all $(a_i)_{1 \leq i \leq n}$ in \mathbb{R} , there exist*

$$\begin{aligned} A_p \left(\sum_1^n \|a_i\|^2 \right)^{1/2} &\leq \left\| \sum_1^n a_i \varphi_i \right\|_{L_p^{2n}} \leq B_p \left(\sum_1^n |a_i|^2 \right)^{1/2}, \\ \sup_{1 \leq i \leq n} \|\varphi_i\|_{L_\infty^n} &\leq C(\log n)^{1/2}. \end{aligned} \quad (2.6)$$

Remark 2.2. It is well known that if X is a finite dimensional space, then, all the norms are equivalent. But what is most remarkable in Theorem 2.1 is that the constants are independent of the dimension n . It is also true for all p in $]0, 2]$. We can and do choose the φ_i to be orthonormal, that is what we do in the sequel.

Let E_n be the subspace of L_1^{2n} spanned by the functions $(\varphi_i)_{1 \leq i \leq n}$ and let $\mathbf{e}_n : E_n \rightarrow L_1^{2n}$ be the natural injection. By the above theorem, E_n is isomorphic to l_2^n , we denote by $\beta_n : l_2^n \rightarrow E_n$ the isomorphism which maps e_i onto φ_i , where (e_i) the unit vector basis of l_2^n . We have by (2.6) that $\|\beta_n\| \leq B_1$ and $\|\beta_n^{-1}\| \leq A_1^{-1}$.

Now we give the following definition which is introduced in [9].

Definition 2.3. Let X and Y be Banach spaces and let $u : X \rightarrow Y$ be a linear operator. Say that u is of cotype q -Kašin for $2 \leq q < +\infty$, if there is a positive constant K such that for all integer n and for all finite sequence $(x_i)_{1 \leq i \leq n}$ in X , there exists

$$\left(\sum_{i=1}^n \|u(x_i)\|^q \right)^{1/q} \leq K \left\| \sum_{i=1}^n \varphi_i x_i \right\|_{L_1^{2n}(X)}. \quad (2.7)$$

Denote by $K_q(u)$ the smallest constant for which this holds. X is of cotype q -Kašin if the identity of X is of cotype q -Kašin.

For example L_p ($1 \leq p \leq 2$) is of cotype 2-Kašin.

For being complete, we add (see [13, page 115]) that there is an orthonormal basis (φ_n) of $L_2([0, 1], \nu)$ (ν is the Lebesgue measure) such that the L_1 and L_2 norms are equivalent on each of the spans of $\{\varphi_n, n \text{ odd}\}$ and $\{\varphi_n, n \text{ even}\}$. Let E_0 be the space spanned by one of these sequences in $L_1([0, 1], \nu)$ and let $e : E_0 \rightarrow L_1([0, 1], \nu)$ be the isometric embedding. We denote also by E_0^n the space spanned by the n first φ_i .

Given two Banach spaces X and Y , denote by $X \otimes_\epsilon Y$ their injective tensor product, that is, the completion of $X \otimes Y$ under the cross norm:

$$\left\| \sum_{i=1}^n x_i \otimes y_i \right\|_\epsilon = \sup \left\{ \left\| \sum_{i=1}^n x_i(\xi) y_i(\eta) \right\| : \|\xi\|_{X^*} \leq 1, \|\eta\|_{Y^*} \leq 1 \right\}. \quad (2.8)$$

Let $u : X \rightarrow Y$ be a linear operator. We will say that u is *absolutely p -summing*, $0 < p < \infty$ (we write $u \in \Pi_p(X, Y)$), if there exists a positive constant C such that for every n in \mathbb{N} ,

4 Weak Grothendieck's theorem

the mappings

$$\begin{aligned} I_n \otimes u : l_p^n \otimes_\epsilon X &\longrightarrow l_p^n(Y), \\ \sum_{i=1}^n e_i \otimes x_i &\longmapsto (u(x_i))_{1 \leq i \leq n} \end{aligned} \quad (2.9)$$

are uniformly bounded by C (i.e., $\|I_n \otimes u\|_{l_p^n \otimes_\epsilon X \rightarrow l_p^n(Y)} \leq C$).

We define the p -summing norm of an operator u by

$$\pi_p(u) = \sup_n \|I_n \otimes u\|_{l_p^n \otimes_\epsilon X \rightarrow l_p^n(Y)}. \quad (2.10)$$

The following proposition is a characterization of spaces of cotype 2-Kašin.

PROPOSITION 2.4. *Let C be a positive constant. Then the following properties of a Banach space X are equivalent.*

- (i) *The space X^* (X^* is the Banach space dual of X) is of cotype 2-Kašin and $K_2(X^*) \leq C$.*
- (ii) *For all integers n and for all finite sequences $(x_i)_{1 \leq i \leq n}$ in X , the operator $u : E_n \rightarrow X$ defined by $u(\varphi_i) = x_i$ admits an extension $\tilde{u} : L_1^{2n} \rightarrow X$ such that $\tilde{u}/E_n = u$ and $\|\tilde{u}\| \leq C(\sum_{i=1}^n \|x_i\|^2)^{1/2}$.*

Proof. Let n be a fixed integer. Since X^* is of cotype 2-Kašin, hence for all $(\xi_i)_{1 \leq i \leq n} \subset X^*$ we have

$$\left(\sum_{i=1}^n \|\xi_i\|_{X^*}^2 \right)^{1/2} \leq C \left\| \sum_{i=1}^n \varphi_i \xi_i \right\|_{L_1^{2n}(X^*)}. \quad (2.11)$$

Let $E = \{ \sum_{i=1}^n \varphi_i \xi_i, (\xi_i)_{1 \leq i \leq n} \subset X^* \}$, which is a closed subspace of $L_1^{2n}(X^*)$. We now define the operators

$$\begin{aligned} T : E &\longrightarrow l_2^n(X^*), \\ \sum_{i=1}^n \varphi_i \xi_i &\longmapsto (\xi_i)_{1 \leq i \leq n}. \end{aligned} \quad (2.12)$$

This definition is unambiguous (indeed, $\sum_{i=1}^n \varphi_i \xi_i = \sum_{i=1}^n \varphi_i \eta_i$ implies that $\xi_i = \eta_i$ for all $1 \leq i \leq n$ because the φ_i are orthogonal and consequently $(\xi_i)_{1 \leq i \leq n} = (\eta_i)_{1 \leq i \leq n}$).

Observe that

$$\|T\| \leq C. \quad (2.13)$$

By duality we have

$$\begin{aligned} T^* : l_2^n(X) &\longrightarrow \frac{L_\infty^{2n}(X)}{E^\perp}, \\ (x_i)_{1 \leq i \leq n} &\longmapsto \sum_{i=1}^n x_i \varphi_i + E^\perp, \end{aligned} \quad (2.14)$$

where $E^\perp = \{\sum_{i=n+1}^{2n} \varphi_i x'_i, (x'_i)_{n+1 \leq i \leq 2n} \subset X\}$ is the subspace of $L_\infty^{2n}(X)$ which is orthogonal to E .

Since $\|T\| = \|T^*\|$ (T^* is the adjoint operator of T), hence we have

$$\inf_{R \in E^\perp} \left\| \sum_{i=1}^n x_i \varphi_i + R \right\|_{L_\infty^{2n}(X)} \leq C \left(\sum_{i=1}^n \|x_i\|^2 \right)^{1/2}. \quad (2.15)$$

If now $\tilde{u} : L_1^{2n} \rightarrow X$ is an extension of u , by Riesz representation theorem then there is Ψ in L_∞^{2n} such that

$$\begin{aligned} \forall \varphi \in L_1^{2n}, \quad \tilde{u}(\varphi) &= \frac{1}{2n} \sum_{i=1}^{2n} \varphi_i \Psi_i, \\ \|\tilde{u}\| &= \|\Psi\|_{L_\infty^{2n}}. \end{aligned} \quad (2.16)$$

Since $\tilde{u}(\varphi_i) = x_i$, we have

$$\Psi = \sum_{i=1}^n x_i \varphi_i + R. \quad (2.17)$$

The correspondence $\tilde{u} \rightarrow \Psi$ is bijective and this implies that

$$\inf \|\tilde{u}\| = \inf_{R \in E^\perp} \left\| \sum_{i=1}^n x_i \varphi_i + R \right\|_{L_\infty^{2n}(X)}. \quad (2.18)$$

This concludes the proof. \square

We say now that a Banach space X is of cotype strongly 2-Kašin if there is a positive constant C such that, for all integers n and for all finite sequences $(x_i)_{1 \leq i \leq n}$ in X , we have

$$\pi_2(v) \leq C \left\| \sum_{i=1}^n \varphi_i x_i \right\|_{L_1^{2n}(X)}, \quad (2.19)$$

where $v : l_2^n \rightarrow X$ is the operator defined by $v(e_i) = x_i$ for all $1 \leq i \leq n$.

We denote by

$$K_2^{\text{strong}}(X) = \inf \{C : (2.19) \text{ holds } \forall (x_i)_{1 \leq i \leq n}, n \geq 1\}. \quad (2.20)$$

COROLLARY 2.5. *Let X be a Banach space and let C be a positive constant. The following assertions are equivalent.*

- (i) *The space X^* is of cotype strongly 2-Kašin and $K_2^{\text{strong}}(X^*) \leq C$.*
- (ii) *For all integers n and any $u : l_2^n \rightarrow X$, u admits an extension \tilde{u} to L_1^{2n} such that $\tilde{u}/E_n = u\beta_n^{-1}$ and $\|\tilde{u}\| \leq C\pi_2(u^*)$.*

6 Weak Grothendieck's theorem

Proof. Fixed n in \mathbb{N} , let $E = \{\sum_{i=1}^n \varphi_i \xi_i, (\xi_i)_{1 \leq i \leq n} \subset X^*\}$ which is a closed subspace of $L_1^{2n}(X^*)$. We now define the operators

$$\begin{aligned} T : E &\longrightarrow \pi_2(l_2^n, X^*), \\ \sum_{i=1}^n \varphi_i \xi_i &\longmapsto v, \end{aligned} \quad (2.21)$$

where $v : l_2^n \rightarrow X^*$ defined by $v(e_i) = \xi_i$.

We have

$$\left\| T \left(\sum_{i=1}^n \varphi_i \xi_i \right) \right\| = \pi_2(v) \leq C \left\| \sum_{i=1}^n \varphi_i \xi_i \right\|. \quad (2.22)$$

By duality, we obtain

$$\begin{aligned} T^* : \pi_2(X^*, l_2^n) &\longrightarrow \frac{L_\infty^{2n}(X)}{E^\perp}, \\ w &\longmapsto \sum_{i=1}^n x_i \varphi_i + E^\perp, \end{aligned} \quad (2.23)$$

where $w : X^* \rightarrow l_2^n$ is a linear operator defined by $w(\xi) = \langle x_i, \xi \rangle$.

Let $u(e_i) = x_i$. We have

$$\inf_{R \in E^\perp} \left\| \sum_{i=1}^n x_i \varphi_i + R \right\|_{L_\infty^{2n}(X)} \leq C \pi_2(u^*). \quad (2.24)$$

We conclude directly by using (2.18). \square

Remark 2.6. Let X be a Banach space. If X has Gaussian (resp., Rademacher) cotype 2, then (2.19) holds with (g_i) (resp., (r_i)) and conversely. The space X is of cotype strongly 2-Kašin implies that X is of cotype 2-Kašin. We do not know if the converse is true.

Let us introduce the following definition.

Definition 2.7. Let X be a Banach space. Say that X satisfies weak Grothendieck's theorem if there is a positive constant C such that for all n in \mathbb{N} and any linear operator u from X into l_2^n , there exists

$$\pi_2(u) \leq C \|u\|. \quad (2.25)$$

Remark 2.8. (1) X satisfies W.G.T. if and only if X^{**} satisfies WGT.

(2) L_1 and L_∞ verify weak Grothendieck's theorem. The spaces S_1 (see below) and $B(l_2)$ (see [8, Corollary 4.2]) fail this.

(3) The classical definition is let X be a Banach space. We will say that X satisfies Grothendieck's theorem if there is a constant C such that, for any linear operator u from X into a Hilbert space H , we have

$$\pi_1(u) \leq C \|u\|. \quad (2.26)$$

(4) We can replace H by l_2^n for any integer n (i.e., there is a constant C such that for any integer n and any $u : X \rightarrow l_2^n$ we have $\pi_1(u) \leq C\|u\|$). Also, this is equivalent to the dual property (i.e., there is a constant C' such that for every linear operator from X^* into an L_1 -space, we have $\pi_2(u) \leq C'\|u\|$). GT implies WGT. If X is of (classical) cotype 2, then we have equivalence between GT and WGT because $\pi_p(X, Y) = \pi_2(X, Y)$ for any Banach space Y and for all $p \leq 2$ (see [7]).

(5) The space L_1 verifies Grothendieck's theorem. In [2] Bourgain proved that L_1/H_1 is of cotype 2 and verifies Grothendieck's theorem (L_1 is the L_1 -space relative to the circle group and H_1 the subspace of L_1 spanned by all functions $\{e^{int}, n \geq 0\}$).

(6) Suppose that X is a subspace of $C(K)$ and that $C(K)/X$ is reflexive. Then every operator with domain X and range a cotype 2 space is 2-summing [6, 11]. As corollary, let X be a reflexive subspace of an L_1 . Then, every operator $u : L_1/X \rightarrow l_2$ is 1-summing.

(7) For any Banach E of cotype 2, Pisier has constructed in [12] a Banach space X which contains isometrically E such that, X and X^* are both of cotype 2 and verify Grothendieck's theorem.

3. S_p fails WGT for all $1 \leq p \leq 2$

We recall (see [14]) the noncommutative analogues of l_p which is the Schatten class S_p . Let $0 < p < \infty$. We will denote by $B(l_2)$ the space of all bounded linear operators $u : l_2 \rightarrow l_2$ and by S_p the subspace of all compact operators such that $\text{tr } |u|^p < \infty$ (where $|u| = (uu^*)^{1/2}$). We equip it with the norm if $1 \leq p < \infty$ and the p -norm if $0 < p < 1$:

$$\|u\|_p = (\text{tr } |u|^p)^{1/p} \quad (3.1)$$

for which it becomes a Banach space if $1 \leq p < \infty$ and a quasi-Banach if $0 < p < 1$. If $p = \infty$, S_∞ is the subspace of all compact operators on l_2 equipped with operator norm. We have $(S_p)^* = S_q$ for $1 < p \leq \infty$ and $1/p + 1/q = 1$, and also $S_1^* = B(l_2)$. We do not know if the Schatten spaces S_p are of the same cotype Kašin as the usual l_p -spaces for $1 \leq p \leq 2$.

Finally, we denote by S_p^n and $B(l_2^n)$ the finite dimensional version of S_p and $B(l_2)$, respectively.

Let $0 < p \leq q \leq \infty$. We have for $u \in B(l_2^n)$,

$$\|u\|_q \leq \|u\|_p \leq n^{1/p-1/q} \|u\|_q. \quad (3.2)$$

Let R_n denote the subspace of S_p^n consisting of all $n \times n$ matrices u such that $u_{i,j} = 0$ when $i \neq 1$ (first row matrices). Then $a = uu^*$ is the matrix with $a_{1,1} = \sum_{j=1}^n |u_{1,j}|^2 = \|u\|_2^2$ and $a_{i,j} = 0$ when $(i,j) \neq (1,1)$. Hence $|u|$ is the rank one operator $\|u\|_2 e_1 \otimes e_1$. Its norm in all spaces S_p^n , $0 < p \leq \infty$ is equal to $\|u\|_2$. In particular R_n equipped with the S_p^n -norm is isometric to l_2^n . We denote by p_n the natural projection from S_p^n into R_n defined by $p_n(u) = v$ such that $v_{1,j} = u_{1,j}$ for $1 \leq j \leq n$. We have $\|p_n\| \leq 1$.

The proposition to be proved now is the finite dimensional version of the theorem of extension.

PROPOSITION 3.1. *Suppose that for some $p > 0$, there exists a constant C_p such that for every n and every linear operator u from E_n to S_p^n , there is an extension $\tilde{u} \in B(L_1^{2n}, S_p^n)$ of u with*

$\|\tilde{u}\| \leq C_p \|u\|$. Then

$$C_p \geq C\sqrt{n}, \quad (3.3)$$

where C is an absolute constant.

Proof. Let u_n be the operator sending the n vector basis of E_n to the n vector basis of R_n ($u_n(\varphi_i) = e_{1,i}, 1 \leq i \leq n$). This operator is an isomorphism, by the above remark and (2.6). We have $\|u_n\| \leq B_1$ and $\|u_n^{-1}\| \leq A_1$. Let \tilde{u}_n be an extension of u_n to an operator from L_1^{2n} to S_p^n , with $\|\tilde{u}_n\| \leq C_p \|u_n\|$. Consider now the following commutative diagram:

$$\begin{array}{ccccc} L_1^{2n} & \xrightarrow{\tilde{u}_n} & S_p^n & & \\ \uparrow e_n & & \downarrow p_n & & \\ E_n & \xrightarrow{u_n} & R_n & \xrightarrow{u_n^{-1}} & E_n \end{array} \quad (3.4)$$

Let $q_n = u_n^{-1} p_n \tilde{u}_n$. Then q_n is a projection from L_1^{2n} to E_n . Since E_n is $A_1 B_1$ -isomorphic to l_2^n (Theorem 2.1), we get by Grothendieck's theorem [4] that q_n is 1-summing with $\pi_1(q_n) \leq A_1 K_G \|p_n \tilde{u}_n\|$. Restricting q_n to E_n we obtain for the identity i_n of E_n the estimation

$$\sqrt{n} = \pi_2(i_n) \leq \pi_2(q_n) \leq \pi_1(q_n) \leq A_1 K_G \|\tilde{u}_n\| \leq A_1 K_G C_p \|u_n\| \leq A_1 B_1 K_G C_p. \quad (3.5)$$

This completes the proof. \square

Let now \mathcal{B}_n be the σ -algebra on $[0, 1]$ generated by the Rademacher functions $\{r_1, \dots, r_n\}$ ($r_n(t) = \text{sign}(\sin 2^n \pi t)$). The space $L_p([0, 1], \mathcal{B}_n, \nu)$, where ν is the Lebesgue measure on $[0, 1]$, is isometric to $L_p^{2^n}$.

We denote by G (resp., G_n) the closed linear subspace in $L_1([0, 1], \nu)$ (resp., $L_1^{2^n}$) of the Rademacher functions $\{r_n\}_{n \in \mathbb{N}}$ (resp., $\{r_i, 1 \leq i \leq n\}$). Let $g : G \rightarrow L_1([0, 1], \nu)$ (resp., $g_n : G_n \rightarrow L_1^{2^n}$) be the isometric embedding. By Khinchine's inequalities, there are positive constants A'_1 and B'_1 such that for every (a_n) in l_2 we have

$$A'_1 \left(\sum_{n \geq 1} |a_n|^2 \right)^{1/2} \leq \left(\int_{[0,1]} \left| \sum_{n \geq 1} a_n r_n(t) \right| d\nu \leq B'_1 \left(\sum_{n \geq 1} |a_n|^2 \right)^{1/2}. \quad (3.6)$$

Hence G (resp., G_n) is isomorphic to l_2 (resp., l_2^n). We will denote by $\alpha : l_2 \rightarrow G$ (resp., $\alpha_n : l_2^n \rightarrow G_n$) the isomorphism which maps e_i onto r_i . We have $\|\alpha\| \leq B'_1$, $\|\alpha^{-1}\| \leq A'_1$, and also the same for α_n .

PROPOSITION 3.2. *Suppose that for some $p > 0$, there exists a constant C_p such that for every n and every linear operator u from G_n to S_p^n there is an extension $\tilde{u} \in B(L_1^{2^n}, S_p^n)$ of u with $\|\tilde{u}\| \leq C_p \|u\|$. Then*

$$C_p \geq C\sqrt{n}, \quad (3.7)$$

where C is an absolute constant.

Proof. The same proof as in Proposition 3.1. \square

THEOREM 3.3. *Let $0 < p \leq \infty$. Let $u : G \rightarrow S_p$ be a compact linear operator. In general, there is no continuous linear operator \tilde{u} extending u to $L_1([0, 1], \nu)$.*

Proof. Suppose that for any compact linear operator $u : G \rightarrow S_p$ there is a bounded linear operator $\tilde{u} : L_1([0, 1], \nu) \rightarrow S_p$ extending u . It follows from the open mapping theorem that there is an absolute constant C_p such that

$$\|\tilde{u}\| \leq C_p \|u\| \quad (3.8)$$

for any u . This implies by Proposition 3.2 that $C_p \geq C\sqrt{n}$ for any integer n . This is impossible when n is large enough. \square

THEOREM 3.4. *Let $0 < p \leq \infty$. Let $u : E_0 \rightarrow S_p$ be a compact linear operator. In general, there is no continuous linear operator \tilde{u} extending u .*

Proof. Using the same proof as in Proposition 3.2 (we take E_0^n instead of G_n) and Theorem 3.3, we show that the extension property concerning $(L_1([0, 1], \nu), E_0)$ fails for all $0 < p \leq \infty$. \square

The following result shows that space S_p fails GT.

THEOREM 3.5. *The space S_p fails GT for all $1 \leq p \leq \infty$ and consequently WGT for $1 \leq p \leq 2$.*

Proof. Consider the following diagram:

$$R_n \xrightarrow{i_n} S_p^n \xrightarrow{p_n} R_n, \quad (3.9)$$

where i_n is the canonical injection. We have $\text{id}_{R_n} = p_n \circ i_n$. Since $\sqrt{n} \leq \pi_1(\text{id}_{R_n}) \leq \pi_1(p_n)$ and $\|p_n\| \leq 1$, hence S_p fails GT for all $1 \leq p \leq \infty$. As S_p is of cotype 2 for $1 \leq p \leq 2$ then, by one result of Maurey, we have $\pi_1(p_n) \leq C\pi_2(p_n)$ for some constant C . This implies the proof. \square

Remark 3.6. The space $B(l_2)$ fails weak Grothendieck's theorem because by [8, Corollary 4.2] we have $\pi_2(B(l_2), l_2) \neq B(B(l_2), l_2)$.

4. Characterization of spaces which satisfy WGT

We start this section by recalling some notations and facts. We denote by $l_p^\omega(X)$ (resp., $l_p^{m\omega}(X)$) the space of all sequences (x_i) (resp., $(x_i)_{1 \leq i \leq n}$) in X with the norm

$$\begin{aligned} \|(x_i)\|_{l_p^\omega(X)} &= \sup_{\|\xi\|_{X^*}=1} \left(\sum_1^\infty |\langle x_i, \xi \rangle|^p \right)^{1/p} < \infty, \\ \left(\text{resp., } \|(x_i)\|_{l_p^{m\omega}(X)} &= \sup_{\|\xi\|_{X^*}=1} \left(\sum_1^n |\langle x_i, \xi \rangle|^p \right)^{1/p} \right). \end{aligned} \quad (4.1)$$

We know (see [3]) that $l_p(X) = l_p^\omega(X)$ for some $1 \leq p < \infty$ if and only if $\dim(X)$ is finite. If $p = \infty$, we have $l_\infty(X) = l_\infty^\omega(X)$. We have also if $1 < p \leq \infty$, $l_p^\omega(X) \equiv B(l_{p^*}, X)$, and $l_1^\omega(X) \equiv B(c_0, X)$ isometrically (where p^* is the conjugate of p , i.e., $1/p + 1/p^* = 1$). In other words, let $v: l_{p^*} \rightarrow X$ be a linear operator such that $v(e_i) = x_i$ (namely, $v = \sum_1^\infty e_j \otimes x_j$, e_j denotes the unit vector basis of l_{p^*}), then

$$\|v\| = \|(x_i)\|_{l_p^\omega(X)} = \left\| \sum_1^\infty e_j \otimes x_j \right\|_{l_p \otimes_\epsilon X}. \quad (4.2)$$

We prove in the following theorem that the spaces which satisfy WGT and which happen to be also of cotype strongly 2-Kašin can be characterized by an extension property.

THEOREM 4.1. *The following properties of a Banach space X are equivalent:*

- (i) *the space X^* is of cotype strongly 2-Kašin and verifies WGT;*
- (ii) *there is a positive constant C such that for every $n \in \mathbb{N}$ and every $u: E_n \rightarrow X$, then u admits an extension $\tilde{u}: L_1^{2n} \rightarrow X$ such that $\tilde{u}/E_n = u$ and $\|\tilde{u}\| \leq C\|u\|$.*

Proof. We prove that (ii) \Rightarrow (i). Let $v: l_2^n \rightarrow X$ be a linear operator. Consider $u = v\beta_n^{-1}: E_n \rightarrow X$, then u admits an extension $\tilde{u}: L_1^{2n} \rightarrow X$ such that

$$\|\tilde{u}\| \leq C\|u\| \leq C\|\beta_n^{-1}\|\|v\| \leq C/A_1\pi_2(v^*). \quad (4.3)$$

From Corollary 2.5, we obtain that X^* is of cotype strongly 2-Kašin and $K_2^{\text{strong}}(X^*) \leq C/A_1$. Let now $u: X^* \rightarrow l_2^n$ be an operator. First, we notice that $B(l_2^n, X^{**}) \equiv B(l_2^n, X)^{**} \equiv B(X^*, l_2^n)$ isometrically. Since $u: X^* \rightarrow l_2^n$ is in $B(l_2^n, X)^{**}$, then by Goldstine's theorem, there is a net of operators $u_i^*: X^* \rightarrow l_2^n$ which are w^* -continuous with $\|u_i\| \leq \|u\|$ for all i and $\{u_i^*\}$ converges to u in w^* -topology of $B(l_2^n, X)^{**}$. As u_i^* is 2-summing this implies that u is 2-summing and $\pi_2(u) = \lim_i \pi_2(u_i^*)$. Indeed,

$$\begin{aligned} \pi_2(u) &= \sup \{ \text{Tr}(uv), v: l_2^n \rightarrow X^{***} \pi_2(v) \leq 1 \} \\ &= \sup \{ \lim_i \text{Tr}(u_i^* v), v: l_2^n \rightarrow X^{***} \pi_2(v) \leq 1 \} \\ &= \lim_i \sup \{ \text{Tr}(u_i^* v), v: l_2^n \rightarrow X^{***} \pi_2(v) \leq 1 \} \\ &= \lim_i \pi_2(u_i^*). \end{aligned} \quad (4.4)$$

Let us consider the following commutative diagram:

$$\begin{array}{ccc} L_1^{2n} & & \\ \uparrow e_n & \searrow \tilde{u}_i & \\ E_n & \xrightarrow{\beta_n^{-1}} l_2^n & \xrightarrow{u_i} X \end{array} \quad (4.5)$$

by duality, we have

$$\begin{array}{ccccc}
 & & & L_\infty^{2n} & \\
 & & \nearrow \tilde{u}_i^* & \downarrow \mathbf{e}_n^* & \\
 X^* & \xrightarrow{u_i^*} & l_2^n & \xrightarrow{(\beta_n^{-1})^*} & E_n^*
 \end{array} \quad (4.6)$$

hence

$$\begin{aligned}
 \pi_2(u_i^*) &= \pi_2(\beta_n^*(\beta_n^{-1})^* u_i^*) \leq \|\beta_n^*\| \pi_2((\beta_n^{-1})^* u_i^*) \\
 &\leq \|\beta_n^*\| \|\tilde{u}_i^*\| \pi_2(\mathbf{e}_n^*) \leq \|\beta_n^*\| \|\beta_n^{-1}\| \|u_i\| \pi_2(\mathbf{e}_n^*) \\
 &\leq A_1^{-1} B_1 \|u_i\| \pi_2(\mathbf{e}_n^*).
 \end{aligned} \quad (4.7)$$

Thus

$$\lim_i \pi_2(u_i^*) \leq A_1^{-1} B_1 \pi_2(\mathbf{e}_n^*) \lim_i \|u_i\| \leq A_1^{-1} B_1 \pi_2(\mathbf{e}_n^*) \|u\|. \quad (4.8)$$

Consequently

$$\pi_2(u) \leq A_1^{-1} B_1 \pi_2(\mathbf{e}_n^*) \|u\|. \quad (4.9)$$

This shows that X has WGT because the numbers $\pi_2(\mathbf{e}_n^*)$ are uniformly bounded by Maurey's theorem [7].

(i) \Rightarrow (ii). The space X^* is of cotype strongly 2-Kašin which implies by Corollary 2.5 that for any $u : l_2^n \rightarrow X$, u admits an extension \tilde{u} to L_1^{2n} such that $\tilde{u}/E_n = u\beta_n^{-1}$ and $\|\tilde{u}\| \leq K_2^{\text{strong}}(X^*) \pi_2(u^*)$. As X^* verifies WGT, then $\pi_2(u^*) \leq C' \|u\|$ and hence

$$\|\tilde{u}\| \leq C' K_2(X^*) \|u\| \leq C \|u\| \quad (C = C' K_2(X^*)) \quad (4.10)$$

which gives the extension. \square

We end this paper by the following remark.

Remark 4.2. We do not know if S_p for $1 \leq p \leq 2$ is of cotype 2-Kašin.

Acknowledgment

The author is very grateful to the referees for pointing out some mistakes in the first version and for several valuable suggestions and comments which improved the paper.

References

- [1] G. W. Anderson, *Integral Kašin splittings*, Israel Journal of Mathematics **138** (2003), 139–156.
- [2] J. Bourgain, *New Banach space properties of the disc algebra and H^∞* , Acta Mathematica **152** (1984), no. 1-2, 1–48.

- [3] J. Diestel, H. Jarchow, and A. Tonge, *Absolutely Summing Operators*, Cambridge Studies in Advanced Mathematics, vol. 43, Cambridge University Press, Cambridge, 1995.
- [4] A. Grothendieck, *Résumé de la théorie métrique des produits tensoriels topologiques*, Boletín de la Sociedad Matemática São Paulo **8** (1956), 1–79.
- [5] B. S. Kašin, *Sections of finite dimensional sets closes of smooths functions*, Izvestiya Akademii Nauk SSSR. Seriya Matematicheskaya **41** (1977), 334–351.
- [6] S. V. Kisliakov, *On spaces with “small” annihilators*, Seminarov Leningradskogo Otdeleniya Matematicheskogo Instituta imeni V. A. Steklova Akademii Nauk SSSR (LOMI) **73** (1978), 91–101.
- [7] B. Maurey, *Théorèmes de factorisation pour les opérateurs linéaires à valeurs dans les espaces L^p* , Astérisque, no. 11, Société Mathématique de France, Paris, 1974.
- [8] L. Mezrag, *Comparison of non-commutative 2- and p -summing operators from $B(l_2)$ into OH* , Zeitschrift für Analysis und ihre Anwendungen **21** (2002), no. 3, 709–717.
- [9] ———, *Factorization of operator valued L_p for $0 \leq p < 1$* , Mathematische Nachrichten **266** (2004), no. 1, 60–67.
- [10] ———, *On the L_1^n -extension properties*, Quaestiones Mathematicae **27** (2004), no. 3, 297–309.
- [11] G. Pisier, *Une nouvelle classe d'espaces de Banach vérifiant le théorème de Grothendieck*, Annales de l'Institut Fourier (Grenoble) **28** (1978), no. 1, x, 69–90.
- [12] ———, *Counterexamples to a conjecture of Grothendieck*, Acta Mathematica **151** (1983), no. 1, 181–208.
- [13] ———, *Factorization of Linear Operators and Geometry of Banach Spaces*, CBMS Regional Conference Series in Mathematics, vol. 60, American Mathematical Society, Rhode Island, 1986, reprinted with corrections 1987.
- [14] J. R. Retherford, *Hilbert Space: Compact Operators and the Trace Theorem*, London Mathematical Society Student Texts, vol. 27, Cambridge University Press, Cambridge, 1993.
- [15] G. Schechtman, *Special orthogonal splittings of L_1^{2k}* , Israel Journal of Mathematics **139** (2004), 337–347.
- [16] N. Tomczak-Jaegermann, *The moduli of smoothness and convexity and the Rademacher averages of trace classes S_p ($1 \leq p < \infty$)*, Studia Mathematica **50** (1974), 163–182.

Lahcène Mezrag: Department of Mathematics, M'sila University, P.O. Box 166, M'sila 28000, Algeria
 E-mail address: lmezrag@yahoo.fr

