

Research Article

The Modified Negative Decision Number in Graphs

Changping Wang

Department of Mathematics, Ryerson University, Toronto, ON, Canada M5B 2K3

Correspondence should be addressed to Changping Wang, cpwang@ryerson.ca

Received 14 December 2010; Accepted 11 January 2011

Academic Editor: Dalibor Froncek

Copyright © 2011 Changping Wang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A mapping $x : V \rightarrow \{-1, 1\}$ is called *negative* if $\sum_{u \in N[v]} x(u) \leq 1$ for every $v \in V$. The maximum of the values of $\sum_{v \in V} x(v)$ taken over all negative mappings x , is called the *modified negative decision number* and is denoted by $\beta'_D(G)$. In this paper, several sharp upper bounds of this number for a general graph are presented. Exact values of these numbers for cycles, paths, cliques and bicliques are found.

1. Introduction

Let $G = (V, E)$ be a graph with vertex set V and edge set E . For a vertex $v \in V$, the *open neighbourhood* of v in G is $N(v) = \{u \in V \mid uv \in E\}$, and $N[v] = N(v) \cup \{v\}$ is its *closed neighbourhood*. For a graph G , and a subset $S \subseteq V$, we let $\deg_S(v)$ denote the number of vertices in S joined to v . In particular, $\deg_V(v) = \deg(v)$, the degree of v in G . For disjoint subsets S and T of vertices, we use $E(S, T)$ for the set of edges between S and T , and let $e(S, T) = |E(S, T)|$. Let $G[S]$ denote the subgraph of G induced by S . For an integer $k \geq 3$, the bipartite graph $K_{1,k}$ is called a *star*, and the vertex of degree k is called the *central vertex*. Let $x : V \rightarrow \mathbb{R}$ be a real-valued function. For convenience, we write $x(S)$ for $\sum_{v \in S} x(v)$ for $S \subseteq V$.

In [1], we initiated the study of the *negative decision number* in a graph. A function $x : V(G) \rightarrow \{-1, 1\}$ is called a *bad function* of G if $x(N(v)) \leq 1$ for every $v \in V(G)$. The maximum of the values of $x(V(G))$, taken over all bad functions f , is called the *negative decision number* and is denoted by $\beta_D(G)$.

The motivation for studying this parameter may be explained from a modelling perspective. For instance, by assigning the values -1 or 1 to the vertices of a graph, one can model networks of people in which global decisions must be made (e.g., positive or negative responses). In certain circumstances, a positive decision can be made only if there are significantly more people voting for than those voting against. We assume that each

individual has one vote, and each has an initial opinion. We assign 1 to vertices (individuals) which have a positive opinion and -1 to vertices which have a negative opinion. A voter votes “good” if there are at least two more vertices in its open neighbourhood with positive opinion than with negative opinion, otherwise the vote is “bad”. We seek an assignment of opinions that guarantee a unanimous decision; namely, for which every vertex votes “bad”. Such an assignment of opinions is called a uniformly negative assignment. Among all uniformly negative assignments of opinions, we are particularly interested in the minimum number of vertices (individuals) which have a negative opinion. The negative decision number is the maximum possible sum of all opinions, 1 for a positive opinion and -1 for a negative opinion, in a uniformly negative assignment of opinions. The negative decision number corresponds to the minimum number of individuals who can have negative opinions and in doing so force every individual to vote bad.

In the present paper, we study a variation of the negative decision number in a graph. A mapping $x : V \rightarrow \{-1, 1\}$ is called *negative* if $\sum_{u \in N[v]} x(u) \leq 1$ for every $v \in V$. The maximum of the values of $\sum_{v \in V} x(v)$, taken over all negative mappings x , is called the *modified negative decision number* and is denoted by $\beta'_D(G)$.

The parameter β'_D differs significantly from β_D . For instance, for a cycle C_n of order n , $\beta_D(C_n) \in \{-2, -1, 0\}$ ([1] see Theorem 5), while $\beta'_D(C_n) \in \{\lfloor n/3 \rfloor, \lfloor n/3 - 1 \rfloor\}$ (see Corollary 2.6).

Throughout this paper, if x is a negative mapping of G , then we let P and Q denote the sets of those vertices of G which are assigned (under x) the values $+1$ and -1 , respectively, and we let $p = |P|$ and $q = |Q|$. Hence, $x(V) = p - q$.

All graphs considered in this paper are simple, finite, and undirected. For a general reference on graph theory, the reader is referred to [2, 3]. Notice that $\beta'_D(G \cup H) = \beta'_D(G) + \beta'_D(H)$ for two disjoint graphs G and H . Hence, we assume that all graphs are connected in this paper.

In the next section, we present several sharp upper bounds on the modified negative decision number for a general graph. We also establish sharp upper bounds on this number for bipartite graphs and regular graphs. Exact values of these numbers for cycles, paths, cliques, and bicliques are found.

2. Main Results

In this section, we investigate the modified negative decision number of a graph. We first present an upper bound on this number for a general graph in terms of its order.


For this purpose, we define a family \mathcal{F} of graphs as follows. Let $F_3 = K_{1,2}$, and for $k \geq 4$, let F_k be the graph obtained from the disjoint union of k stars $K_{1,k+1}$ by adding all possible edges between the central vertices of the k stars. See Figure 1 for an example of F_4 . Let $\mathcal{F} = \{F_k \mid k \geq 3\}$.

Theorem 2.1. *If G is a graph of order $n \geq 2$, then*

$$\beta'_D(G) \leq n + 2 - 2\sqrt{n+1}, \quad (2.1)$$

and this bound is sharp.

Proof. Let x be a negative mapping such that $\beta'_D(G) = x(V)$. Then $\beta'_D(G) = |P| - |Q| = n - 2q$. Note that every vertex in P must be joined to at least one vertex in Q . By the pigeonhole

Figure 1: A copy of the graph F_4 .

principle, at least one vertex v in Q is joined to at least $|P|/|Q| = (n - q)/q$ vertices in P . Hence,

$$1 \geq x(N[v]) \geq -|Q| + \frac{n - q}{q} = -q - 1 + \frac{n}{q}. \quad (2.2)$$

That is,

$$q^2 + 2q - n \geq 0. \quad (2.3)$$

Solving the above inequality, we obtain that

$$q \geq -1 + \sqrt{n + 1}. \quad (2.4)$$

Thus $\beta'_D(G) = n - 2q \leq n + 2 - 2\sqrt{n + 1}$.

To see this bound is sharp, let $G \in \mathcal{F}$. Thus, $G = F_k$ for some $k \geq 4$. Hence, G has order $n = k(k + 2)$, and so $k = \sqrt{n + 1} - 1$. Assigning the value -1 to each of the k central vertices of the stars, and $+1$ to all other vertices, we define a negative mapping x of G satisfying $x(V) = k^2 = n - 2k = n + 2 - 2\sqrt{n + 1}$. Thus, $\beta'_D(G) \geq n + 2 - 2\sqrt{n + 1}$. It follows that $\beta'_D(G) = n + 2 - 2\sqrt{n + 1}$. \square

Next we give an upper bound of the modified negative decision number for a general graph in terms of its order and size.

Theorem 2.2. *If G is a graph of order $n \geq 2$ and size m , then*

$$\beta'_D(G) \leq \frac{1}{5}(4m - n), \quad (2.5)$$

and this bound is sharp.

Proof. Let x be a negative mapping such that $\beta'_D(G) = x(V)$. Then $\beta'_D(G) = |P| - |Q| = n - 2q$. As every vertex in P must be joined to at least one vertex in Q , $e(P, Q) \geq p = n - q$. For each

vertex v of Q , $\deg_Q(v) \geq \deg_P(v) - 2$, and hence,

$$n - q \leq e(P, Q) = \sum_{v \in Q} \deg_P(v) \leq \sum_{v \in Q} (\deg_Q(v) + 2). \quad (2.6)$$

Namely,

$$n - q \leq 2|E(G[Q])| + 2q. \quad (2.7)$$

Hence,

$$|E(G[Q])| \geq \frac{n - 3q}{2}. \quad (2.8)$$

Thus, the total number of edges in G is

$$m \geq |E(G[Q])| + e(P, Q) \geq \frac{n - 3q}{2} + n - q. \quad (2.9)$$

So, $q \geq (3n - 2m)/5$. Hence,

$$\beta'_D(G) = n - 2q \leq \frac{1}{5}(4m - n). \quad (2.10)$$

To see this bound is sharp, let $G = F_k \in \mathcal{F}$ for some $k \geq 4$ and let x be the negative mapping of G defined in the proof of Theorem 2.1. As $x(V) = k^2$, $\beta'_D(G) \geq k^2$. On the other hand, as G has order $n = k(k + 2)$ and size $m = k(k + 1) + k(k - 1)/2$, $\beta'_D(G) \leq (1/5)(4m - n) = k^2$. Therefore, $\beta'_D(G) = (1/5)(4m - n)$. \square

In the following theorem, we establish an upper bound of the modified negative decision number for a bipartite graph in terms of its order and we characterize the graphs attaining this bound. We define a family \mathcal{H} of bipartite graphs as follows.


For $k \geq 1$, let H_k be the bipartite graph obtained from the disjoint union of $2k$ stars $K_{1,k+2}$ with centres $\{x_1, x_2, \dots, x_k, y_1, y_2, \dots, y_k\}$ by adding all edges of the type $x_i y_j$, $1 \leq i, j \leq k$. See Figure 2 for an example of H_2 . Let $\mathcal{H} = \{H_k \mid k \geq 1\}$.

Theorem 2.3. *If G is a bipartite graph of order $n \geq 2$, then*

$$\beta'_D(G) \leq n + 6 - 2\sqrt{2n + 9}. \quad (2.11)$$

The equality holds if and only if $G \in \mathcal{H}$.

Proof. Let x be a negative mapping of G such that $\beta'_D(G) = x(V)$. Let X and Y be the partite sets of G . Further, let X^+ and X^- be the sets of vertices in X that are assigned the value $+1$ and -1 (under x), respectively. Let Y^+ and Y^- be defined analogously. Then $P = X^+ \cup Y^+$ and $Q = X^- \cup Y^-$. For convenience, let $|X^+| = k$, $|X^-| = s$, $|Y^+| = l$ and $|Y^-| = t$. Hence, $\beta'_D(G) = k + l - s - t = n - 2(s + t)$.

Figure 2: A copy of the graph H_2 .

As every vertex in X^+ must be joined to at least one vertex in Y^- , by the pigeonhole principle, at least one vertex v in Y^- is joined to at least $|X^+|/|Y^-| = k/t$ vertices in X^+ . Hence,

$$1 \geq x(N[v]) \geq -|X^-| - 1 + \frac{|X^+|}{|Y^-|} = -s - 1 + \frac{k}{t}. \quad (2.12)$$

Namely,

$$t(s + 2) \geq k. \quad (2.13)$$

By a similar argument, one may show that

$$s(t + 2) \geq l. \quad (2.14)$$

Adding (2.13) and (2.14), we have that

$$2st + 2(s + t) \geq k + l. \quad (2.15)$$

Thus,

$$\begin{aligned} n &= k + l + s + t \\ &\leq 2st + 3(s + t) \\ &\leq \frac{1}{2}(s + t)^2 + 3(s + t). \end{aligned} \quad (2.16)$$

Solving the above inequality for $s + t$, we obtain that

$$s + t \geq -3 + \sqrt{2n + 9}. \quad (2.17)$$

Thus, $\beta'_D(G) = n - 2(s + t) \leq n + 6 - 2\sqrt{2n + 9}$.

If $\beta'_D(G) = n + 6 - 2\sqrt{2n+9}$, then by the above analysis, we have $s = t$ and $k = l = s(s + 2)$. Moreover, each vertex of X^+ (resp., Y^+) has degree 1 and is joined to one vertex of Y^- (resp., X^-), while each vertex of Y^- is joined to all vertices of X^- and $s + 2$ vertices of X^+ and each vertex of X^- is joined to all vertices of Y^- and $s + 2$ vertices of Y^+ . Thus, $\beta'_D(G) = n + 6 - 2\sqrt{2n+9}$ implies that $G \in \mathcal{H}$.

If $G \in \mathcal{H}$, then $G = H_k$ for some $k \geq 1$. Hence, G has order $n = 2k(k + 3)$, and so $k = (1/2)(\sqrt{2n+9} - 3)$. Assigning to the k central vertices of the stars the value -1 , and to all other vertices $+1$, we define a negative mapping x of G satisfying $x(V) = 2k(k + 1) = n + 6 - 2\sqrt{2n+9}$. Hence, $\beta'_D(G) \geq n + 6 - 2\sqrt{2n+9}$. It follows that $\beta'_D(G) = n + 6 - 2\sqrt{2n+9}$. \square

Finally, we present an upper bound of the modified negative decision number for a regular graph in terms of its order in Theorem 2.5. We encourage the reader to verify our next theorem first.

Theorem 2.4. *For any integer $n \geq 2$, one has*

$$\beta'_D(K_n) = \begin{cases} 0 & \text{for } n \text{ even;} \\ 1 & \text{for } n \text{ odd.} \end{cases} \quad (2.18)$$

For a general k -regular graph, we have the following.

Theorem 2.5. *If G is k -regular graph of order n , then*

$$\beta'_D(G) \leq \begin{cases} \left\lfloor \frac{n}{k+1} \right\rfloor & \text{for } k \text{ even;} \\ 0 & \text{for } k \text{ odd.} \end{cases} \quad (2.19)$$

This bound is sharp.

Proof. Let x be any negative mapping of G . As G is a k -regular graph,

$$\sum_{v \in V} x(N[v]) = (k+1)x(V). \quad (2.20)$$

We discuss the following two cases.

Case 1. k is even. Since x is a negative mapping, for every vertex $v \in V$, $x(N[v]) \leq 1$. By (2.20), it follows that

$$(k+1)x(V) \leq n. \quad (2.21)$$

Hence, $\beta'_D(G) \leq n/(k+1)$.

Case 2. k is odd. In this case, $|N[v]|$ is even for every $v \in V$. So, for each $v \in V$, $x(N[v]) \leq 1$ implies that $x(N[v]) \leq 0$. Thus,

$$\sum_{v \in V} x(N[v]) \leq 0. \quad (2.22)$$

By (2.20), it follows that $(k+1)x(V) \leq 0$. Hence, $\beta'_D(G) \leq 0$.

The sharpness follows from Theorem 2.4. \square

Corollary 2.6. For any integer $n \geq 3$, one has

$$\beta'_D(C_n) = \begin{cases} \left\lfloor \frac{n}{3} \right\rfloor - 1 & n \equiv 1 \pmod{3}; \\ \left\lfloor \frac{n}{3} \right\rfloor & \text{otherwise.} \end{cases} \quad (2.23)$$

Proof. As C_n is a 2-regular graph, by Theorem 2.5, we have $\beta'_D(C_n) \leq n/3$. Let v_1, v_2, \dots, v_n be the n vertices of C_n in a clockwise order. We discuss the following three cases.

Case 1. $n = 3k$ for some integer k . To show that $\beta'_D(C_n) = k$, it suffices to show that there is a negative mapping x of C_n such that $x(V(C_n)) = k$. In fact, assigning $+1, +1, -1$ starting with v_1 clockwise, and repeating, we produce a negative mapping x of C_n satisfying $x(V(C_n)) = k$.

Case 2. $n = 3k + 2$ for some integer k . To show that $\beta'_D(C_n) = k$, it suffices to show that there is a negative mapping x of C_n such that $x(V(C_n)) = k$. In fact, if we assign $1, 1, -1, 1, -1$ to the vertices in a clockwise order when $n = 5$, or assign $1, 1, -1, \dots, 1, 1, -1, 1, -1$ starting with v_1 clockwise when $n > 5$, then we produce a negative mapping x of C_n satisfying $x(V(C_n)) = k$.

Case 3. $n = 3k + 1$ for some integer k . To show that $\beta'_D(C_n) \geq k - 1$, it suffices to show that there is a negative mapping x of C_n such that $x(V(C_n)) = k - 1$. In fact, if we assign $1, -1, 1, -1$ to the vertices in a clockwise order when $n = 4$, or assign $1, 1, -1, \dots, 1, 1, -1, -1$ starting with v_1 clockwise when $n > 4$, then we produce a negative mapping x of C_n satisfying $x(V(C_n)) = k - 1$.

To show $\beta'_D(C_n) \leq k - 1$, we let x be a negative mapping of C_n such that $x(V(C_n)) = \beta'_D(C_n)$. So $\beta'_D(C_n) = |P| - |Q| = 2p - n \leq n/3$. Hence, $p \leq 2k$, implying that $\beta'_D(C_n) = 2p - n \leq 2 \cdot 2k - (3k + 1) = k - 1$. \square

Corollary 2.7. For any integer $n \geq 2$, one has

$$\beta'_D(P_n) = \begin{cases} \left\lfloor \frac{n}{3} \right\rfloor - 1 & n \equiv 1 \pmod{3}; \\ \left\lfloor \frac{n}{3} \right\rfloor & \text{otherwise.} \end{cases} \quad (2.24)$$

Proof. It is easy to see that Corollary 2.7 is true for $2 \leq n \leq 4$. So we may assume that $n \geq 5$. Let v_1, v_2, \dots, v_n be the n vertices of P_n with end vertices v_1 and v_n . We only show that $\beta'_D(P_n) \leq n/3$. The rest of the proof is similar to that of Corollary 2.6, so is omitted.

Take any negative mapping x of P_n . For $i = 1$ or n , as $\deg(v_i) = 1$, we have that

$$x(N[v_i]) \leq 0. \quad (2.25)$$

For $1 < i < n$, $\deg(v_i) = 2$. Hence,

$$x(N[v_i]) \leq 1. \quad (2.26)$$

Thus,

$$\sum_{v \in V} x(N[v]) \leq n - 2. \quad (2.27)$$

Note that $\sum_{v \in V} x(N[v]) = 3x(V) - x(v_1) - x(v_n)$. So,

$$3x(V) - x(v_1) - x(v_n) \leq n - 2, \quad (2.28)$$

implying that $x(V) \leq (n - 2 + x(v_1) + x(v_n))/3 \leq n/3$. \square

For the biclique $K_{n,n}$, we have the following result.

Theorem 2.8. *For any integer $n \geq 2$, one has*

$$\beta'_D(K_{n,n}) = \begin{cases} 0 & \text{for } n \text{ even;} \\ -2 & \text{for } n \text{ odd.} \end{cases} \quad (2.29)$$

Proof. Let $X = \{u_1, u_2, \dots, u_n\}$ and $Y = \{v_1, v_2, \dots, v_n\}$ be the partite sets of $K_{n,n}$. As $K_{n,n}$ is an n -regular graph, by Theorem 2.5, $\beta'_D(K_{n,n}) \leq 2n/(n+1) < 2$. Hence, $\beta'_D(K_{n,n}) \leq 1$. As $K_{n,n}$ has order $2n$, which is even,

$$\beta'_D(K_{n,n}) \leq 0. \quad (2.30)$$

To prove the case when n is even, it suffices to show that there exists a negative mapping x of $K_{n,n}$ such that $x(V(K_{n,n})) = 0$. In fact, we can produce such x by assigning $x(v_i) = x(u_i) = (-1)^i$, $1 \leq i \leq n$.

Now we prove the case when n is odd. Let x be a negative mapping of $K_{n,n}$ such that $\beta'_D(K_{n,n}) = x(V(K_{n,n}))$. Let X^+, X^-, Y^+ , and Y^- be defined the same as in the proof of Theorem 2.3. Then $P = X^+ \cup Y^+$ and $Q = X^- \cup Y^-$. Let $|X^+| = k$, $|X^-| = s$, $|Y^+| = l$ and $|Y^-| = t$. Thus, $\beta'_D(K_{n,n}) = k + l - s - t = 2n - 2(s + t)$. As $\beta'_D(K_{n,n}) \leq 0$, by (2.30), we have that

$$k + l \leq s + t. \quad (2.31)$$

Notice that we may assume $k \leq s$ and $l \leq t$. If $k + l = s + t$, then $k = s$ and $l = t$ contradicting to the fact $k + s = l + t = n$, where n is odd. Hence, $\beta'_D(K_{n,n}) \leq -2$.

To show that $\beta'_D(K_{n,n}) = -2$, it suffices to show that there is a negative mapping x of $K_{n,n}$ such that $x(V(K_{n,n})) = -2$. In fact, we can produce such x by assigning $x(u_n) = -1$, $x(u_i) = (-1)^{i-1}$ for $1 \leq i \leq n-1$ and $x(v_j) = (-1)^j$, $1 \leq j \leq n$. \square

References

- [1] C. Wang, "The negative decision number in graphs," *The Australasian Journal of Combinatorics*, vol. 41, pp. 263–272, 2008.
- [2] G. Chartrand and L. Lesniak, *Graphs & Digraphs*, Chapman & Hall, Boca Raton, Fla, USA, 3rd edition, 2000.
- [3] D. B. West, *Introduction to Graph Theory*, Prentice Hall, Upper Saddle River, NJ, USA, 2nd edition, 2001.

