

Research Article

Certain Conditions for Starlikeness of Analytic Functions of Koebe Type

Saibah Siregar¹ and Maslina Darus²

¹ Faculty of Science and Biotechnology, Universiti Selangor, Bestari Jaya,
 45600 Selangor Darul Ehsan, Malaysia

² School of Mathematical Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia,
 Bangi, 43600 Selangor Darul Ehsan, Malaysia

Correspondence should be addressed to Maslina Darus, maslina@ukm.my

Received 1 May 2011; Revised 29 June 2011; Accepted 5 July 2011

Academic Editor: A. Zayed

Copyright © 2011 S. Siregar and M. Darus. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

For $\alpha \geq 0$, $\lambda > 0$, we consider the $M(\alpha, \lambda)_b$ of normalized analytic $\alpha - \lambda$ convex functions defined in the open unit disc \mathbb{U} . In this paper, we investigate the class $M(\alpha, \lambda)_b$, that is, $\operatorname{Re}\{(zf'_b(z)/f_b(z))[1 - \alpha + \alpha(1 - \lambda)(zf'_b(z)/f_b(z)) + \alpha\lambda(1 + (zf''_b(z)/f'_b(z)))]\} > 0$, with f_b is Koebe type, that is, $f_b(z) := z/(1 - z^n)^b$. The subordination result for the aforementioned class will be given. Further, by making use of Jack's Lemma as well as several differential and other inequalities, the authors derived sufficient conditions for starlikeness of the class $M(\alpha, \lambda)_b$ of n -fold symmetric analytic functions of Koebe type. Relevant connections of the results presented here with those given in the earlier works are also indicated.

1. Introduction

Let \mathcal{A} denote the class of normalized analytic functions of the form

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k, \quad (1.1)$$

which are analytic in the open unit disk $\mathbb{U} = \{z : |z| < 1\}$. Also, as usual, let

$$\begin{aligned} S^* &= \left\{ f : f \in \mathcal{A}, \operatorname{Re}\left(\frac{zf'(z)}{f(z)}\right) > 0, (z \in \mathbb{U}) \right\}, \\ K &= \left\{ f : f \in \mathcal{A}, \operatorname{Re}\left(1 + \frac{zf''(z)}{f'(z)}\right) > 0, (z \in \mathbb{U}) \right\}, \end{aligned} \quad (1.2)$$

be the familiar classes of *starlike functions* in \mathbb{U} and *convex functions* in \mathbb{U} , respectively.

If the functions f and g are analytic in \mathbb{U} , then we say that the function f is *subordinate* to g , or g is *superordinate* to f (written as $f < g$) if there exist a function $w(z)$ analytic \mathbb{U} , such that $|w(z)| < 1$ and $z \in \mathbb{U}$, and $w(0) = 0$ with $f(z) = g(w(z))$ in \mathbb{U} . If g is univalent in \mathbb{U} , then $f < g$ is equivalent to $f(0) = g(0)$ and $f(\mathbb{U}) \subseteq g(\mathbb{U})$.

Next, we let the $M(\alpha)$, that is,

$$M(\alpha) = \left\{ f(z) \in A : \operatorname{Re} \left\{ (1-\alpha) \frac{zf'(z)}{f(z)} + \alpha \left(1 + \frac{zf''(z)}{f'(z)} \right) \right\} > 0, (z \in \mathbb{U}) \right\}. \quad (1.3)$$

The class $M(\alpha)$ was first introduced by Mocanu [1], which was then known as the class of α convex (or α -starlike) functions. Later, Miller et al. [2] studied this class and showed that $M(\alpha)$ is a subclass of S^* for any real number α and also that $M(\alpha)$ is a subclass of K for $\alpha \geq 1$. We note that $M(0) = S^*$ and $M(1) = K$. Note also that Mocanu introduced $M(\alpha)$ with $f(z) \cdot f'(z)/0$. But Sakaguchi and Fukui [3] later showed that this condition was not needed.

Motivated essentially by the aforementioned earlier works, we aim here at deriving sufficient conditions for starlikeness of n -fold symmetric function f_b of the Koebe type, defined by

$$f_b(z) := \frac{z}{(1-z^n)^b} \quad (b \geq 0; n \in \mathbb{N} := \{1, 2, 3, \dots\}), \quad (1.4)$$

which obviously corresponds to the familiar Koebe function when $n = 1$ and $b = 2$.

Definition 1.1. A function $f(z)$ given by (1.1) is said to be in the class $M(\alpha, \lambda)_b$, for $\alpha \geq 0$, $\lambda > 0$, if the following conditions are satisfied:

$$\operatorname{Re} \left\{ \frac{zf'_b(z)}{f_b(z)} \left[1 - \alpha + \alpha(1-\lambda) \frac{zf'_b(z)}{f_b(z)} + \alpha\lambda \left(1 + \frac{zf''_b(z)}{f'_b(z)} \right) \right] \right\} > 0, \quad (z \in \mathbb{U}). \quad (1.5)$$

In this paper, we consider the class of functions $M(\alpha, \lambda)_b$.

In addition, in this paper, authors investigate the subordination of the class denoted by $M(\alpha, \lambda)_b$.

We have the following inclusion relationships:

$$(i) \quad M(0, \lambda)_1 \subset S^*$$

$$(ii) \quad M(\alpha, 1)_1 \subset \mathcal{H}(\alpha) \subset S^*, \text{ which } \mathcal{H}(\alpha) \text{ has studied by [4].}$$

The work of Siregar et al. [5] and Bansal and Raina [6] have also motivated us to come to these problems. Look also at [7, 8] for different studies.

The following result (popularly known as Jack's Lemma) will also be required in the derivation of our result (Theorem 4.1 below).

2. Preliminaries

Lemma 2.1 (see [9]). Let $q(z)$ be univalent in \mathbb{U} and let the function θ and ϕ be analytic in a domain D containing $q(\mathbb{U})$, with $\phi(w) \neq 0$ when $w \in q(\mathbb{U})$. Set

$$\begin{aligned} Q(z) &= \gamma z q'(z) \phi(q(z)), \quad \gamma > 0, \\ h(z) &= \theta(q(z)) + Q(z), \end{aligned} \quad (2.1)$$

and suppose that

- (i) $Q(z)$ is univalent and starlike in \mathbb{U} ;
- (ii) $\operatorname{Re}(zh'(z)/Q(z)) = \operatorname{Re}(\theta'(q(z))/\phi(q(z)) + zQ'(z)/Q(z)) > 0$, $z \in \mathbb{U}$.

If $p(z)$ is analytic in \mathbb{U} with $p(0) = q(0) = 1$, $p(\mathbb{U}) \subset D$, and

$$\theta(p(z)) + \gamma zp'(z) \phi(p(z)) < \theta(q(z)) + \gamma zq'(z) \phi(q(z)) = h(z), \quad (2.2)$$

then

$$p(z) < q(z), \quad (2.3)$$

and q is the best dominant.

Lemma 2.2 (see [10]). Let the (nonconstant) function $w(z)$ be analytic in \mathbb{U} such that $w(0) = 0$. If $|w(z)|$ attains its maximum value on circle $|z| = r < 1$ at a point $z_0 \in \mathbb{U}$, we have

$$z_0 w'(z_0) = kw(z_0), \quad (2.4)$$

where $k \geq 1$ is a real number.

3. The Subordination Result

Theorem 3.1. Let $f(z) \in A$ satisfy $f(z) \neq 0$ ($z \in \mathbb{U}$). Also, let the function $q(z)$ be univalent in \mathbb{U} , with $q(0) = 1$ and $q(z) \neq 0$, for $\lambda > 0$ and $\alpha \geq 0$, such that

$$\begin{aligned} \operatorname{Re}\left(1 + \frac{zf_b''(z)}{q'(z)}\right) &> 0 \quad (z \in \mathbb{U}), \\ \operatorname{Re}\left\{\lambda + 2q(z) + \frac{zf_b''(z)}{q'(z)}\right\} &> 0 \quad (z \in \mathbb{U}). \end{aligned} \quad (3.1)$$

If

$$\frac{zf_b'(z)}{f_b(z)} \left[1 - \alpha + \alpha(1 - \lambda) \frac{zf_b'(z)}{f_b(z)} + \alpha \lambda \left(1 + \frac{zf_b''(z)}{f_b'(z)} \right) \right] < h(z) \quad (z \in \mathbb{U}), \quad (3.2)$$

where

$$h(z) = \alpha [q(z)]^2 + (1 - \alpha)(q(z)) + \alpha \lambda z q'(z), \quad (3.3)$$

then

$$\frac{zf'_b(z)}{f_b(z)} < q(z) \quad (z \in \mathbb{U}), \quad (3.4)$$

and $q(z)$ is the best dominant of (3.2).

Proof. We first choose

$$p(z) = \frac{zf'_b(z)}{f_b(z)}, \quad \theta(w) = w(1 - \alpha + \alpha w), \quad \phi(w) = 1, \quad (3.5)$$

then $\theta(w)$ and $\phi(w)$ are analytic inside the domain \mathbb{D}^* , which contains $q(\mathbb{U})$, $q(0) = 1$, and $\phi(w) \neq 0$ when $w \in q(\mathbb{U})$.

Now, if we define the functions $Q(z)$ and $h(z)$ by

$$\begin{aligned} Q(z) &= \alpha \lambda z q'(z) \phi(q(z)) = \alpha \lambda z q'(z), \\ h(z) &= \theta(q(z)) + Q(z) = \alpha [q(z)]^2 + (1 - \alpha)(q(z)) + \alpha \lambda z q'(z), \end{aligned} \quad (3.6)$$

then it follows from (3.1) that $Q(z)$ is starlike in \mathbb{U} and

$$\operatorname{Re} \left(\frac{zh'(z)}{Q(z)} \right) > 0 \quad (z \in \mathbb{U}). \quad (3.7)$$

We also note that the function $p(z)$ is analytic in \mathbb{U} , with $p(0) = q(0) = 1$. Since $0 \notin p(\mathbb{U})$, therefore, $p(\mathbb{U}) \subset \mathbb{D}^*$, $\gamma = \alpha \lambda > 0$ and hence, the hypothesis of Lemma 2.1 are satisfied.

Applying Lemma 2.1, we find that

$$\begin{aligned} & \frac{zf'_b(z)}{f_b(z)} \left[1 - \alpha + \alpha(1 - \lambda) \frac{zf'_b(z)}{f_b(z)} + \alpha \lambda \left(1 + \frac{zf''_b(z)}{f'_b(z)} \right) \right] \\ &= \alpha [p(z)]^2 + (1 - \alpha)(p(z)) + \alpha \lambda z p'(z) = \theta(p(z)) + \alpha \lambda z p'(z) < h(z) \\ &= \alpha [q(z)]^2 + (1 - \alpha)(q(z)) + \alpha \lambda z q'(z) = \theta(q(z)) + \alpha \lambda z q'(z), \quad (z \in \mathbb{U}), \end{aligned} \quad (3.8)$$

which implies that

$$\frac{zf'_b(z)}{f_b(z)} < q(z) \quad (z \in \mathbb{U}), \quad (3.9)$$

and $q(z)$ is the best dominant of (3.2). □

4. The Properties of the Class $M(\alpha, \lambda)_b$

We begin by proving a stronger result than what we indicated in the preceding section.

Theorem 4.1. *Let the n -fold symmetric function $f_b(z)$, defined by (1.4), be analytic in \mathbb{U} , with*

$$\frac{f_b(z)}{z} \neq 0 \quad (z \in \mathbb{U}). \quad (4.1)$$

If $f_b(z)$ satisfies the inequality:

$$\operatorname{Re} \left\{ \frac{zf'_b(z)}{f_b(z)} \left[1 - \alpha + \alpha(1 - \lambda) \frac{zf'_b(z)}{f_b(z)} + \alpha\lambda \left(1 + \frac{zf''_b(z)}{f'_b(z)} \right) \right] \right\} > \left(1 - \frac{nb}{2} \right) \left(1 - \frac{\alpha nb}{2} \right) - \frac{\alpha\lambda nb}{4}, \quad (4.2)$$

($z \in \mathbb{U}$), then $f_b(z)$ is starlike in \mathbb{U} for

$$\alpha > 0, \quad \lambda > 0, \quad \left(\frac{\alpha\lambda + 2\alpha + 2 - \sqrt{\Delta}}{2\alpha} \leq nb \leq \frac{\alpha\lambda + 2\alpha + 2 + \sqrt{\Delta}}{2\alpha} \right), \quad (4.3)$$

$$(\Delta := \alpha^2(\lambda + 2)^2 + 4\alpha(\lambda - 2) + 4).$$

If $f_b(z)$ satisfies the inequality (4.2) with $\lambda = 1$, that is, if

$$\operatorname{Re} \left(\frac{\alpha z^2 f''_b(z)}{f_b(z)} + \frac{zf'_b(z)}{f_b(z)} \right) > -\frac{\alpha nb}{4} + \left(1 - \frac{nb}{2} \right) \left(1 - \frac{\alpha nb}{2} \right), \quad (z \in \mathbb{U}) \quad (4.4)$$

then $f_b(z)$ is starlike in \mathbb{U} for

$$\alpha > 0, \quad \frac{3\alpha + 2 - \sqrt{\Delta^*}}{2\alpha} \leq nb \leq \frac{3\alpha + 2 + \sqrt{\Delta^*}}{2\alpha}, \quad (\Delta^* := 9\alpha^2 - 4\alpha + 4). \quad (4.5)$$

Proof. Let $\alpha > 0$, $\lambda > 0$ and $f_b(z)$ satisfy the hypothesis of Theorem 4.1. We put

$$\frac{zf'_b(z)}{f_b(z)} = \frac{1 + (nb - 1)w(z)}{1 - w(z)}, \quad (4.6)$$

where $w(z)$ is analytic in \mathbb{U} , with

$$w(0) = 0, \quad w(z) \neq 1, \quad (z \in \mathbb{U}), \quad (4.7)$$

such that, we can write

$$1 + \frac{zf''_b(z)}{f'_b(z)} = \frac{nbzw'(z)}{[1 - w(z)][1 + (nb - 1)w(z)]} + \frac{1 + (nb - 1)w(z)}{1 - w(z)}, \quad (4.8)$$

which, in turn, implies that

$$\begin{aligned}
 & \frac{zf'_b(z)}{f_b(z)} \left[1 - \alpha + \alpha(1 - \lambda) \frac{zf'_b(z)}{f_b(z)} + \alpha\lambda \left(1 + \frac{zf''_b(z)}{f'_b(z)} \right) \right] \\
 &= \left(\frac{1 + (nb - 1)w(z)}{1 - w(z)} \right) \\
 & \times \left[1 - \alpha + \alpha(1 - \lambda) \left(\frac{1 + (nb - 1)w(z)}{1 - w(z)} \right) \right. \\
 & \quad \left. + \alpha\lambda \left(\frac{nbzw'(z)}{(1 - w(z))[1 + (nb - 1)w(z)]} + \frac{1 + (nb - 1)w(z)}{1 - w(z)} \right) \right] \\
 &= (1 - \alpha) \left(\frac{1 + (nb - 1)w(z)}{1 - w(z)} \right) + \alpha \left(\frac{[1 + (nb - 1)w(z)]^2 + \lambda nbzw'(z)}{(1 - w(z))^2} \right).
 \end{aligned} \tag{4.9}$$

Now, we claim that $|w(z)| < 1$ ($z \in \mathbb{U}$). If there exists a z_o in \mathbb{U} such that $|w(z_o)| = 1$, then (by Jack's Lemma) Lemma 2.2, we have

$$z_o w'(z_o) = kw(z_o), \tag{4.10}$$

where $k \geq 1$ is a real number.

By setting $w(z_o) = e^{i\theta}$ ($0 \leq \theta \leq 2\pi$), thus, we find that

$$\begin{aligned}
 & \operatorname{Re} \left\{ \frac{z_o f'_b(z_o)}{f_b(z_o)} \left[1 - \alpha + \alpha(1 - \lambda) \frac{z_o f'_b(z_o)}{f_b(z_o)} + \alpha\lambda \left(1 + \frac{z_o f''_b(z_o)}{f'_b(z_o)} \right) \right] \right\} \\
 &= \operatorname{Re} \left\{ (1 - \alpha) \left(\frac{1 + (nb - 1)w(z_o)}{1 - w(z_o)} \right) + \alpha \left(\frac{\lambda nb z_o w'(z_o) + [1 + (nb - 1)w(z_o)]^2}{(1 - w(z_o))^2} \right) \right\} \\
 &= \operatorname{Re} \left\{ (1 - \alpha) \left(\frac{1 + (nb - 1)e^{i\theta}}{1 - e^{i\theta}} \right) + \alpha \left(\frac{\lambda nb k e^{i\theta} + [1 + (nb - 1)e^{i\theta}]^2}{(1 - e^{i\theta})^2} \right) \right\} \\
 &= \operatorname{Re} \left\{ \frac{1 + (\alpha \lambda nb k + \alpha nb + nb - 2)e^{i\theta} + (nb - 1)(\alpha(nb - 1) - (1 - \alpha))e^{i2\theta}}{(1 - e^{i\theta})^2} \right\} \\
 &= \frac{[2\alpha nb(\lambda k + 2 - nb) + 4nb - 8] \cos \theta + [(nb - 1)(\alpha nb - 1) + 1] \cos 2\theta - \alpha nb(\lambda k + 3 - 2nb) - 3nb + 6}{2(3 - 4 \cos \theta) + 2 \cos 2\theta} \\
 &\leq \left(1 - \frac{nb}{2} \right) \left(1 - \frac{\alpha nb}{2} \right) - \frac{\alpha \lambda nb}{4}, \quad (z \in \mathbb{U}),
 \end{aligned} \tag{4.11}$$

since $k \geq 1$.

If we let

$$\begin{aligned} & \operatorname{Re} \left\{ \frac{z_0 f'_b(z_0)}{f_b(z_0)} \left[1 - \alpha + \alpha(1 - \lambda) \frac{z_0 f'_b(z_0)}{f_b(z_0)} + \alpha \lambda \left(1 + \frac{z_0 f''_b(z_0)}{f'_b(z_0)} \right) \right] \right\} \\ & \leq \left(1 - \frac{nb}{2} \right) \left(1 - \frac{\alpha nb}{2} \right) - \frac{\alpha \lambda nb}{4} \\ & = \tau(nb), \end{aligned} \quad (4.12)$$

then

$$\tau(nb) \leq 0, \quad \left(\frac{\alpha \lambda + 2\alpha + 2\sqrt{\Delta}}{2\alpha} \leq nb \leq \frac{\alpha \lambda + 2\alpha + 2\sqrt{\Delta}; \Delta}{2\alpha} := \alpha^2(\lambda + 2)^2 + 4\alpha(\lambda - 2) + 4 \right). \quad (4.13)$$

Thus, we have

$$\operatorname{Re} \left\{ \frac{z f'_b(z)}{f_b(z)} \left[1 - \alpha + \alpha(1 - \lambda) \frac{z f'_b(z)}{f_b(z)} + \alpha \lambda \left(1 + \frac{z f''_b(z)}{f'_b(z)} \right) \right] \right\} \leq 0 \quad (z \in \mathbb{U}), \quad (4.14)$$

$$\left(\frac{\alpha \lambda + 2\alpha + 2\sqrt{\Delta}}{2\alpha} \leq nb \leq \frac{\alpha \lambda + 2\alpha + 2\sqrt{\Delta}}{2\alpha}; \Delta := \alpha^2(\lambda + 2)^2 + 4\alpha(\lambda - 2) + 4 \right), \quad (4.15)$$

which is a contradiction to the hypotheses of (4.2).

Therefore, $|w(z)| < 1$ for all z in \mathbb{U} . Hence f_b is starlike in \mathbb{U} , then by proving the assertion (i) of Theorem 4.1, this completes the proof of our theorem. \square

Next, we arrive to the following remark which was given by Fukui et al. [11], and so we omit the detail here.

Remark 4.2. Let the n -fold symmetric function $f_b(z)$, defined by (1.4), be analytic in U , with

$$\frac{f_b(z)}{z} \neq 0 \quad (z \in \mathbb{U}). \quad (4.16)$$

If $f_b(z)$ satisfies the inequality (4.2) with $\alpha = 0$, that is, if

$$\operatorname{Re} \left(\frac{z f'_b(z)}{f_b(z)} \right) > 1 - \frac{nb}{2} \quad (z \in \mathbb{U}), \quad (4.17)$$

then $f_b(z)$ is starlike in \mathbb{U} for $0 \leq nb < 2$.

The following remark was obtained by Kamali and Srivastava [12].

Remark 4.3. Let the n -fold symmetric function $f_b(z)$, defined by (1.4), be analytic in U , with

$$\frac{f_b(z)}{z} \neq 0 \quad (z \in \mathbb{U}). \quad (4.18)$$

If $f_b(z)$ satisfies the inequality (4.2) with $\lambda = 1$, that is, if

$$\operatorname{Re} \left(\frac{\alpha z^2 f_b''(z)}{f_b(z)} + \frac{z f_b'(z)}{f_b(z)} \right) > -\frac{\alpha n b}{4} + \left(1 - \frac{n b}{2} \right) \left(1 - \frac{\alpha n b}{2} \right) \quad (z \in \mathbb{U}), \quad (4.19)$$

then $f_b(z)$ is starlike in \mathbb{U} for

$$\alpha > 0, \quad \frac{3\alpha + 2 - \sqrt{\Delta^*}}{2\alpha} \leq n b \leq \frac{3\alpha + 2\sqrt{\Delta^*}}{2\alpha}, \quad (\Delta^* := 9\alpha^2 - 4\alpha + 4). \quad (4.20)$$

5. Applications of Differential Inequalities

We apply the following result involving differential inequalities with a view to deriving several further sufficient conditions for starlikeness of the n -fold symmetric function f_b defined by (1.4).

Lemma 5.1 (Miller and Mocanu [13]). *Let $\Theta(u, v)$ be a complex-valued function such that*

$$\Theta : \mathbb{D} \longrightarrow \mathbb{C}, \quad (\mathbb{D} \subset \mathbb{C} \times \mathbb{C}), \quad (5.1)$$

\mathbb{C} being (as usual) the complex plane, and let

$$u = u_1 + iu_2, \quad v = v_1 + iv_2. \quad (5.2)$$

Suppose that the functions $\Theta(u, v)$ satisfies each of the following conditions.

- (i) $\Theta(u, v)$ is continuous in \mathbb{D} .
- (ii) $(1, 0) \in \mathbb{D}$ and $\operatorname{Re}(\Theta(1, 0)) > 0$.
- (iii) $\operatorname{Re}(\Theta(iu_2, v_1)) \leq 0$ for all $(iu_2, v_1) \in \mathbb{D}$ such that

$$v_1 \leq -\frac{1}{2}(1 + u_2^2). \quad (5.3)$$

Let

$$p(z) = 1 + p_1 z + p_2 z^2 + \cdots \quad (5.4)$$

be analytic (regular) in \mathbb{U} such that

$$(p(z), zp'(z)) \in \mathbb{D} \quad (z \in \mathbb{U}). \quad (5.5)$$

If

$$\operatorname{Re}(\Theta(p(z), zp'(z))) \in \mathbb{D} \quad (z \in \mathbb{U}), \quad (5.6)$$

then

$$\operatorname{Re}(p(z)) > 0 \quad (z \in \mathbb{U}). \quad (5.7)$$

Let us now consider the following implication.

Theorem 5.2. Let n -fold symmetric function f_b , defined by (1.4) and analytic in \mathbb{U} with $(f_b(z))/z, \neq 0$, ($z \in \mathbb{U}$), satisfy the following inequality:

$$\operatorname{Re} \left\{ \frac{zf'_b(z)}{f_b(z)} \left[1 - \alpha + \alpha(1 - \lambda) \frac{zf'_b(z)}{f_b(z)} + \alpha\lambda \left(1 + \frac{zf''_b(z)}{f'_b(z)} \right) \right] \right\} > \left(1 - \frac{nb}{2} \right) \left(1 - \frac{\alpha nb}{2} \right) - \frac{\alpha\lambda nb}{4}, \quad (5.8)$$

then

$$\operatorname{Re} \left\{ \left(\frac{zf'_b(z)}{f_b(z)} \right)^\mu \right\} > 0, \quad (5.9)$$

$$\left(z \in \mathbb{U}; \left(1 - \frac{nb}{2} \right) \left(1 - \frac{\alpha nb}{2} \right) - \frac{\alpha\lambda nb}{4} < 1; \alpha \geq 0, \lambda > 0; \mu \geq 1 \right).$$

Proof. If we put

$$p(z) = \left\{ \frac{zf'_b(z)}{f_b(z)} \right\}^\mu, \quad (5.10)$$

then (5.8) is equivalent to

$$\begin{aligned} & \operatorname{Re} \left\{ \frac{\alpha\lambda}{\mu} \{p(z)\}^{(1-\mu)/\mu} zp'(z) + \alpha \{p(z)\}^{2/\mu} + (1 - \alpha)p(z)^{1/\mu} \right. \\ & \quad \left. - \left(1 - \frac{nb}{2} \right) \left(1 - \frac{\alpha nb}{2} \right) + \frac{\lambda\alpha nb}{4} \right\} > 0 \\ & \implies \operatorname{Re}(p(z)) > 0 \quad (z \in \mathbb{U}). \end{aligned} \quad (5.11)$$

By setting $p(z) = u$ and $zp'(z) = v$ and letting

$$\Theta(z) = \frac{\alpha\lambda}{\mu} u^{(1-\mu)/\mu} v + \alpha u^{2/\mu} + (1 - \alpha)u^{1/\mu} - \left(1 - \frac{nb}{2} \right) \left(1 - \frac{\alpha nb}{2} \right) + \frac{\lambda\alpha nb}{4}, \quad (5.12)$$

for $\alpha \geq 0$ and $\mu \geq 1$, we have the following.

- (i) $\Theta(u, v)$ is continuous in $\mathbb{D} = (\mathbb{C} \setminus \{0\} \times \mathbb{C})$.
- (ii) $(1, 0) \in \mathbb{D}$ and

$$\operatorname{Re}(\Theta(1, 0)) = \frac{\alpha \lambda n b}{4} + \frac{\alpha n b}{2} + \frac{n b}{2} - \frac{\alpha n^2 b^2}{4} > 0, \quad (5.13)$$

since

$$\left(1 - \frac{n b}{2}\right) \left(1 - \frac{\alpha n b}{2}\right) - \frac{\lambda \alpha n b}{4} < 1. \quad (5.14)$$

Thus, the conditions (i) and (ii) of Lemma 5.1 are satisfied. Moreover, for $(iu_2, v_1) \in \mathbb{D}$ and $v_1 \leq (-1/2)(1 + u_2^2)$, we obtain

$$\begin{aligned} \operatorname{Re}(\Theta(iu_2, v_1)) &= \frac{\alpha \lambda}{\mu} |u_2|^{(1-\mu)/\mu} v_1 \cos\left(\frac{(1-\mu)\pi}{2\mu}\right) + \alpha |u_2|^{2/\mu} \cos\left(\frac{\pi}{\mu}\right) \\ &\quad + (1-\alpha) |u_2|^{1/\mu} \cos\left(\frac{\pi}{2\mu}\right) - \left(1 - \frac{n b}{2}\right) \left(1 - \frac{\alpha n b}{2}\right) + \frac{\lambda \alpha n b}{4} \\ &\leq -\frac{\alpha \lambda}{2\mu} (1 + u_2^2) |u_2|^{(1-\mu)/\mu} \sin\left(\frac{\pi}{2\mu}\right) + \alpha |u_2|^{2/\mu} \cos\left(\frac{\pi}{\mu}\right) \\ &\quad + (1-\alpha) |u_2|^{1/\mu} \cos\left(\frac{\pi}{2\mu}\right) - \left(1 - \frac{n b}{2}\right) \left(1 - \frac{\alpha n b}{2}\right) + \frac{\lambda \alpha n b}{4}, \end{aligned} \quad (5.15)$$

which, upon putting $|u_2| = \zeta$ ($\zeta > 0$), yields

$$\operatorname{Re}(\Theta(iu_2, v_1)) \leq \Phi(\zeta), \quad (5.16)$$

where

$$\begin{aligned} \Phi(\zeta) &:= -\frac{\alpha \lambda}{2\mu} (1 + \zeta^2) \zeta^{(1-\mu)/\mu} \sin\left(\frac{\pi}{2\mu}\right) + \alpha \zeta^{2/\mu} \cos\left(\frac{\pi}{\mu}\right) + (1-\alpha) \zeta^{1/\mu} \cos\left(\frac{\pi}{2\mu}\right) \\ &\quad - \left(1 - \frac{n b}{2}\right) \left(1 - \frac{\alpha n b}{2}\right) + \frac{\lambda \alpha n b}{4}. \end{aligned} \quad (5.17)$$

□

Remark 5.3. If, for some choices of the parameters α , λ , μ , and $n b$, we find that

$$\Phi(\zeta) \leq 0 \quad (\zeta > 0), \quad (5.18)$$

then we can conclude from (5.16) and Lemma 5.1 that the corresponding implication (5.8) holds true.

First of all, for the choice: $\mu = 1$ and $nb = 2$, we have the following.

Theorem 5.4. *If n -fold symmetric function f_b , defined by (1.4) and analytic in \mathbb{U} with*

$$\frac{f_b(z)}{z} \neq 0, \quad (z \in \mathbb{U}), \quad (5.19)$$

satisfies the following inequality:

$$\operatorname{Re} \left\{ \frac{zf'_b(z)}{f_b(z)} \left[1 - \alpha + \alpha(1 - \lambda) \frac{zf'_b(z)}{f_b(z)} + \alpha \lambda \left(1 + \frac{zf''_b(z)}{f'_b(z)} \right) \right] \right\} > -\frac{\alpha \lambda}{2}, \quad (5.20)$$

then $f_b \in S^$ for any real $\alpha \geq 0$ and $\lambda > 0$.*

Proof. For $\mu = 1$, $nb = 2$, we find from (5.17) that

$$\Phi(\zeta) := -\alpha \lambda \left(\frac{1}{2} + s^2 \right) - \alpha s^2 \leq 0, \quad (\zeta \in \mathbb{R}), \quad (5.21)$$

which implies Theorem 5.4 in view of the remark. \square

Remark 5.5. For $\lambda = 1$, we will obtain the results by Kamali and Srivastava [12].

Acknowledgment

The work presented here was supported by UKM-ST-06-FRGS0244-2010.

References

- [1] P. T. Mocanu, "Une propriété de convexité généralisée dans la théorie de la représentation conforme," *Mathematica*, vol. 11, no. 34, pp. 127–133, 1969.
- [2] S. S. Miller, P. Mocanu, and M. O. Reade, "All α -convex functions are univalent and starlike," *Proceedings of the American Mathematical Society*, vol. 37, pp. 553–554, 1973.
- [3] K. Sakaguchi and S. Fukui, "On alpha-starlike functions and related functions," *Bulletin of Nara University of Education*, vol. 28, no. 2, pp. 5–12, 1979.
- [4] C. Ramesha, S. Kumar, and K. S. Padmanabhan, "A sufficient condition for starlikeness," *Chinese Journal of Mathematics*, vol. 23, no. 2, pp. 167–171, 1995.
- [5] S. Siregar, M. Darus, and B. A. Frasin, "Subordination and superordination for certain analytic functions," *Bulletin of Mathematical Analysis and Applications*, vol. 2, no. 2, pp. 42–49, 2010.
- [6] D. Bansal and R. K. Raina, "Some sufficient conditions for starlikeness using subordination criteria," *Bulletin of Mathematical Analysis and Applications*, vol. 2, no. 4, pp. 1–6, 2010.
- [7] S. Siregar, M. Darus, and T. Bulboacă, "A class of superordination-preserving convex integral operator," *Mathematical Communications*, vol. 14, no. 2, pp. 379–390, 2009.
- [8] B. A. Frasin and M. Darus, "Subordination results on subclasses concerning Sakaguchi functions," *Journal of Inequalities and Applications*, vol. 2009, Article ID 574014, 7 pages, 2009.
- [9] S. S. Miller and P. T. Mocanu, *Differential Subordinations: Theory and Applications*, vol. 225 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 2000.

- [10] I. S. Jack, "Functions starlike and convex of order α ," *Journal of the London Mathematical Society*, vol. 3, pp. 469–474, 1971.
- [11] S. Fukui, S. Owa, and K. Sakaguchi, "Some properties of analytic functions of Koebe type," in *Current topics in Analytic Function Theory*, pp. 106–117, World Scientific Publishing Company, New Jersey, NJ, USA, 1992.
- [12] M. Kamali and H. M. Srivastava, "A sufficient condition for starlikeness of analytic functions of Koebe type," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 5, no. 3, pp. 1–8, 2004.
- [13] S. S. Miller and P. T. Mocanu, "Second-order differential inequalities in the complex plane," *Journal of Mathematical Analysis and Applications*, vol. 65, no. 2, pp. 289–305, 1978.

