REISSNER-MINDLIN PLATE THEORY FOR ELASTODYNAMICS

SHEN RONG WU

Received 14 January 2004 and in revised form 11 May 2004

Existence and uniqueness of solution are proved for elastodynamics of Reissner-Mindlin plate model. Higher regularity is proved under the assumptions of smoother data and certain compatibility conditions. A mass scaling is introduced. When the thickness approaches zero, the solution of the clamped Reissner-Mindlin plate is shown to approach the solution of a Kirchhoff-Love plate.

1. Introduction

The Reissner-Mindlin (R-M) theory [13, 14, 15] has been popularly applied to thin-walled structures with moderate thickness. Transient response plays an important role in many aspects of structural analysis. The governing equation of the elastodynamics problem of R-M plate is of an evolutionary type with second-order time derivatives. In this paper, we apply a priori estimate to investigate the elastodynamics problem of R-M plate. This method has been successfully used in developing the theory of various partial differential equations, for example, [7, 10, 11]. Following the line of [7, 10, 11], we prove the existence and uniqueness of the H^1 solution. We then apply the approaches of [8] to prove the H^2 regularity and higher regularity when the data is smoother and certain compatibility conditions are satisfied.

For static problem under the assumption of load scaling, it is proved in [3] that the solution of the clamped R-M plate approaches the solution of the Kirchhoff-Love (K-L) plate when the thickness approaches zero. This fact has been employed to investigate the finite element method of R-M plate, such as locking-free and uniform convergence, cf. [2, 3, 5, 12, 16, 18]. For dynamics problem, with the introduction of mass scaling [4], we prove that when thickness approaches zero, the H^2 strong solution of the clamped R-M plate approaches the H^2 weak solution of K-L plate (whose classical solution requires H^4 smoothness).

In what follows, we describe the system of equations in Section 2 and prove the existence, uniqueness, and regularity in Section 3. Then we discuss the relation between R-M plate and K-L plate in Section 4. This is followed by a summary in Section 5.

2. Governing equations of Reissner-Mindlin plate for elastodynamics

For elastodynamic bending shear problem modeled by R-M plate theory [13, 14, 15], the displacement components of a generic point at a distance z to the midsurface are expressed by the deflection w at the midsurface and the rotations (β_1, β_2) of the normal to the midsurface,

$$U_1 = -z\beta_1, \quad U_2 = -z\beta_2, \quad U_3 = w, \quad (2.1)$$

$|z| \leq \zeta/2$. ζ is the thickness of the plate. For dynamics problems, the velocity and acceleration, traditionally denoted by $\dot{U}_i$ and $\ddot{U}_i$, respectively, have the same format of (2.1) after differentiation with respect to time. The motion equation of R-M plate can be derived from the general three-dimensional elastodynamics by integration through thickness, or from the energy method using Hamilton's principle, for example, [9]

$$\begin{aligned} I\ddot{\beta}_1 + EA_1(\boldsymbol{\beta}) - \lambda\zeta^{-2}(w_{,1} - \beta_1) &= m_1, \\ I\ddot{\beta}_2 + EA_2(\boldsymbol{\beta}) - \lambda\zeta^{-2}(w_{,2} - \beta_2) &= m_2, \\ \rho\zeta^{-2}\ddot{w} - \lambda\zeta^{-2}\nabla \cdot (\nabla w - \boldsymbol{\beta}) &= g = f_3\zeta^{-2}, \end{aligned} \quad (2.2)$$

where we define

$$\begin{aligned} A_1(\boldsymbol{\beta}) &= \frac{-((1+\nu)(\beta_{\alpha,\alpha})_{,1} + (1-\nu)\nabla^2\beta_1)}{24(1-\nu^2)}, \\ A_2(\boldsymbol{\beta}) &= \frac{-((1+\nu)(\beta_{\alpha,\alpha})_{,2} + (1-\nu)\nabla^2\beta_2)}{24(1-\nu^2)}. \end{aligned} \quad (2.3)$$

Here, E is the Young's modulus, ρ is the density, and ν is the Poisson ratio. We denote $I = \rho/12$ and $\lambda = G\kappa$, with the shear modulus G and a shear correction factor κ , which is introduced to balance the zero shear stress at the top and bottom surfaces. As analyzed for static problem [2, 3, 5], the lateral loading force f_3 (per unit volume) is scaled to $\zeta^2 g$. The convention of summation on repeated indices is also applied, with the Greek index running over the range from 1 to 2. The bold-faced variables are used to denote a two-dimensional vector, for example, $\boldsymbol{\beta} = (\beta_1, \beta_2)$, and β_α is used to indicate all of the two components involved when the indication is clear. Here, $w_{,1}$ in (2.2) indicates the partial derivative $\partial w / \partial x_1$. The same applies for all the similar cases.

For simplicity, we consider the equations defined on a smooth bounded domain Ω in $\mathbb{R}^2$, with homogeneous Dirichlet boundary conditions and general initial conditions

$$\begin{aligned} \beta_\alpha(t, \mathbf{x}) &= 0; \quad w(t, \mathbf{x}) = 0 \quad \text{on } \partial\Omega, \\ \beta_\alpha(0, \mathbf{x}) &= \beta_\alpha^0(\mathbf{x}); \quad w(0, \mathbf{x}) = W^0(\mathbf{x}), \\ \dot{\beta}_\alpha(0, \mathbf{x}) &= \dot{\beta}_\alpha^1(\mathbf{x}); \quad \dot{w}(0, \mathbf{x}) = W^1(\mathbf{x}). \end{aligned} \quad (2.4)$$

We adopt the usual notations of Sobolev spaces. The Galerkin method yields the following variational equation. For any $t \in [0, T]$, find $\beta_\alpha, w \in V = H_0^1(\Omega)$ such that

$$I\langle \ddot{\beta}_\alpha, \eta_\alpha \rangle + \rho \zeta^{-2} \langle \ddot{w}, v \rangle + Ea(\beta, \eta) + \lambda \zeta^{-2} (w_{,\alpha} \beta_\alpha, v_{,\alpha} \eta_\alpha) = (m_\alpha, \eta_\alpha) + (g, v), \quad \forall \eta_\alpha, v \in V. \quad (2.5)$$

Here, $(\cdot, \cdot)$ denotes the usual L^2 inner product, and $\langle \cdot, \cdot \rangle$ denotes the duality on $V' \otimes V$. $a(\cdot, \cdot)$ is a bilinear form on $V \otimes V$ defined as

$$a(\beta, \eta) = \frac{1}{24(1-\nu^2)} ((1+\nu)(\beta_{\alpha,\alpha}, \eta_{\alpha,\alpha}) + (1-\nu)(\nabla \beta_\alpha, \nabla \eta_\alpha)). \quad (2.6)$$

It is associated with the operators A_1 and A_2 such that

$$a(\beta, \eta) = \langle A_1(\beta), \eta_1 \rangle + \langle A_2(\beta), \eta_2 \rangle = \langle \mathbf{A}(\beta), \boldsymbol{\eta} \rangle, \quad \forall \beta_\alpha, \eta_\alpha \in V. \quad (2.7)$$

In fact, $a(\cdot, \cdot)$ is symmetric, the same as the two-dimensional elasticity operator with a scalar factor. With Dirichlet boundary conditions, $a(\cdot, \cdot)$ is equivalent to the H^1 -norm on V [6] and there exist constants $\alpha_1, \alpha_2 > 0$ such that

$$\begin{aligned} \alpha_1 \|\boldsymbol{\eta}\|_1^2 &\leq a(\boldsymbol{\eta}, \boldsymbol{\eta}), \quad \forall \eta_\alpha \in V, \\ a(\beta, \eta) &\leq \alpha_2 \|\beta\|_1 \|\eta\|_1, \quad \forall \beta_\alpha, \eta_\alpha \in V. \end{aligned} \quad (2.8)$$

We denote $\|\boldsymbol{\eta}\|_x^2 = \|\eta_1\|_x^2 + \|\eta_2\|_x^2$ for η_α of a functional space X . Note that for time-dependent problems, the norm $\|\nu\|_X$ of a function $\nu: [0, T] \rightarrow X$ is a function of time. We use the following notations for the functional spaces and the measure in time:

$$\begin{aligned} L^2(X) &= L^2(0, T; X) \\ &= \left\{ \nu: [0, T] \rightarrow X \mid \nu(t, \mathbf{x}) \in X, \|\nu\|_{L^2(0, T; X)} = \left(\int_0^T (\|\nu\|_X)^2 dt \right)^{1/2} < \infty \right\}, \\ L^\infty(X) &= L^\infty(0, T; X) \\ &= \left\{ \nu: [0, T] \rightarrow X \mid \nu(t, \mathbf{x}) \in X, \|\nu\|_{L^\infty(0, T; X)} = \operatorname{esssup}_{0 \leq t \leq T} (\|\nu\|_X) < \infty \right\}. \end{aligned} \quad (2.9)$$

3. Existence, uniqueness, and regularity

For linear hyperbolic equations of the second order in time with one function, the existence and uniqueness are proven (see, e.g., [7, 10, 11]) using the method of a priori estimate. The method is also employed in [17] for Navier-Stokes problem whose steady-state case has close relation to the static problems of R-M plate, (cf. [5]). We extend the scheme to the dynamic problems of R-M plate. For the time being, we keep the material parameters explicitly expressed for later use in Section 4.

THEOREM 3.1. *If $m_\alpha, g \in L^2(L^2)$; $B_\alpha^0, W^0 \in H_0^1$; and $B_\alpha^1, W^1 \in L^2$, then there exists a solution (β_α, w) of (2.5) (a weak solution of (2.2)) with initial conditions (2.4), $\beta_\alpha, w \in L^\infty(H_0^1)$, $\dot{\beta}_\alpha, \dot{w} \in L^\infty(L^2)$, and $\ddot{\beta}_\alpha, \ddot{w} \in L^\infty(H^{-1})$. Moreover, there exists a constant $C > 0$, independent*

of the material parameters, such that

$$\begin{aligned}
 & I \|\dot{\boldsymbol{\beta}}\|_0^2 + \rho \zeta^{-2} \|\dot{w}\|_0^2 + E \|\boldsymbol{\beta}\|_1^2 + \lambda \zeta^{-2} \|\nabla w - \boldsymbol{\beta}\|_0^2 \\
 & \leq C \left(I \|\mathbf{B}^1\|_0^2 + \rho \zeta^{-2} \|W^1\|_0^2 + E \|\mathbf{B}^0\|_1^2 \right. \\
 & \quad \left. + \lambda \zeta^{-2} \|\nabla W^0 - \mathbf{B}^0\|_0^2 + I^{-1} \|\mathbf{m}\|_{L^2(L^2)}^2 + \rho^{-1} \zeta^2 \|g\|_{L^2(L^2)}^2 \right).
 \end{aligned} \tag{3.1}$$

Proof. We apply the scheme for hyperbolic equations developed in [7, 10, 11]. The space V is separable. We construct an approximation of order n , with a countable basis $\{\psi_i(\mathbf{x}), i = 1, 2, \dots\}$ of V :

$$\begin{aligned}
 \beta_{an}(t, \mathbf{X}) &= \sum_{j=1}^n \beta_{an}^j(t) \psi_j(\mathbf{X}), & w_n(t, \mathbf{X}) &= \sum_{j=1}^n w_n^j(t) \psi_j(\mathbf{X}), \\
 I(\ddot{\beta}_{an}, \eta_\alpha) + \rho \zeta^{-2}(\dot{w}_n, \nu) + Ea(\boldsymbol{\beta}_n, \boldsymbol{\eta}) + \lambda \zeta^{-2}(w_{n,\alpha} - \beta_{an}, \nu_\alpha - \eta_\alpha) \\
 &= (g, \nu) + (m_\alpha, \eta_\alpha), \quad \forall \eta_\alpha, \nu \in V_n = \text{span}\{\psi_1, \dots, \psi_n\}.
 \end{aligned} \tag{3.2}$$

The approximation problem (3.2) leads to a linear system of second-order ordinary differential equations. With the approximations of (2.4) for the initial conditions

$$\begin{aligned}
 \sum_{j=1}^n B_{an}^{0j} \psi_j(\mathbf{X}) &= B_{an}^0 \longrightarrow B_\alpha^0, & \sum_{j=1}^n W_n^{0j} \psi_j(\mathbf{X}) &= W_n^0 \longrightarrow W^0, \\
 \sum_{j=1}^n B_{an}^{1j} \psi_j(\mathbf{X}) &= B_{an}^1 \longrightarrow B_\alpha^1, & \sum_{j=1}^n W_n^{1j} \psi_j(\mathbf{X}) &= W_n^1 \longrightarrow W^1,
 \end{aligned} \tag{3.3}$$

we have unique solution

$$\{\beta_{1n}^j(t), \beta_{2n}^j(t), w_n^j(t), j = 1, \dots, n\} \in H^2([0, T]). \tag{3.4}$$

Now using $\eta_\alpha = \dot{\beta}_{an}(t, \mathbf{X})$ and $\nu = \dot{w}_n(t, \mathbf{X})$ in (3.2), then integrating from $t = 0$ to T , we have

$$\begin{aligned}
 & I \|\dot{\boldsymbol{\beta}}_n\|_0^2 + \rho \zeta^{-2} \|\dot{w}_n\|_0^2 + Ea(\boldsymbol{\beta}_n, \boldsymbol{\beta}_n) + \lambda \zeta^{-2} \|\nabla w_n - \boldsymbol{\beta}_n\|_0^2 \\
 & = I \|\dot{\boldsymbol{\beta}}_n(0)\|_0^2 + \rho \zeta^{-2} \|\dot{w}_n(0)\|_0^2 + Ea(\boldsymbol{\beta}_n(0), \boldsymbol{\beta}_n(0)) \\
 & \quad + \lambda \zeta^{-2} \|\nabla w_n(0) - \boldsymbol{\beta}_n(0)\|_0^2 + 2 \int_0^T ((m_\alpha, \dot{\beta}_{an}) + (g, \dot{w}_n)) dt \\
 & \leq I \|\mathbf{B}_n^1\|_0^2 + \rho \zeta^{-2} \|W_n^1\|_0^2 + E \alpha_2 \|\mathbf{B}_n^0\|_1^2 + \lambda \zeta^{-2} \|\nabla W_n^0 - \mathbf{B}_n^0\|_0^2 \\
 & \quad + \int_0^T (I^{-1} \|\mathbf{m}\|_0^2 + \rho^{-1} \zeta^2 \|g\|_0^2) dt + \int_0^T (I \|\dot{\boldsymbol{\beta}}_n\|_0^2 + \rho \zeta^{-2} \|\dot{w}_n\|_0^2) dt.
 \end{aligned} \tag{3.5}$$

Applying Gronwall inequality, we obtain

$$\begin{aligned}
 & I\|\dot{\beta}_n\|_0^2 + \rho\zeta^{-2}\|\dot{w}_n\|_0^2 + \alpha_1 E\|\beta_n\|_1^2 + \lambda\zeta^{-2}\|\nabla w_n - \beta_n\|_0^2 \\
 & \leq C\left(I\|\mathbf{B}_n^1\|_0^2 + \rho\zeta^{-2}\|W_n^1\|_0^2 + \alpha_2 E\|\mathbf{B}_n^0\|_1^2\right. \\
 & \quad \left.+ \lambda\zeta^{-2}\|\nabla W_n^0 - \mathbf{B}_n^0\|_0^2 + I^{-1}\|\mathbf{m}\|_{L^2(L^2)}^2 + \rho^{-1}\zeta^2\|g\|_{L^2(L^2)}^2\right).
 \end{aligned} \tag{3.6}$$

The right-hand side has limit as $n \rightarrow \infty$ due to (3.3). Therefore, the left-hand side is bounded. Note that $\|w_n\|_1 \leq C\|\nabla w_n\|_0 \leq C(\|\nabla w_n - \beta_n\|_0 + \|\beta_n\|_0)$. By compactness, we can find convergent subsequences, still denoted by subscript n , such that

$$\begin{aligned}
 \beta_{\alpha n} &\rightharpoonup \beta_\alpha, & w_n &\rightharpoonup w \quad \text{weakly star in } L^\infty(H_0^1), \\
 \dot{\beta}_{\alpha n} &\rightharpoonup \varphi_\alpha, & \dot{w}_n &\rightharpoonup \chi \quad \text{weakly star in } L^\infty(L^2).
 \end{aligned} \tag{3.7}$$

It is a straightforward task to verify that $\dot{\beta}_\alpha = \varphi_\alpha$, $\dot{w} = \chi$, $\ddot{\beta}_{\alpha n} \rightharpoonup \ddot{\beta}_\alpha$, $\ddot{w}_n \rightharpoonup \ddot{w}$ weakly star in $L^\infty(H^{-1})$, and $\{\beta_\alpha, w\}$ satisfy the initial conditions (2.4) and the variational equation (2.5), thus form a weak solution of (2.2). $\square$

THEOREM 3.2. *Under the conditions of Theorem 3.1, the solution $\{\beta_\alpha, w\}$ is unique, that is, if $g = 0$, $m_\alpha = 0$, $B_\alpha^0 = w^0 = B_\alpha^1 = w^1 = 0$, then $\beta_\alpha = w = 0$.*

Proof. Following the line of [10, 11], we can prove the uniqueness, but omit the details. $\square$

THEOREM 3.3. *Under the conditions of Theorem 3.1, if $\dot{m}_\alpha, \dot{g} \in L^2(L^2)$, $B_\alpha^0, W^0 \in H^2$, and $B_\alpha^1, W^1 \in H_0^1$, then the solution (β_α, w) of (2.2) with the initial conditions (2.4) satisfies $\ddot{\beta}_\alpha, \ddot{w} \in L^\infty(L^2)$, $\dot{\beta}_\alpha, \dot{w} \in L^\infty(H_0^1)$, $\beta_\alpha, w \in L^\infty(H^2)$, and*

$$\begin{aligned}
 & I\|\ddot{\beta}\|_0^2 + \rho\zeta^{-2}\|\ddot{w}\|_0^2 + E\|\dot{\beta}\|_1^2 + \zeta^{-2}\|\nabla \dot{w} - \dot{\beta}\|_0^2 \\
 & \leq C\left(I^{-1}\left(E^2\|\mathbf{B}^0\|_2^2 + \lambda^2\zeta^{-4}\|\nabla W^0 - \mathbf{B}^0\|_0^2 + \|\mathbf{m}(0)\|_0^2\right)\right. \\
 & \quad \left.+ \rho^{-1}\zeta^2\left(\lambda^2\zeta^{-4}\|\nabla W^0 - \mathbf{B}^0\|_1^2 + \|g(0)\|_0^2\right)\right. \\
 & \quad \left.+ E\|\mathbf{B}^1\|_1^2 + \lambda\zeta^{-2}\|\nabla W^1 - \mathbf{B}^1\|_0^2 + I^{-1}\|\dot{\mathbf{m}}\|_{L^2(L^2)}^2 + \rho^{-1}\zeta^2\|\dot{g}\|_{L^2(L^2)}^2\right),
 \end{aligned} \tag{3.8}$$

$$E\|\beta\|_2 \leq C(I\|\ddot{\beta}\|_0 + \lambda\zeta^{-2}\|\nabla \dot{w} - \dot{\beta}\|_0 + \|\mathbf{m}\|_0), \tag{3.9}$$

$$\lambda\zeta^{-2}\|w\|_2 \leq C(\rho\zeta^{-2}\|\ddot{w}\|_0 + \lambda\zeta^{-2}\|\beta\|_1 + \|g\|_0), \tag{3.10}$$

where the bounds of $\|\nabla w - \beta\|_0$ and $\|\beta\|_1$ are established in (3.1).

Proof. We apply the method for hyperbolic equations demonstrated in [8]. From Theorem 3.1, we have $\dot{\beta}_{\alpha n}^j, \dot{w}_n^j \in H^1([0, T])$. Differentiating (3.2) with respect to t , we obtain $\ddot{\beta}_{\alpha n}^j, \ddot{w}_n^j \in L^2([0, T])$. The a priori estimate like (3.6) holds:

$$\begin{aligned}
 & I\|\ddot{\beta}_n\|_0^2 + \rho\zeta^{-2}\|\ddot{w}_n\|_0^2 + \alpha_1 E\|\dot{\beta}_n\|_1^2 + \lambda\zeta^{-2}\|\nabla \dot{w}_n - \dot{\beta}_n\|_0^2 \\
 & \leq C\left(I\|\ddot{\beta}_n(0)\|_0^2 + \rho\zeta^{-2}\|\ddot{w}_n(0)\|_0^2 + \alpha_2 E\|\dot{\beta}_n(0)\|_1^2\right. \\
 & \quad \left.+ \lambda\zeta^{-2}\|\nabla \dot{w}_n(0) - \dot{\beta}_n(0)\|_0^2 + I^{-1}\|\dot{\mathbf{m}}\|_{L^2(L^2)}^2 + \rho^{-1}\zeta^2\|\dot{g}\|_{L^2(L^2)}^2\right).
 \end{aligned} \tag{3.11}$$

From (2.2), we obtain $\ddot{\beta}_\alpha(0), \ddot{w}(0) \in L^2$,

$$\begin{aligned} I \|\ddot{\beta}_\alpha(0)\|_0 &\leq E \|\mathbf{B}^0\|_2 + \lambda \zeta^{-2} \|\nabla W^0 - \mathbf{B}^0\|_0 + \|m_\alpha(0)\|_0, \\ \rho \zeta^{-2} \|\ddot{w}(0)\|_0 &\leq \lambda \zeta^{-2} \|\nabla W^0 - \mathbf{B}^0\|_1 + \|g(0)\|_0. \end{aligned} \quad (3.12)$$

The argument of boundedness and compactness leads to the conclusion that $\ddot{\beta}_{\alpha n} \rightarrow \ddot{\beta}_\alpha$, $\ddot{w}_n \rightarrow \ddot{w}$ weakly star in $L^\infty(L^2)$. Hence, (3.11) implies (3.8).

On the other hand, we rewrite (2.2):

$$\begin{aligned} EA_1(\boldsymbol{\beta}) &= m_1 - I\ddot{\beta}_1 + \lambda \zeta^{-2} (w_{,1} - \beta_1), \\ EA_2(\boldsymbol{\beta}) &= m_2 - I\ddot{\beta}_2 + \lambda \zeta^{-2} (w_{,2} - \beta_2), \\ -\lambda \zeta^{-2} \nabla^2 w &= g - \rho \zeta^{-2} \ddot{w} - \lambda \zeta^{-2} \beta_{\alpha,\alpha}. \end{aligned} \quad (3.13)$$

For any fixed time t , the right-hand sides of these equations are in L^2 . We have the elasticity operator and the Laplace operator in the left-hand side. According to the theory of elliptic equations, with a smooth domain Ω , we have $\beta_{\alpha,w} \in H^2$, and the bounds (3.9) and (3.10). $\square$

We are ready to extend the method for higher regularity of hyperbolic equation [8] to the transient dynamics of R-M plate. For simplicity, the dependence on the material parameters is not explicitly expressed and will have more discussion in Section 4.

THEOREM 3.4. *Assume for any integer $P \geq 0$,*

$$\begin{aligned} B_\alpha^0, W^0 &\in H^{P+1} \cap H_0^1, \quad B_\alpha^1, W^1 \in H^P \cap H_0^1, \\ \frac{\partial^k m_\alpha}{\partial t^k}, \quad \frac{\partial^k g}{\partial t^k} &\in L^2(H^{P-k}), \quad k = 0, 1, \dots, P, \end{aligned} \quad (3.14)$$

and that the following compatibility conditions hold for $P \geq 2$:

$$\begin{aligned} B_\alpha^{k+2} &= I^{-1} \left(\frac{\partial^k m_\alpha(0)}{\partial t^k} - EA_\alpha(\mathbf{B}^k) + \lambda \zeta^{-2} (\nabla W^k - \mathbf{B}^k) \right) \in H_0^1, \\ W^{k+2} &= (\rho \zeta^{-2})^{-1} \left(\frac{\partial^k g(0)}{\partial t^k} + \lambda \zeta^{-2} \nabla \cdot (\nabla W^k - \mathbf{B}^k) \right) \in H_0^1, \end{aligned} \quad k = 0, 1, \dots, P-2. \quad (3.15)$$

Then the solution of (2.2) with (2.4) satisfy, for $k = 0, 1, \dots, P+1$,

$$\frac{\partial^k \beta_\alpha}{\partial t^k}, \quad \frac{\partial^k w}{\partial t^k} \in L^\infty(H^{P+1-k}), \quad (3.16)$$

$$\begin{aligned} &\left\| \frac{\partial^k \beta_\alpha}{\partial t^k} \right\|_{P+1-k} + \left\| \frac{\partial^k w}{\partial t^k} \right\|_{P+1-k} \\ &\leq C \left(\sum_{j=0}^P \left(\left\| \frac{\partial^j m_\alpha}{\partial t^j} \right\|_{L^2(H^{P-j})} + \left\| \frac{\partial^j g}{\partial t^j} \right\|_{L^2(H^{P-j})} \right) \right. \\ &\quad \left. + \|\mathbf{B}^0\|_{P+1} + \|W^0\|_{P+1} + \|\mathbf{B}^1\|_P + \|W^1\|_P \right). \end{aligned} \quad (3.17)$$

Proof. The cases of $P = 0$ and $P = 1$ are proved in Theorems 3.1 and 3.3, respectively. Using the method of induction, we assume that the theorem is true for $P \leq Q$ and assume that the conditions (3.14) and (3.15) are valid for $P = Q + 1$. Denote

$$\begin{aligned}\tilde{B}_\alpha &= \dot{\beta}_\alpha, & \tilde{w} &= \dot{w}, \\ \tilde{m}_\alpha &= \dot{m}_\alpha, & \tilde{g} &= \dot{g}, \\ \tilde{B}_\alpha^k &= B_\alpha^{k+1}, & \tilde{W}^k &= W^{k+1}, \quad k = 0, 1, \dots, Q.\end{aligned}\tag{3.18}$$

Then $\tilde{B}_\alpha^k, \tilde{W}^k$, $k = 0, 1, \dots, Q$, satisfy (3.15) for $P = Q$. $\tilde{B}_\alpha^0 = B_\alpha^1$ and $\tilde{W}^0 = W^1 \in H^{Q+1} \cap H_0^1$. For $k = 0, 1, \dots, Q$, $\partial^k \tilde{m}_\alpha / \partial t^k = \partial^{k+1} m_\alpha / \partial t^{k+1}$ and $\partial^k \tilde{g} / \partial t^k = \partial^{k+1} g / \partial t^{k+1} \in L^2(H^{Q-k})$. From (3.14) and (3.15) with $P = Q + 1$,

$$\begin{aligned}\|\tilde{\mathbf{B}}^1\|_Q &= \|\mathbf{B}^2\|_Q \leq C(\|\mathbf{m}(0)\|_Q + \|\mathbf{B}^0\|_{Q+2} + \|W^0\|_{Q+1} + \|\mathbf{B}^0\|_Q) \leq \infty, \\ \|\tilde{W}^1\|_Q &= \|W^2\|_Q \leq C(\|g(0)\|_Q + \|W^0\|_{Q+2} + \|\mathbf{B}^0\|_{Q+1}) \leq \infty.\end{aligned}\tag{3.19}$$

Hence $\tilde{B}_\alpha^k, \tilde{W}^k$, $k = 0$ and 1 , satisfy (3.14) for $P = Q$. We apply the assumption of induction and obtain from (3.16) and (3.17) with $P = Q$, for $k = 0, 1, \dots, Q + 1$,

$$\begin{aligned}\frac{\partial^k \tilde{B}_\alpha}{\partial t^k}, \quad \frac{\partial^k \tilde{w}}{\partial t^k} &\in L^\infty(H^{Q+1-k}), \\ \left\| \frac{\partial^k \tilde{\beta}_\alpha}{\partial t^k} \right\|_{Q+1-k} + \left\| \frac{\partial^k \tilde{w}}{\partial t^k} \right\|_{Q+1-k} \\ &\leq C \left(\sum_{j=0}^Q \left(\left\| \frac{\partial^j \tilde{m}_\alpha}{\partial t^j} \right\|_{L^2(H^{Q-j})} + \left\| \frac{\partial^j \tilde{g}}{\partial t^j} \right\|_{L^2(H^{Q-j})} \right) \right. \\ &\quad \left. + \|\tilde{\mathbf{B}}^0\|_{Q+1} + \|\tilde{W}^0\|_{Q+1} + \|\tilde{\mathbf{B}}^1\|_Q + \|\tilde{W}^1\|_Q \right).\end{aligned}\tag{3.20}$$

It implies that, for $k = 1, \dots, Q + 2$,

$$\begin{aligned}\frac{\partial^k \beta_\alpha}{\partial t^k}, \quad \frac{\partial^k w}{\partial t^k} &\in L^\infty(H^{Q+2-k}), \\ \left\| \frac{\partial^k \beta_\alpha}{\partial t^k} \right\|_{Q+2-k} + \left\| \frac{\partial^k w}{\partial t^k} \right\|_{Q+2-k} \\ &\leq C \left(\sum_{j=0}^{Q+1} \left(\left\| \frac{\partial^j m_\alpha}{\partial t^j} \right\|_{L^2(H^{Q+1-j})} + \left\| \frac{\partial^j g}{\partial t^j} \right\|_{L^2(H^{Q+1-j})} \right) \right. \\ &\quad \left. + \|\mathbf{B}^1\|_{Q+1} + \|W^1\|_{Q+1} + \|\mathbf{B}^2\|_Q + \|W^2\|_Q \right).\end{aligned}\tag{3.21}$$

We can use (3.19) to estimate $\mathbf{B}^2$ and W^2 in (3.23) with

$$\begin{aligned}\|\mathbf{m}(0)\|_Q &\leq C\|\mathbf{m}\|_{C^0(H^Q)} \leq C(\|\mathbf{m}\|_{L^2(H^Q)} + \|\dot{\mathbf{m}}\|_{L^2(H^Q)}), \\ \|g(0)\|_Q &\leq C\|g\|_{C^0(H^Q)} \leq C(\|g\|_{L^2(H^Q)} + \|\dot{g}\|_{L^2(H^Q)}).\end{aligned}\tag{3.24}$$

Therefore, (3.17) is true for $P = Q + 1$ and $k = 1, \dots, Q + 2$. Now the right-hand sides of (3.13) are bounded in H^Q . We have

$$\begin{aligned} \|\boldsymbol{\beta}\|_{Q+2}^2 &\leq C \|\mathbf{I}\ddot{\boldsymbol{\beta}} - \lambda\zeta^{-2}(\nabla w - \boldsymbol{\beta}) - \mathbf{m}\|_Q^2 \leq C(\|\ddot{\boldsymbol{\beta}}\|_Q^2 + \|w\|_{Q+1}^2 + \|\boldsymbol{\beta}\|_Q^2 + \|\mathbf{m}\|_Q^2) \leq \infty, \\ \|w\|_{Q+2}^2 &\leq C \|\rho\zeta^{-2}\ddot{w} + \lambda\zeta^{-2}\beta_{\alpha,\alpha} - g\|_Q^2 \leq C(\|\ddot{w}\|_Q^2 + \|\boldsymbol{\beta}\|_{Q+1}^2 + \|g\|_Q^2) \leq \infty. \end{aligned} \quad (3.25)$$

Therefore,

$$\begin{aligned} &\|\boldsymbol{\beta}\|_{Q+2} + \|w\|_{Q+2} \\ &\leq C \left(\sum_{j=0}^{Q+1} \left(\left\| \frac{\partial^j m_\alpha}{\partial t^j} \right\|_{L^2(H^{Q+1-j})} + \left\| \frac{\partial^j g}{\partial t^j} \right\|_{L^2(H^{Q+1-j})} \right) \right. \\ &\quad \left. + \|\mathbf{B}^0\|_{Q+2} + \|W^0\|_{Q+2} + \|\mathbf{B}^1\|_{Q+1} + \|W^1\|_{Q+1} \right). \end{aligned} \quad (3.26)$$

Thus, (3.17) also holds for $P = Q + 1$ and $k = 0$. The case of $P = Q + 1$ of the induction is true. $\square$

4. Relation to Kirchhoff-Love plate

For static problem, it is understood that when the thickness $\zeta \rightarrow 0$, the solution of the clamped R-M plate approaches the solution of a K-L plate (see, e.g., [3] for a proof). The convergence is for the systems with load scaling, in the sense that $\beta_\alpha \rightarrow \tilde{\beta}_\alpha$, $w \rightarrow \tilde{w}$, and

$$\tilde{\boldsymbol{\beta}} = \nabla \tilde{w}, \quad (4.1)$$

$$D_0 \nabla^4 \tilde{w} = g, \quad (4.2)$$

where $D_0 = E/12(1 - \nu^2) = D\zeta^{-3}$. D is the usual bending stiffness. Due to the load scaling, the K-L equation (4.2) is independent of thickness. Physically, when the thickness approaches zero, the bending stiffness approaches zero faster with a factor of ζ^3 . The unscaled loading, which contributes to the external work, is proportional to the thickness and will not give a meaningful solution. This fact is used for investigating the thickness-independent convergence of finite element method, for example, [2, 3, 5, 12, 16, 18] (see [12, 16, 18] for numerical examples).

For dynamic problem, due to the appearance of the inertia term, which contributes to the kinetic energy, the equation of K-L plate is no longer thickness independent. To keep K-L plate as a reference model, a possible approach is then to scale the mass density [4] along with the load. Assume

$$\begin{aligned} \rho &= \zeta^2 \rho_0, \\ I &= \zeta^2 I_0, \quad I_0 = \frac{\rho_0}{12}. \end{aligned} \quad (4.3)$$

We consider the scaled R-M equation (2.2) with $m_\alpha = 0$, which does not appear in K-L plate:

$$\begin{aligned} I_0 \zeta^2 \ddot{\beta}_1 + EA_1(\boldsymbol{\beta}) - \lambda \zeta^{-2} (w_{,1} - \beta_1) &= 0, \\ I_0 \zeta^2 \ddot{\beta}_2 + EA_2(\boldsymbol{\beta}) - \lambda \zeta^{-2} (w_{,2} - \beta_2) &= 0, \\ \rho_0 \ddot{w} - \lambda \zeta^{-2} \nabla \cdot (\nabla w - \boldsymbol{\beta}) &= g, \end{aligned} \quad (4.4)$$

or the variational equation (2.5):

$$I_0 \zeta^2 \langle \ddot{\beta}_\alpha, \eta_\alpha \rangle + \rho_0 \langle \ddot{w}, \nu \rangle + Ea(\boldsymbol{\beta}, \boldsymbol{\eta}) + \lambda \zeta^{-2} (w_{,\alpha} - \beta_\alpha, \nu_{,\alpha} - \eta_\alpha) = (g, \nu), \quad \forall \eta_\alpha, \nu \in V. \quad (4.5)$$

As a parallel study to the static problem, we consider a special case of elastodynamics with zero initial conditions:

$$B_\alpha^0 = B_\alpha^1 = W^0 = W^1 = 0. \quad (4.6)$$

THEOREM 4.1. Assume $g \in H^1(L^2)$, $\dot{g} \in L^2(L^2)$, and $(\beta_\alpha, w) \in L^\infty(H_0^1)$ is the solution of (4.4) (or (4.5)) with initial conditions (4.6). Then as $\zeta \rightarrow 0$, there exists a sequence of (β_α, w) with the same notation for simplicity such that

$$\begin{aligned} w &\rightharpoonup \tilde{w} \quad \text{weakly star in } L^\infty(H^2), \\ \beta_\alpha &\rightharpoonup \tilde{\beta}, \quad \dot{w} \rightharpoonup \dot{\tilde{w}} \quad \text{weakly star in } L^\infty(H^1), \\ \ddot{w} &\rightharpoonup \ddot{\tilde{w}} \quad \text{weakly star in } L^\infty(L^2). \end{aligned} \quad (4.7)$$

Moreover,

$$\tilde{\boldsymbol{\beta}} = \nabla \tilde{w} \quad (4.8)$$

and $\tilde{w}$ is the solution of a K-L plate problem of elastodynamics with clamped boundary conditions

$$\begin{aligned} \rho_0 \ddot{\tilde{w}} + D_0 \nabla^4 \tilde{w} &= g \quad \text{or} \\ \rho_0 (\ddot{\tilde{w}}, \nu) + D_0 (\nabla^2 \tilde{w}, \nabla^2 \nu) &= (g, \nu), \quad \forall \nu \in H_0^2, \\ \tilde{w}|_{\partial\Omega} &= \frac{\partial \tilde{w}}{\partial n}|_{\partial\Omega} = 0, \\ \tilde{w}(0, \mathbf{x}) &= \dot{\tilde{w}}(0, \mathbf{x}) = 0. \end{aligned} \quad (4.9)$$

Proof. By Theorems 3.1 to 3.3, we have a unique solution $(\beta_\alpha, w) \in L^\infty(H_0^1) \cap L^\infty(H^2)$ for (4.4) (or (4.5)), where the generic constant $C > 0$ used in the a priori estimates is independent of material parameters. With (4.6), the a priori estimate (3.1) is reduced to

$$\sqrt{I_0 \zeta} \|\dot{\boldsymbol{\beta}}\|_0 + \sqrt{\rho_0} \|\dot{w}\|_0 + \sqrt{E} \|\boldsymbol{\beta}\|_1 + \sqrt{\lambda \zeta^{-2}} \|\nabla w - \boldsymbol{\beta}\|_0 \leq C \|g\|_{L^2(L^2)}. \quad (4.10)$$

Inequality (3.8) yields

$$\sqrt{I_0 \zeta} \|\ddot{\boldsymbol{\beta}}\|_0 + \sqrt{\rho_0} \|\ddot{w}\|_0 + \sqrt{E} \|\ddot{\boldsymbol{\beta}}\|_1 + \sqrt{\lambda \zeta^{-2}} \|\nabla \dot{w} - \dot{\boldsymbol{\beta}}\|_0 \leq C (\|g(0)\|_0 + \|\dot{g}\|_{L^2(L^2)}). \quad (4.11)$$

Inequality (3.10) results in

$$\|w\|_2 \leq C(\|g(0)\|_0 + \|\dot{g}\|_{L^2(L^2)} + \|g\|_{L^2(L^2)}). \quad (4.12)$$

Furthermore, $\|\dot{w}\|_1 \leq C(\|\dot{w}\|_0 + \|\nabla \dot{w}\|_0) \leq C(\|\dot{w}\|_0 + \|\nabla \dot{w} - \dot{\beta}\|_0 + \|\dot{\beta}\|_0)$. Therefore, the boundedness of w in $L^\infty(H^2)$, $\beta_\alpha, \dot{\beta}_\alpha, \dot{w}$ in $L^\infty(H^1)$, and $\zeta \dot{\beta}_\alpha, \ddot{w}$ in $L^\infty(L^2)$ are all uniform with respect to the thickness. We can extract the convergent sequences

$$\begin{aligned} w &\rightharpoonup \tilde{w} \quad \text{weakly star in } L^\infty(H^2), \\ \beta_\alpha &\rightharpoonup \tilde{\beta}_\alpha, \quad \dot{\beta}_\alpha \rightharpoonup \dot{\tilde{\beta}}_\alpha, \quad \dot{w} \rightharpoonup \dot{\tilde{w}} \quad \text{weakly star in } L^\infty(H^1), \\ \ddot{w} &\rightharpoonup \ddot{\tilde{w}} \quad \text{weakly star in } L^\infty(L^2), \end{aligned} \quad (4.13)$$

where, for simplicity, no ζ -dependence notation is used for the sequences. The relation with time differentiation is trivial.

The initial conditions $\tilde{w}(0, \mathbf{x}) = \dot{\tilde{w}}(0, \mathbf{x}) = 0$ are a direct result of (4.6). Inequality (4.10) implies $\sqrt{\lambda} \|\nabla w - \beta\|_0 \leq C\zeta \|g\|_{L^2(L^2)} \rightarrow 0 \Rightarrow \nabla \tilde{w} - \tilde{\beta} = 0$. Meanwhile, the boundary conditions on (β_α, w) lead to $\tilde{\beta}_\alpha|_{\partial\Omega} = \tilde{w}|_{\partial\Omega} = \nabla \tilde{w}|_{\partial\Omega} = 0$. The last equation implies, for smooth domain, $\partial \tilde{w} / \partial n|_{\partial\Omega} = 0$.

On the other hand, by (4.11), $\zeta^2 \|\dot{\beta}\|_0 \rightarrow 0$. Thus, the first two equations of (4.4) yield $EA_\alpha(\beta) - \lambda \zeta^{-2}(w_{,\alpha} - \beta_\alpha) \rightarrow 0$. That means $\lambda \zeta^{-2}(w_{,\alpha} - \beta_\alpha) \rightarrow EA_\alpha(\tilde{\beta}) = EA_\alpha(\nabla \tilde{w})$. Then the third equation of (4.4) gives $\rho_0 \ddot{\tilde{w}} - E \nabla \cdot (\mathbf{A}(\nabla \tilde{w})) = \rho_0 \ddot{\tilde{w}} + D_0 \nabla^4 \tilde{w} = g$. The last equality can be easily verified with the definition of the operator $\mathbf{A}$ and considered in the weak sense. Similar statement for the variational equation is straightforward. $\square$

Remark 4.2. The generic constant C involved in the inequalities derived in Theorem 3.4 is thickness dependent. So the boundedness of β and w in higher spaces may not be uniform with respect to the thickness. It is worth noting that the boundary layer is found for static problems of R-M plate [1]. With clamped boundary conditions, as in our case, $\partial^3 \beta / \partial n^3 = O(\zeta^{-1})$ near the boundary, that is, $\|\beta\|_3 = O(\zeta^{-1/2})$. The boundary layer is expected for the dynamics problem too, which warrants further investigation. Since $\tilde{\beta} = \nabla \tilde{w}$, it is not optimistic that we can have the convergence of $w \rightarrow \tilde{w}$ in the sense of H^4 , although the corresponding K-L plate can have a strong solution $\tilde{w}$ in H^4 .

5. Summary

Existence and uniqueness of H^1 solution of R-M plate for elastodynamics with homogeneous Dirichlet boundary conditions and general initial conditions were proved. The solution with smoother data was further investigated and proved to be in H^2 . Furthermore, with higher smoothness of data and certain compatibility requirements satisfied, higher regularity of the solution was proved. With the introduction of mass scaling, along with the load scaling, the H^2 solution of R-M plate was proved to approach the H^2 weak solution of K-L plate when the thickness approaches zero.

Acknowledgments

The author is greatly indebted to Professor F. Brezzi for the fruitful discussion about the dynamics of Reissner-Mindlin plate and the mass scaling. The author is also pleased to acknowledge the referee's helpful comments.

References

- [1] D. N. Arnold and R. S. Falk, *Edge effect in the Reissner-Mindlin plate theory*, Analytic and Computational Models of Shells (A. K. Noor, T. Belytschko, and J. C. Simo, eds.), A.S.M.E., New York, 1989, pp. 71–90.
- [2] ———, *A uniformly accurate finite element method for the Reissner-Mindlin plate*, SIAM J. Numer. Anal. **26** (1989), no. 6, 1276–1290.
- [3] K. J. Bathe and F. Brezzi, *On the convergence of a four-node plate bending element based on Mindlin-Reissner plate theory and a mixed interpolation*, The Mathematics of Finite Elements and Applications V (Uxbridge, 1984), Academic Press, London, 1985, pp. 491–503.
- [4] F. Brezzi, *On scaling of Reissner-Mindlin plate*, Private communication (2003).
- [5] F. Brezzi and M. Fortin, *Numerical approximation of Mindlin-Reissner plates*, Math. Comp. **47** (1986), no. 175, 151–158.
- [6] P. G. Ciarlet, *Mathematical Elasticity. Vol. I, Three-Dimensional Elasticity*, Elsevier Science Publishing, New York, 1988.
- [7] G. Duvaut and J.-L. Lions, *Inequalities in Mechanics and Physics*, Springer-Verlag, Berlin, 1976.
- [8] L. C. Evans, *Partial Differential Equations*, Graduate Studies in Mathematics, vol. 19, American Mathematical Society, Rhode Island, 1998.
- [9] K. M. Liew, C. M. Wang, Y. Xiang, and S. Kitipornchai, *Vibration of Mindlin Plates, Programming the P-Erson Ritz Method*, Elsevier, Oxford, 1998.
- [10] J. L. Lions and E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications. Vol. I*, Springer-Verlag, New York, 1972.
- [11] ———, *Non-Homogeneous Boundary Value Problems and Applications. Vol. II*, Springer-Verlag, New York, 1972.
- [12] M. Lyly, R. Stenberg, and T. Vihinen, *A stable bilinear element for the Reissner-Mindlin plate model*, Comput. Methods Appl. Mech. Engrg. **110** (1993), no. 3–4, 343–357.
- [13] R. D. Mindlin, *Influence of rotatory inertia and shear on flexural motions of isotropic elastic plates*, J. Appl. Mech. **18** (1951), 31–38.
- [14] E. Reissner, *The effect of transverse shear deformation on the bending of elastic plates*, J. Appl. Mech. **12** (1945), A-69–A-77.
- [15] ———, *On bending of elastic plates*, Quart. Appl. Math. **5** (1947), 55–68.
- [16] M. Suri, I. Babuška, and C. Schwab, *Locking effects in the finite element approximation of plate models*, Math. Comp. **64** (1995), no. 210, 461–482.
- [17] R. Temam, *Navier-Stokes Equations*, 3rd ed., Theory and Numerical Analysis, Elsevier, Amsterdam, 1984.
- [18] Z. Zhang and S. Zhang, *Wilson's element for the Reissner-Mindlin plate*, Comput. Methods Appl. Mech. Engrg. **113** (1994), no. 1–2, 55–65.

Shen Rong Wu: Research and Advanced Engineering, Ford Motor Company, 2101 Village Road, SRL-MD 2115, Dearborn, MI 48124, USA

E-mail address: swu@ford.com