Research Article

Periodic Solutions for Autonomous (*q*, *p*)**-Laplacian System with Impulsive Effects**

Xiaoxia Yang and Haibo Chen

School of Mathematical Sciences and Computing Technology, Central South University, Changsha, Hunan 410083, China

Correspondence should be addressed to Haibo Chen, hbchen2003@gmail.com

Received 17 July 2011; Accepted 31 August 2011

Academic Editor: Yongkun Li

Copyright © 2011 X. Yang and H. Chen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

By using the variational method, some existence theorems are obtained for periodic solutions of autonomous (q, p)-Laplacian system with impulsive effects.

1. Introduction

Let $B = \{1, 2, ..., l\}, C = \{1, 2, ..., k\}, l, k \in \mathbb{N}$. In this paper, we consider the following system:

$$\begin{aligned} \frac{d}{dt} \Phi_q(\dot{u}_1(t)) &= \nabla_{u_1} F(u_1(t), u_2(t)), \quad \text{a.e. } t \in [0, T], \\ \frac{d}{dt} \Phi_p(\dot{u}_2(t)) &= \nabla_{u_2} F(u_1(t), u_2(t)), \quad \text{a.e. } t \in [0, T], \\ u_1(0) - u_1(T) &= \dot{u}_1(0) - \dot{u}_1(T) = 0, \\ u_2(0) - u_2(T) &= \dot{u}_2(0) - \dot{u}_2(T) = 0, \\ \Delta \Phi_q(\dot{u}_1(t_j)) &= \Phi_q(\dot{u}_1(t_j^+)) - \Phi_q(\dot{u}_1(t_j^-)) = \nabla I_j(u_1(t_j)), \quad j \in B, \\ \Delta \Phi_p(\dot{u}_2(s_m)) &= \Phi_p(\dot{u}_2(s_m^+)) - \Phi_p(\dot{u}_2(s_m^-)) = \nabla K_m(u_2(s_m)), \quad m \in C, \end{aligned}$$
(1.1)

where p > 1, q > 1, T > 0, $u(t) = (u_1(t), u_2(t)) = (u_1^1(t), u_1^2(t), \dots, u_1^N(t), u_2^1(t), u_2^2(t), \dots, u_2^N(t))^{\tau}$, $t_j(j = 1, 2, \dots, l)$, and $s_m(m = 1, 2, \dots, k)$ are the instants where the impulses occur

and $0 = t_0 < t_1 < t_2 < \cdots < t_l < t_{l+1} = T$, $0 = s_0 < s_1 < s_2 < \cdots < s_k < s_{k+1} = T$, $I_j : \mathbb{R}^N \to \mathbb{R}$ $(j \in B)$, and $K_m : \mathbb{R}^N \to \mathbb{R}$ $(m \in C)$ are continuously differentiable

$$\Phi_{\mu}(z) = |z|^{\mu-2} z = \left(\sum_{i=1}^{N} z_{i}^{2}\right)^{(\mu-2)/2} \begin{pmatrix} z_{1} \\ \vdots \\ z_{N} \end{pmatrix}, \quad \mu \in \mathbb{R}, \ \mu > 1,$$

$$\nabla I_{j}(x) = \begin{pmatrix} \frac{\partial I_{j}}{\partial x_{1}} \\ \vdots \\ \frac{\partial I_{j}}{\partial x_{N}} \end{pmatrix}, \quad \nabla K_{m}(x) = \begin{pmatrix} \frac{\partial K_{m}}{\partial x_{1}} \\ \vdots \\ \frac{\partial K_{m}}{\partial x_{N}} \end{pmatrix}, \quad (1.2)$$

and $F : \mathbb{R}^N \times \mathbb{R}^N \to \mathbb{R}$ satisfies the following assumption.

(A) F(x) is continuously differentiable in (x_1, x_2) , and there exist $a_1, a_2 \in C(\mathbb{R}^+, \mathbb{R}^+)$ such that

$$|F(x_1, x_2)| \le a_1(|x_1|) + a_2(|x_2|) , \qquad |\nabla F(x_1, x_2)| \le a_1(|x_1|) + a_2(|x_2|),$$

$$|I_j(x_1)| \le a_1(|x_1|), \quad |\nabla I_j(x_1)| \le a_1(|x_1|), \quad j \in B,$$

$$|K_m(x_2)| \le a_2(|x_2|), \quad |\nabla K_m(x_2)| \le a_2(|x_2|), \quad m \in C,$$

(1.3)

for all $x = (x_1, x_2) \in \mathbb{R}^N \times \mathbb{R}^N$.

When p = q = 2, $I_j \equiv 0$ $(j \in B)$, $K_m \equiv 0$ $(m \in C)$, and $F(u_1, u_2) = F_1(u_1)$, system (1.1) reduces to the following autonomous second-order Hamiltonian system:

$$\begin{aligned} \ddot{u}_1(t) &= \nabla_{u_1} F_1(u_1(t)), \quad \text{a.e. } t \in [0, T], \\ u_1(0) &- u_1(T) = \dot{u}_1(0) - \dot{u}_1(T) = 0. \end{aligned}$$
(1.4)

There have been lots of results about the existence of periodic solutions for system (1.4) and nonautonomous second order Hamiltonian system

$$\begin{aligned} \ddot{u}_1(t) &= \nabla_{u_1} F_1(t, u_1(t)), \quad \text{a.e. } t \in [0, T], \\ u_1(0) &- u_1(T) = \dot{u}_1(0) - \dot{u}_1(T) = 0, \end{aligned}$$
(1.5)

(e.g., see [1–21]). Many solvability conditions have been given, for instance, coercive condition, subquadratic condition, superquadratic condition, convex condition, and so on.

When p = q = 2, $\nabla I_j \neq 0$ ($j \in B$), $K_m \equiv 0$ ($m \in C$), and $F(u_1, u_2) = F_1(u_1)$, system (1.1) reduces to the following autonomous second-order Hamiltonian system with impulsive effects:

$$\begin{aligned} \ddot{u}_{1}(t) &= \nabla_{u_{1}} F_{1}(u_{1}(t)), \quad \text{a.e. } t \in [0,T], \\ u_{1}(0) &- u_{1}(T) = \dot{u}_{1}(0) - \dot{u}_{1}(T) = 0, \\ \dot{u}_{1}\left(t_{j}^{+}\right) &- \dot{u}_{1}\left(t_{j}^{-}\right) = \nabla I_{j}(u_{1}(t_{j})). \end{aligned}$$

$$(1.6)$$

Recently, many authors studied the existence of periodic solutions for impulsive differential equations by using variational methods, and lots of interesting results have been obtained. For example, see [22–28]. Especially, nonautonomous second-order Hamiltonian system with impulsive effects is considered in [25, 26] by using the least action principle and the saddle point theorem.

When $I_j \equiv 0$ ($j \in B$) and $K_m \equiv 0$ ($m \in C$), system (1.1) reduces to the following system:

$$\frac{d}{dt}\Phi_{q}(\dot{u}_{1}(t)) = \nabla_{u_{1}}F(u_{1}(t), u_{2}(t)), \quad \text{a.e. } t \in [0, T],$$

$$\frac{d}{dt}\Phi_{p}(\dot{u}_{2}(t)) = \nabla_{u_{2}}F(u_{1}(t), u_{2}(t)), \quad \text{a.e. } t \in [0, T],$$

$$u_{1}(0) - u_{1}(T) = \dot{u}_{1}(0) - \dot{u}_{1}(T) = 0,$$

$$u_{2}(0) - u_{2}(T) = \dot{u}_{2}(0) - \dot{u}_{2}(T) = 0.$$
(1.7)

In [29, 30], Paşca and Tang obtained some existence results for system (1.7) by using the least action principle and saddle point theorem. Motivated by [17, 22–30], in this paper, we are concerned with system (1.1) and also use the least action principle and saddle point theorem to study the existence of periodic solution. Our results still improve those in [17] even if system (1.1) reduces to system (1.4).

A function $G : \mathbb{R}^N \to \mathbb{R}$ is called to be (λ, μ) -quasiconcave if

$$G(\lambda(x+y)) \ge \mu(G(x) + G(y)), \tag{1.8}$$

for some $\lambda, \mu > 0$ and $x, y \in \mathbb{R}^N$.

Next, we state our main results.

Theorem 1.1. Let q' and p' be such that 1/q + 1/q' = 1 and 1/p + 1/p' = 1. Suppose F satisfies assumption (A) and the following conditions:

(F1) there exist

$$0 < r_1 < \frac{(q'+1)^{q/q'}}{T^q \Theta(q,q')}, \qquad 0 < r_2 < \frac{(p'+1)^{p/p'}}{T^p \Theta(p,p')}, \tag{1.9}$$

such that

$$(\nabla_{x_1} F(x_1, x_2) - \nabla_{y_1} F(y_1, y_2), x_1 - y_1) \ge -r_1 |x_1 - y_1|^q, \quad \forall (x_1, x_2), (y_1, y_2) \in \mathbb{R}^N \times \mathbb{R}^N, (\nabla_{x_2} F(x_1, x_2) - \nabla_{y_2} F(y_1, y_2), x_2 - y_2) \ge -r_2 |x_2 - y_2|^p, \quad \forall (x_1, x_2), (y_1, y_2) \in \mathbb{R}^N \times \mathbb{R}^N,$$

$$(1.10)$$

where

$$\Theta(q,q') = \int_0^1 \left[s^{q'+1} + (1-s)^{q'+1} \right]^{q/q'} ds,$$

$$\Theta(p,p') = \int_0^1 \left[s^{p'+1} + (1-s)^{p'+1} \right]^{p/p'} ds,$$
(1.11)

- (F2) $F(x) \to +\infty$, as $|x| \to \infty$, where $x = (x_1, x_2)$,
- (I1) there exists $\beta \in \mathbb{R}$ such that

$$I_{j}(x) \ge \beta, \quad \forall x \in \mathbb{R}^{N}, \ j \in B,$$

$$K_{m}(x) \ge \beta, \quad \forall x \in \mathbb{R}^{N}, \ m \in C.$$
(1.12)

Then, system (1.1) has at least one solution in $W_T^{1,q} \times W_T^{1,p}$, where $W_T^{1,s} = \{u : [0,T] \to \mathbb{R}^N \mid u \text{ is absolutely continuous, } u(0) = u(T) \text{ and } \dot{u} \in L^s(0,T;\mathbb{R}^N)\}, s \in \mathbb{R}.$

Furthermore, if $I_j \equiv 0$ $(j \in B)$, $K_m \equiv 0$ $(m \in C)$ and the following condition holds:

(F3) there exist $\delta > 0$, $a \in [0, (q'+1)^{q/q'}/qT^q\Theta(q,q'))$ and $b \in [0, (p'+1)^{p/p'}/(pT^p\Theta(p,p')))$ such that

$$-a|x_1|^q - b|x_2|^p \le F(x_1, x_2) \le 0, \quad \forall |x_1| \le \delta, \quad |x_2| \le \delta,$$
(1.13)

then system (1.7) has at least two nonzero solutions in $W_T^{1,q} \times W_T^{1,p}$.

When p = q = 2, $F(x_1, x_2) = F_1(x_1)$, by Theorem 1.1, it is easy to get the following corollary.

Corollary 1.2. *Suppose* F_1 *satisfies the following conditions:*

(A)' $F_1(z)$ is continuously differentiable in z and there exists $a_1 \in C(\mathbb{R}^+, \mathbb{R}^+)$ such that

$$|F_1(z)| \le a_1(|z|), \qquad |\nabla F_1(z)| \le a_1(|z|),$$

$$|I_j(z)| \le a_1(|z|), \qquad |\nabla I_j(z)| \le a_1(|z|), \quad j \in B,$$
(1.14)

for all $z \in \mathbb{R}^N$.

(F1)' there exists $0 < r < 6/T^2$ such that

$$(\nabla_z F_1(z) - \nabla_w F_1(w), z - w) \ge -r|z - w|^2, \quad \forall z, w \in \mathbb{R}^N,$$
(1.15)

 $(\mathrm{F2})' \ F_1(z) \ \rightarrow \ +\infty, \ \ as \ \ |z| \ \rightarrow \ \infty, \ \ z \in \mathbb{R}^N;$

(I1)' there exists $\beta \in \mathbb{R}$ such that

$$I_j(z) \ge \beta, \quad \forall z \in \mathbb{R}^N, \quad j \in B.$$
 (1.16)

Then, system (1.6) has at least one solution in $W_T^{1,2}$. Furthermore, if $I_j \equiv 0$ $(j \in B)$ and the following condition holds:

(F3)' there exist $\delta > 0$ and $a \in [0, (3/T^2))$ such that

$$-a|z|^2 \le F_1(z) \le 0, \quad \forall z \in \mathbb{R}^N, \ |z| \le \delta,$$
(1.17)

then system (1.4) has at least two nonzero solutions in $W_T^{1,2}$.

For the Sobolev space $\widetilde{W}_{T}^{1,2}$, one has the following sharp estimates (see in [3, Proposition 1.2]):

$$\int_0^T |u(t)|^2 dt \le \frac{T^2}{4\pi^2} \int_0^T |\dot{u}(t)|^2 dt \quad (\text{Wirtinger's inequality}), \tag{1.18}$$

$$\|u\|_{\infty}^{2} \leq \frac{T}{12} \int_{0}^{T} |\dot{u}(t)|^{2} dt \quad \text{(Sobolev's inequality)}. \tag{1.19}$$

By the above two inequalities, we can obtain the following better results than by Corollary 1.2.

Theorem 1.3. Suppose F_1 satisfies assumption (A)', (F2)', (I1)' and

(F1)" there exists $0 < r < 4\pi^2/T^2$ such that (1.15) holds.

Then, system (1.6) *has at least one solution in* $W_T^{1,2}$. *Furthermore, if* $I_j \equiv 0$ ($j \in B$) *and the following condition holds:*

(F3)" there exist $\delta > 0$ and $a \in [0, (2\pi^2)/T^2)$ such that

$$-a|z|^2 \le F_1(z) \le 0, \quad \forall z \in \mathbb{R}^N, \ |z| \le \delta,$$
(1.20)

then system (1.4) has at least two nonzero solutions in $W_T^{1,2}$.

Moreover, for system (1.6), we have the following additional result.

Theorem 1.4. Suppose F_1 satisfies assumption (A)', (F1)" and the following conditions:

(F4) $F_1(z)$ is (λ, μ) -quasiconcave on \mathbb{R}^N ,

(F5)
$$F_1(z) \to -\infty$$
 as $|z| \to +\infty$, $z \in \mathbb{R}^N$,
(I2) there exist $d_j > 0$ $(j \in B)$ such that

$$|\nabla I_j(z)| \le d_j, \quad \forall z \in \mathbb{R}^N, \ j \in B,$$
 (1.21)

(I3) there exist $b_j > 0$, $c_j > 0$, $\gamma_j \in \mathbb{R}$, $\alpha_j \in [0, 2) (j \in B)$ such that

$$-b_j |z|^{\alpha_j} - c_j \le I_j(z) \le \gamma_j, \quad \forall z \in \mathbb{R}^N, \ j \in B.$$
(1.22)

Then, system (1.6) has at least one solution in $W_T^{1,2}$.

Remark 1.5. In [17], Yang considered the second-order Hamiltonian system with no impulsive effects, that is, system (1.4). When $I_j \equiv 0$ ($j \in B$), our Theorems 1.3 and 1.4 still improve those results in [17]. To be precise, the restriction of r is relaxed, and some unnecessary conditions in [17] are deleted. In [17], the restriction of r is 0 < r < T/12, which is not right. In fact, from his proof, it is easy to see that it should be $0 < r < 12/T^2$. Obviously, our restriction $0 < r < 4\pi^2/T^2$ is better. Moreover, in our Theorem 1.4, we delete such conditions (of in [17, Theorem 1]): $\nabla F_1(0) = 0$, and there exist positive constants M, N such that

$$F_1(z) \ge -M |z|^2 - N, \quad z \in \mathbb{R}^N.$$
 (1.23)

Finally, it is remarkable that Theorems 1.3 and 1.4 are also different from those results in [1–16]. We can find an example satisfying our Theorem 1.3 but not satisfying the results in [1–21]. For example, let

$$F_1(z) = \frac{\pi^2}{2T^2} \left(|z_1|^4 + |z_2|^4 + \dots + |z_N|^4 \right) - \frac{\pi^2}{4T^2} |z|^2,$$
(1.24)

where $z = (z_1, ..., z_N)^{\tau}$. We can also find an example satisfying our Theorem 1.4 but not satisfying the results in [1–21]. For example, let

$$F_1(z) = -\frac{r}{2}|z|^2, \tag{1.25}$$

where $12/T^2 < r < 4\pi^2/T^2$.

2. Variational Structure and Some Preliminaries

The norm in $W_T^{1,p}$ is defined by

$$\|u\|_{W_{T}^{1,p}} = \left[\int_{0}^{T} |u(t)|^{p} dt + \int_{0}^{T} |\dot{u}(t)|^{p} dt\right]^{1/p}.$$
(2.1)

Set

$$\|u\|_{p} = \left(\int_{0}^{T} |u(t)|^{p} dt\right)^{1/p}, \qquad \|u\|_{\infty} = \max_{t \in [0,T]} |u(t)|.$$
(2.2)

Let

$$\widetilde{W}_{T}^{1,p} = \left\{ u \in W_{T}^{1,p} \mid \int_{0}^{T} u(t)dt = 0 \right\} .$$
(2.3)

Obviously, $W_T^{1,p}$ is a reflexive Banach space. It is easy to know that $\widetilde{W}_T^{1,p}$ is a subset of $W_T^{1,p}$ and $W_T^{1,p} = \mathbb{R}^N \oplus \widetilde{W}_T^{1,p}$. In this paper, we will use the space W defined by

$$W = W_T^{1,q} \times W_T^{1,p}, \quad u(t) = (u_1(t), u_2(t)), \tag{2.4}$$

with the norm $\|(u_1, u_2)\|_W = \|u_1\|_{W_T^{1,q}} + \|u_2\|_{W_T^{1,p}}$. It is clear that W is a reflexive Banach space. Let $\widetilde{W} = \widetilde{W}_T^{1,q} \times \widetilde{W}_T^{1,p}$. Then, $W = (\widetilde{W}_T^{1,q} \times \widetilde{W}_T^{1,p}) \oplus (\mathbb{R}^N \times \mathbb{R}^N)$.

Lemma 2.1 (see [31] or [32]). Each $u \in W_T^{1,p}$ and each $v \in W_T^{1,q}$ can be written as $u(t) = \overline{u} + \widetilde{u}(t)$ and $v(t) = \overline{v} + \widetilde{v}(t)$ with

$$\overline{u} = \frac{1}{T} \int_0^T u(t) dt, \qquad \int_0^T \widetilde{u}(t) dt = 0,$$

$$\overline{v} = \frac{1}{T} \int_0^T v(t) dt, \qquad \int_0^T \widetilde{v}(t) dt = 0.$$
(2.5)

Then,

$$\|\widetilde{u}\|_{\infty} \leq \left(\frac{T}{p'+1}\right)^{1/p'} \left(\int_{0}^{T} |\dot{u}(s)|^{p} ds\right)^{1/p}, \qquad \|\widetilde{v}\|_{\infty} \leq \left(\frac{T}{q'+1}\right)^{1/q'} \left(\int_{0}^{T} |\dot{v}(s)|^{q} ds\right)^{1/q},$$
(2.6)

$$\int_{0}^{T} |\tilde{u}(s)|^{p} ds \leq \frac{T^{p}\Theta(p,p')}{\left(p'+1\right)^{p/p'}} \int_{0}^{T} |\dot{u}(s)|^{p} ds, \qquad \int_{0}^{T} |\tilde{v}(s)|^{q} ds \leq \frac{T^{q}\Theta(q,q')}{\left(q'+1\right)^{q/q'}} \int_{0}^{T} |\dot{v}(s)|^{q} ds, \quad (2.7)$$

where

$$\Theta(p,p') = \int_0^1 \left[s^{p'+1} + (1-s)^{p'+1} \right]^{p/p'} ds, \qquad \Theta(q,q') = \int_0^1 \left[s^{q'+1} + (1-s)^{q'+1} \right]^{q/q'} ds.$$
(2.8)

Note that if $u \in W_T^{1,p}$, then u is absolutely continuous. However, we cannot guarantee that \dot{u} is also continuous. Hence, it is possible that $\Delta \Phi_p(\dot{u}(t)) = \Phi_p(\dot{u}(t^+)) - \Phi_p(\dot{u}(t^-)) \neq 0$, which results in impulsive effects.

Following the idea in [22], one takes $v_1 \in W_T^{1,q}$ and multiplies the two sides of

$$\frac{d}{dt}\left(|\dot{u}_1(t)|^{q-2}\dot{u}_1(t)\right) - \nabla_{x_1}F(u_1(t), u_2(t)) = 0,$$
(2.9)

by v_1 and integrate from 0 to *T*, one obtains

$$\int_{0}^{T} \left[\frac{d}{dt} \left(|\dot{u}_{1}(t)|^{q-2} \dot{u}_{1}(t) \right) - \nabla_{x_{1}} F(u_{1}(t), u_{2}(t)) \right] v_{1}(t) dt = 0.$$
(2.10)

Note that $v_1(t)$ is continuous. So, $v_1(t_j^-) = v_1(t_j^+) = v_1(t_j)$. Combining $\dot{u}_1(0) - \dot{u}_1(T) = 0$, one has

$$\begin{split} \int_{0}^{T} \left(\frac{d\Phi_{q}(\dot{u}_{1}(t))}{dt}, v_{1}(t) \right) dt &= \sum_{j=0}^{l} \int_{t_{j}}^{t_{j+1}} \left(\frac{d(\Phi_{q}(\dot{u}_{1}(t)))}{dt}, v_{1}(t) \right) dt \\ &= \sum_{j=0}^{l} \left[\left(\Phi_{q}\left(\dot{u}_{1}\left(t_{j+1}^{-} \right) \right), v_{1}\left(t_{j+1}^{-} \right) \right) - \left(\Phi_{q}\left(\dot{u}_{1}\left(t_{j}^{+} \right) \right), v_{1}\left(t_{j}^{+} \right) \right) \right] dt \\ &- \sum_{j=0}^{l} \int_{t_{j}}^{t_{j+1}} \left(\Phi_{q}(\dot{u}_{1}(t)), \dot{v}_{1}(t) \right) dt \\ &= \left(\Phi_{q}(\dot{u}_{1}(T)), v_{1}(T) \right) - \left(\Phi_{q}(\dot{u}_{1}(0)), v_{1}(0) \right) \\ &- \sum_{j=1}^{l} \left(\Delta \Phi_{q}(\dot{u}_{1}(t_{j})), v_{1}(t_{j}) \right) - \int_{0}^{T} \left(\Phi_{q}(\dot{u}_{1}(t)), \dot{v}_{1}(t) \right) dt \\ &= -\sum_{j=1}^{l} \left(\nabla I_{j}(u_{1}(t_{j})), v_{1}(t_{j}) \right) - \int_{0}^{T} \left(\Phi_{q}(\dot{u}_{1}(t)), \dot{v}_{1}(t) \right) dt. \end{split}$$

$$(2.11)$$

Combining with (2.10), one has

$$\int_{0}^{T} \left(\Phi_{q}(\dot{u}_{1}(t)), \dot{v}_{1}(t) \right) dt + \sum_{j=1}^{l} \left(\nabla I_{j}(u_{1}(t_{j})), v_{1}(t_{j}) \right) + \int_{0}^{T} \left(\nabla_{x_{1}} F(u_{1}(t), u_{2}(t)), v_{1}(t) \right) dt = 0.$$
(2.12)

8

Similarly, one can get

$$\int_{0}^{T} \left(\Phi_{p}(\dot{u}_{2}(t)), \dot{v}_{2}(t) \right) dt + \sum_{m=1}^{k} \left(\nabla K_{m}(u_{2}(s_{m})), v_{2}(s_{m}) \right) + \int_{0}^{T} \left(\nabla_{x_{2}} F(u_{1}(t), u_{2}(t)), v_{2}(t) \right) dt = 0,$$
(2.13)

for all $v_2 \in W_T^{1,p}$. Considering the above equalities, one introduces the following concept of the weak solution for system (1.1).

Definition 2.2. We say that a function $u = (u_1, u_2) \in W_T^{1,q} \times W_T^{1,p}$ is a weak solution of system (1.1) if

$$\int_{0}^{T} \left(\Phi_{q}(\dot{u}_{1}(t)), \dot{v}_{1}(t) \right) dt + \sum_{j=1}^{l} \left(\nabla I_{j}(u_{1}(t_{j})), v_{1}(t_{j}) \right) = -\int_{0}^{T} \left(\nabla_{x_{1}} F(u_{1}(t), u_{2}(t)), v_{1}(t) \right) dt,$$

$$\int_{0}^{T} \left(\Phi_{p}(\dot{u}_{2}(t)), \dot{v}_{2}(t) \right) dt + \sum_{m=1}^{k} \left(\nabla K_{m}(u_{2}(s_{m})), v_{2}(s_{m}) \right) = -\int_{0}^{T} \left(\nabla_{x_{2}} F(u_{1}(t), u_{2}(t)), v_{2}(t) \right) dt$$
(2.14)

holds for any $v = (v_1, v_2) \in W_T^{1,q} \times W_T^{1,p}$. Define the functional $\varphi : W_T^{1,q} \times W_T^{1,p} \to \mathbb{R}$ by

$$\begin{split} \varphi(u_1, u_2) &= \frac{1}{q} \int_0^T |\dot{u}_1(t)|^q dt + \frac{1}{p} \int_0^T |\dot{u}_2(t)|^p dt + \int_0^T F(u_1(t), u_2(t)) dt \\ &+ \sum_{j=1}^l I_j(u_1(t_j)) + \sum_{m=1}^k K_m(u_2(s_m)) \\ &= \phi(u_1, u_2) + \psi(u_1, u_2), \end{split}$$
(2.15)

where $(u_1, u_2) \in W_T^{1,q} \times W_T^{1,p}$,

$$\phi(u_1, u_2) = \frac{1}{q} \int_0^T |\dot{u}_1(t)|^q dt + \frac{1}{p} \int_0^T |\dot{u}_2(t)|^p dt + \int_0^T F(u_1(t), u_2(t)) dt,$$

$$\psi(u_1, u_2) = \sum_{j=1}^l I_j(u_1(t_j)) + \sum_{m=1}^k K_m(u_2(s_m)).$$
(2.16)

By assumption (A) and [33], we know that $\phi \in C^1(W_T^{1,q} \times W_T^{1,p}, \mathbb{R})$. The continuity of $I_j (j \in B)$ and $K_m (m \in C)$ implies that $\psi \in C^1(W_T^{1,p} \times W_T^{1,p}, \mathbb{R})$. So, $\varphi \in C^1(W_T^{1,p}, \mathbb{R})$, and for all $(v_1, v_2) \in W_T^{1,q} \times W_T^{1,p}$, we have

$$\langle \varphi'(u_1, u_2), (v_1, v_2) \rangle = \int_0^T \left(\Phi_q(\dot{u}_1(t)), \dot{v}_1(t) \right) dt + \int_0^T \left(\Phi_p(\dot{u}_2(t)), \dot{v}_2(t) \right) dt + \int_0^T \left(\nabla_{x_1} F(u_1(t), u_2(t)), v_1(t) \right) dt + \int_0^T \left(\nabla_{x_2} F(u_1(t), u_2(t)), v_2(t) \right) dt + \sum_{j=1}^l \left(\nabla I_j(u_1(t_j)), v_1(t_j) \right) + \sum_{m=1}^k \left(\nabla K_m(u_2(s_m)), v_2(s_m) \right).$$

$$(2.17)$$

Definition 2.2 shows that the critical points of φ correspond to the weak solutions of system (1.1).

We will use the following lemma to seek the critical point of φ .

Lemma 2.3 (see [3, Theorem 1.1]). If φ is weakly lower semicontinuous on a reflexive Banach space *X* and has a bounded minimizing sequence, then φ has a minimum on *X*.

Lemma 2.4 (see [34]). Let φ be a C^1 function on $X = X_1 \oplus X_2$ with $\varphi(0) = 0$, satisfying (PS) condition, and assume that for some $\rho > 0$,

$$\begin{aligned} \varphi(u) \ge 0, & \text{for } u \in X_1, \quad \|u\| \le \rho, \\ \varphi(u) \le 0, & \text{for } u \in X_2, \quad \|u\| \le \rho. \end{aligned}$$
(2.18)

Assume also that φ is bounded below and $\inf_X \varphi < 0$, then φ has at least two nonzero critical points.

Lemma 2.5 (see [35, Theorem 4.6]). Let $X = X_1 \oplus X_2$, where X is a real Banach space and $X_1 \neq \{0\}$ and is finite dimensional. Suppose that $\varphi \in C^1(X, \mathbb{R})$ satisfies (PS)-condition and

(φ 1) there is a constant α and a bounded neighborhood D of 0 in X_1 such that $\varphi|_{\partial D} \leq \alpha$,

(φ 2) there is a constant $\beta > \alpha$ such that $\varphi|_{X_2} \ge \beta$.

Then, φ *possesses a critical value* $c \ge \beta$ *. Moreover,* c *can be characterized as*

$$c = \inf_{h \in \Gamma} \max_{u \in \overline{D}} \varphi(h(u)), \tag{2.19}$$

where,

$$\Gamma = \left\{ h \in C\left(\overline{D}, X\right) \mid h = id \text{ on } \partial D \right\}.$$
(2.20)

3. Proof of Theorems

Lemma 3.1. Under assumption (A), φ is weakly lower semicontinuous on $W_T^{1,q} \times W_T^{1,p}$.

Proof. Let

$$\phi_1(u_1, u_2) = \frac{1}{q} \int_0^T |\dot{u}_1(t)|^q dt + \frac{1}{p} \int_0^T |\dot{u}_2(t)|^p dt,$$

$$\phi_2(u_1, u_2) = \int_0^T F(u_1(t), u_2(t)) dt.$$
(3.1)

Since

$$\begin{split} \phi_1\Big(\frac{u_1+v_1}{2},\frac{u_2+v_2}{2}\Big) &= \frac{1}{q} \int_0^T \left|\frac{\dot{u}_1(t)+\dot{v}_1(t)}{2}\right|^q dt + \frac{1}{p} \int_0^T \left|\frac{\dot{u}_2(t)+\dot{v}_2(t)}{2}\right|^p dt \\ &\leq \frac{2^{q-1}}{q} \int_0^T \frac{1}{2^q} |\dot{u}_1(t)|^q dt + \frac{2^{q-1}}{q} \int_0^T \frac{1}{2^q} |\dot{v}_1(t)|^q dt \\ &\quad + \frac{2^{p-1}}{p} \int_0^T \frac{1}{2^p} |\dot{u}_2(t)|^p dt + \frac{2^{p-1}}{p} \int_0^T \frac{1}{2^p} |\dot{v}_2(t)|^p dt \\ &\leq \frac{1}{2q} \int_0^T |\dot{u}_1(t)|^q dt + \frac{1}{2q} \int_0^T |\dot{v}_1(t)|^q dt \\ &\quad + \frac{1}{2p} \int_0^T |\dot{u}_2(t)|^p dt + \frac{1}{2p} \int_0^T |\dot{v}_2(t)|^p dt \\ &= \frac{\phi_1(u_1,u_2) + \phi_1(v_1,v_2)}{2}, \end{split}$$
(3.2)

then ϕ_1 is convex. Moreover, by [33], we know that ϕ_1 is continuous, and so, it is lower semicontinuous. Thus, it follows from [3, Theorem 1.2] that ϕ_1 is weakly lower continuous. By assumption (A), it is easy to verify that $\phi_2(u_1, u_2)$ is weakly continuous. We omit the details. Let

$$\psi_1(u_1) = \sum_{j=1}^l I_j(u_1(t_j)), \qquad \psi_2(u_2) = \sum_{m=1}^k K_m(u_2(s_m)).$$
(3.3)

Next, we show that ψ_1 and ψ_2 are weakly continuous on $W_T^{1,q}$ and $W_T^{1,p}$, respectively. In fact, if

 $u_{1n} \rightarrow u_1$ weakly in $W_T^{1,p}$, as $n \rightarrow \infty$, (3.4)

then by in [3, Proposition 1.2], we know that

$$u_{1n} \longrightarrow u_1$$
 strongly in $C(0,T;\mathbb{R}^N)$, as $n \longrightarrow \infty$. (3.5)

So, there exists $M_1 > 0$ such that $||u_1||_{\infty} \le M_1$ and $||u_{1n}||_{\infty} \le M_1$, for all $n \in \mathbb{N}$. Thus, we have

$$\begin{aligned} |\psi_{1}(u_{1n}) - \psi_{1}(u_{1})| &= \left| \sum_{j=1}^{l} I_{j}(u_{1n}(t_{j})) - \sum_{j=1}^{l} I_{j}(u_{1}(t_{j})) \right| \\ &\leq \sum_{j=1}^{l} \left| I_{j}(u_{1n}(t_{j})) - I_{j}(u_{1}(t_{j})) \right| \\ &= \sum_{j=1}^{l} \left| \int_{0}^{1} (\nabla I_{j}(u_{1}(t_{j}) + s(u_{1n}(t_{j}) - u_{1}(t_{j}))), u_{1n}(t_{j}) - u_{1}(t_{j})) ds \right| \end{aligned}$$
(3.6)
$$&\leq \|u_{1n} - u_{1}\|_{\infty} \sum_{j=1}^{l} \max_{t \in [0, 3M_{1}]} a_{1}(t) \longrightarrow 0.$$

Hence, ψ_1 is weakly continuous on $W_T^{1,q}$. Similarly, we can prove that ψ_2 is also weakly continuous on $W_T^{1,p}$. Thus, we complete the proof.

Proof of Theorem 1.1. It follows from (F1) and (2.7) that

$$\int_{0}^{T} \left[F(u_{1}(t), u_{2}(t)) - F(u_{1}(t), \overline{u}_{2}) \right]$$

$$= \int_{0}^{T} \int_{0}^{1} \frac{1}{s} (\nabla F_{x_{2}}(u_{1}(t), \overline{u}_{2} + s\widetilde{u}_{2}(t)), s\widetilde{u}_{2}(t)) ds dt$$

$$= \int_{0}^{T} \int_{0}^{1} \frac{1}{s} (\nabla F_{x_{2}}(u_{1}(t), \overline{u}_{2} + s\widetilde{u}_{2}(t)) - \nabla F_{x_{2}}(\overline{u}_{1}, \overline{u}_{2}), s\widetilde{u}_{2}(t)) ds dt$$

$$\geq -\frac{r_{2}}{p} \int_{0}^{T} |\widetilde{u}_{2}(t)|^{p} dt$$

$$\geq -\frac{r_{2}T^{p}\Theta(p, p')}{p(p'+1)^{p/p'}} \int_{0}^{T} |\dot{u}_{2}(t)|^{p} dt, \quad \forall (u_{1}, u_{2}) \in W,$$

$$\int_{0}^{T} \left[F(u_{1}(t), \overline{u}_{2}) - F(\overline{u}_{1}, \overline{u}_{2}) \right] dt$$

$$= \int_{0}^{T} \int_{0}^{1} \frac{1}{s} (\nabla_{x_{1}}F(\overline{u}_{1} + s\widetilde{u}_{1}(t), \overline{u}_{2}), s\widetilde{u}_{1}(t)) ds dt$$

$$= \int_{0}^{T} \int_{0}^{1} \frac{1}{s} (\nabla_{x_{1}}F(\overline{u}_{1} + s\widetilde{u}_{1}(t), \overline{u}_{2}) - \nabla_{x_{1}}F(\overline{u}_{1}, \overline{u}_{2}), s\widetilde{u}_{1}(t)) ds dt$$

$$\geq -\frac{r_1}{q} \int_0^T |\tilde{u}_1(t)|^q dt$$

$$\geq -\frac{r_1 T^q \Theta(q, q')}{q (q'+1)^{q/q'}} \int_0^T |\dot{u}_1(t)|^q dt, \quad \forall (u_1, u_2) \in W.$$

(3.8)

Hence, by (I1), (3.7), and (3.8), we have

$$\begin{split} \varphi(u_{1},u_{2}) &= \frac{1}{q} \int_{0}^{T} |\dot{u}_{1}(t)|^{q} dt + \frac{1}{p} \int_{0}^{T} |\dot{u}_{2}(t)|^{p} dt + \int_{0}^{T} [F(u_{1}(t),u_{2}(t)) - F(u_{1}(t),\overline{u}_{2})] dt \\ &+ \int_{0}^{T} [F(u_{1}(t),\overline{u}_{2}) - F(\overline{u}_{1},\overline{u}_{2})] dt + TF(\overline{u}_{1},\overline{u}_{2}) + \sum_{j=1}^{l} I_{j}(u_{1}(t_{j})) + \sum_{m=1}^{k} K_{m}(u_{2}(s_{m})) \\ &\geq \left(\frac{1}{p} - \frac{r_{2}T^{p}\Theta(p,p')}{p(p'+1)^{p/p'}}\right) \int_{0}^{T} |\dot{u}_{2}(t)|^{p} dt + \left(\frac{1}{q} - \frac{r_{1}T^{q}\Theta(q,q')}{q(q'+1)^{q/q'}}\right) \int_{0}^{T} |\dot{u}_{1}(t)|^{q} dt \\ &+ TF(\overline{u}_{1},\overline{u}_{2}) - (l+k)|\beta|. \end{split}$$

$$(3.9)$$

Note that for $u \in W_T^{1,p}$,

$$\|u\|_{W_{T}^{1,p}} \longrightarrow \infty \iff \left(|\overline{u}|^{p} + \int_{0}^{T} |\dot{u}(t)|^{p} dt\right)^{1/p} \longrightarrow \infty,$$
(3.10)

and for $v \in W_T^{1,q}$,

$$\|v\|_{W_T^{1,q}} \longrightarrow \infty \Longleftrightarrow \left(|\overline{v}|^q + \int_0^T |\dot{v}(t)|^q dt \right)^{1/q} \longrightarrow \infty.$$
(3.11)

So, (F2) and (3.9) imply that

$$\varphi(u_1, u_2) \longrightarrow +\infty, \text{ as } ||(u_1, u_2)||_W \longrightarrow \infty.$$
 (3.12)

Thus, by Lemma 2.3, we know that φ has at least one critical point which minimizes φ on W. Furthermore, if $I_j(u_1(t_j)) \equiv 0$ $(j \in B)$ and $K_m(u_2(s_m)) \equiv 0$ $(m \in C)$, then system (1.1)

reduces to (1.7). When (F3) also holds, we will use Lemma 2.4 to obtain more critical points of φ . Let X = W, $X_2 = \mathbb{R}^N \times \mathbb{R}^N$ and $X_1 = \widetilde{W} = \widetilde{W}_T^{1,q} \times \widetilde{W}_T^{1,p}$. By (3.9), we know that $\varphi(u_1, u_2) \to +\infty$ as $||(u_1, u_2)||_W \to \infty$. So, φ satisfies (PS)

condition and is bounded below. Take $\rho = \delta/c_1$, where c_1 is a positive constant such that

 $||u_1||_{\infty} \leq c_1 ||u_1||_{W_T^{1,q}} \leq c_1 ||u||_W$ and $||u_2||_{\infty} \leq c_1 ||u_2||_{W_T^{1,p}} \leq c_1 ||u||_W$ for all $(u_1, u_2) \in W$. It follows from (F3) and Lemma 2.1 that

$$\begin{split} \varphi(u_{1}, u_{2}) &= \frac{1}{q} \int_{0}^{T} |\dot{u}_{1}(t)|^{q} dt + \frac{1}{p} \int_{0}^{T} |\dot{u}_{2}(t)|^{p} dt + \int_{0}^{T} F(u_{1}(t), u_{2}(t)) dt \\ &\geq \frac{1}{q} \int_{0}^{T} |\dot{u}_{1}(t)|^{q} dt + \frac{1}{p} \int_{0}^{T} |\dot{u}_{2}(t)|^{p} dt - a \int_{0}^{T} |u_{1}(t)|^{q} dt - b \int_{0}^{T} |u_{2}(t)|^{p} dt \\ &\geq \frac{1}{q} \int_{0}^{T} |\dot{u}_{1}(t)|^{q} dt + \frac{1}{p} \int_{0}^{T} |\dot{u}_{2}(t)|^{p} dt - a \frac{T^{q} \Theta(q, q')}{(q'+1)^{q/q'}} \int_{0}^{T} |\dot{u}_{1}(t)|^{q} dt \\ &- b \frac{T^{p} \Theta(p, p')}{(p'+1)^{p/p'}} \int_{0}^{T} |\dot{u}_{2}(t)|^{p} dt, \quad \forall (u_{1}, u_{2}) \in X_{1}. \end{split}$$

$$(3.13)$$

Since $a \leq (q'+1)^{q/q'}/(qT^q\Theta(q,q'))$ and $b \leq (p'+1)^{p/p'}/(pT^p\Theta(p,p'))$, (3.13) implies that $\varphi(u_1, u_2) \geq 0$ for all $(u_1, u_2) \in X_1$ with $||u||_W \leq \rho$. By (F3), it is easy to obtain that $\varphi(u_1, u_2) \leq 0$, for all $(u_1, u_2) \in X_2$ with $||u||_W \leq \rho$.

If $\inf\{\varphi(u_1, u_2) : (u_1, u_2) \in W\} = 0$, then from above, we have $\varphi(u_1, u_2) = 0$ for all $(u_1, u_2) \in X_2$ with $||(u_1, u_2)||_W \leq \rho$. Hence, all $(u_1, u_2) \in X_2$ with $||(u_1, u_2)||_W \leq \rho$ are minimal points of φ , which implies that φ has infinitely many critical points. If $\inf\{\varphi(u_1, u_2) : (u_1, u_2) \in W\} < 0$, then by Lemma 2.4, φ has at least two nonzero critical points. Hence, system (1.7) has at least two nontrivial solutions in *W*. We complete our proof.

Proof of Theorem 1.3. We only need to use (1.18) and (1.19) to replace (2.6) and (2.7) in the proof Theorem 1.1 with p = q = 2, $F(t, u_1, u_2) = F_1(u_1)$ and $K_m(u_2) \equiv 0$ ($m \in C$). It is easy. So, we omit it.

Lemma 3.2. Under the assumptions of Theorem 1.4, the functional φ_1 defined by

$$\varphi_1(u_1) = \frac{1}{2} \int_0^T |\dot{u}_1(t)|^2 dt + \int_0^T F_1(u_1(t)) dt + \sum_{j=1}^l I_j(u_1(t_j)) dt$$
(3.14)

satisfies (PS) condition.

Proof. Suppose that $\{u_{1n}\}$ is a (PS) sequence for φ_1 ; that is, there exists $D_1 > 0$ such that

$$|\varphi(u_{1n})| \le D_1, \quad \forall n \in \mathbb{N}, \quad \varphi'(u_{1n}) \longrightarrow 0, \quad \text{as } n \longrightarrow \infty.$$
 (3.15)

Hence, for *n* large enough, we have $\|\varphi'(u_{1n})\| \leq 1$. It follows from (F1)^{*n*}, (I2), and (1.18) that

$$\begin{aligned} \|\widetilde{u}_{1n}\|_{W_{T}^{1,2}} &\geq \left\langle \varphi_{1}'(u_{1n}), \widetilde{u}_{1n} \right\rangle = \int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt + \int_{0}^{T} (\nabla_{x_{1}} F_{1}(u_{1n}(t)), \widetilde{u}_{1n}(t)) dt \\ &+ \sum_{j=1}^{l} (\nabla I_{j}(u_{1n}(t_{j})), \widetilde{u}_{1n}(t_{j})) \end{aligned}$$

$$= \int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt + \int_{0}^{T} (\nabla_{x_{1}} F_{1}(u_{1n}(t)) - \nabla_{x_{1}} F_{1}(\overline{u}_{1n}(t)), \widetilde{u}_{1n}(t)) dt + \sum_{j=1}^{l} (\nabla I_{j}(u_{1n}(t_{j})), \widetilde{u}_{1n}(t_{j})) \geq \int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt - r \frac{T^{2}}{4\pi^{2}} \int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt - \|\widetilde{u}_{1n}\|_{\infty} \sum_{j=1}^{l} d_{j} \geq \left[1 - r \frac{T^{2}}{4\pi^{2}}\right] \int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt - \left(\frac{T}{12}\right)^{1/2} \left(\int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt\right)^{1/2} \sum_{j=1}^{l} d_{j},$$
(3.16)

for n large enough. By (1.18), we have

$$\|\tilde{u}_{1n}\|_{W_{T}^{1,2}} \leq \left[\frac{T^{2}}{4\pi^{2}} + 1\right]^{1/2} \left(\int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt\right)^{1/2},$$
(3.17)

and (3.16), (3.17), and $r < 4\pi^2/T^2$ imply that there exists $D_2, D_3 > 0$ such that

$$\int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt \leq D_{2}, \quad \|\widetilde{u}_{1n}\|_{W_{T}^{1,2}} \leq D_{3}.$$
(3.18)

It follows from (F4), (3.15), (I3), (1.18), and (3.18) that

$$\begin{aligned} -\mathrm{D}_{1} &\leq \varphi_{1}(u_{1n}) = \frac{1}{2} \int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt + \int_{0}^{T} F_{1}(u_{1n}(t)) dt + \sum_{j=1}^{l} I_{j}(u_{1}(t_{j})) \\ &\leq \frac{1}{2} \int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt + \frac{1}{\mu} \int_{0}^{T} F_{1}(\lambda \overline{u}_{1n}) dt - \int_{0}^{T} F_{1}(-\widetilde{u}_{1n}(t)) dt + \sum_{j=1}^{l} \gamma_{j} \\ &= \frac{1}{2} \int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt + \frac{T}{\mu} F_{1}(\lambda \overline{u}_{1n}) - TF_{1}(0) - \int_{0}^{T} [F_{1}(-\widetilde{u}_{1n}(t)) - F_{1}(0)] dt + \sum_{j=1}^{l} \gamma_{j} \\ &= \frac{1}{2} \int_{0}^{T} |\dot{u}_{1n}(t)|^{2} dt + \frac{T}{\mu} F_{1}(\lambda \overline{u}_{1n}) - TF_{1}(0) + \sum_{j=1}^{l} \gamma_{j} \end{aligned}$$

$$\begin{split} &-\int_{0}^{T}\int_{0}^{1}\frac{1}{s}(\nabla F_{1}(-s\widetilde{u}_{1n}(t))-\nabla F_{1}(0),-s\widetilde{u}_{1n}(t))ds\,dt\\ &\leq \frac{1}{2}\int_{0}^{T}|\dot{u}_{1n}(t)|^{2}dt+\frac{T}{\mu}F_{1}(\lambda\overline{u}_{1n})+r\int_{0}^{T}\int_{0}^{1}s|\widetilde{u}_{1n}(t)|^{2}ds\,dt-TF_{1}(0)+\sum_{j=1}^{l}\gamma_{j}\\ &\leq \frac{1}{2}\int_{0}^{T}|\dot{u}_{1n}(t)|^{2}dt+\frac{T}{\mu}F_{1}(\lambda\overline{u}_{1n})+\frac{r}{2}\int_{0}^{T}|\widetilde{u}_{1n}(t)|^{2}dt-TF_{1}(0)+\sum_{j=1}^{l}\gamma_{j}\\ &\leq \frac{\max\{1,r\}}{2}\|\widetilde{u}_{1n}\|_{W_{T}^{1,2}}^{2}+\frac{T}{\mu}F_{1}(\lambda\overline{u}_{1n})-TF_{1}(0)+\sum_{j=1}^{l}\gamma_{j}\\ &\leq \frac{\max\{1,r\}}{2}D_{3}^{q}+\frac{T}{\mu}F_{1}(\lambda\overline{u}_{1n})-TF_{1}(0)+\sum_{j=1}^{l}\gamma_{j}, \end{split}$$

$$(3.19)$$

for all *n* and (3.19) and (F5) imply that $\{\overline{u}_{1n}\}\)$ is bounded. Combining (3.18), we know that $\{u_{1n}\}\)$ is a bounded sequence. Similar to the argument in [25], it is easy to obtain that φ satisfies (PS) condition.

Proof of Theorem 1.4. From (I3) and (F5), it is easy to see that for $x_1 \in \mathbb{R}^N$,

$$\varphi_1(x_1) \longrightarrow -\infty, \quad \text{as } |x_1| \longrightarrow \infty.$$
 (3.20)

For all $u_1 \in \widetilde{W}_T^{1,2}$, by (1.18), (F1)["] and (I3), we have

$$\begin{split} \varphi_{1}(u_{1}) &= \frac{1}{2} \int_{0}^{T} |\dot{u}_{1}(t)|^{2} dt + \int_{0}^{T} F_{1}(u_{1}(t)) dt + \sum_{j=1}^{l} I_{j}(u_{1}(t_{j})) \\ &= \frac{1}{2} \int_{0}^{T} |\dot{u}_{1}(t)|^{2} dt + \int_{0}^{T} [F_{1}(u_{1}(t)) - F_{1}(0)] dt + TF_{1}(0) + \sum_{j=1}^{l} I_{j}(u_{1}(t_{j})) \\ &= \frac{1}{2} \int_{0}^{T} |\dot{u}_{1}(t)|^{2} dt + \int_{0}^{T} \int_{0}^{1} (\nabla F_{1x_{1}}(su_{1}(t)), u_{1}(t)) ds dt + \sum_{j=1}^{l} I_{j}(u_{1}(t_{j})) + TF_{1}(0) \\ &= \frac{1}{2} \int_{0}^{T} |\dot{u}_{1}(t)|^{2} dt + \int_{0}^{T} \int_{0}^{1} \frac{1}{s} (\nabla F_{1x_{1}}(su_{1}(t)) - \nabla F_{1x_{1}}(0), su_{1}(t)) ds dt \\ &+ \sum_{j=1}^{l} I_{j}(u_{1}(t_{j})) + TF_{1}(0) \end{split}$$

$$\geq \frac{1}{2} \int_{0}^{T} |\dot{u}_{1}(t)|^{2} dt - \frac{r_{1}}{2} \int_{0}^{T} |u_{1}(t)|^{2} dt + TF_{1}(0) - \sum_{j=1}^{l} b_{j} |u_{1}(t_{j})|^{\alpha_{j}} - \sum_{j=1}^{l} c_{j}$$

$$\geq \frac{1}{2} \int_{0}^{T} |\dot{u}_{1}(t)|^{2} dt - \frac{r_{1}T^{2}}{4\pi^{2}} \int_{0}^{T} |\dot{u}_{1}(t)|^{2} dt + TF_{1}(0) - \sum_{j=1}^{l} b_{j} ||u_{1}||_{\infty}^{\alpha_{j}} - \sum_{j=1}^{l} c_{j}$$

$$\geq \left(\frac{1}{2} - \frac{r_{1}T^{2}}{4\pi^{2}}\right) \int_{0}^{T} |\dot{u}_{1}(t)|^{2} dt + TF_{1}(0)$$

$$- \left(\frac{T}{12}\right)^{\alpha_{j}/2} \sum_{j=1}^{l} b_{j} \left(\int_{0}^{T} |\dot{u}_{1}(t)|^{2} dt\right)^{\alpha_{j}/2} - \sum_{j=1}^{l} c_{j}.$$
(3.21)

Note that for all $u_1 \in \widetilde{W}_T^{1,2}$, $||u_1||_{W_T^{1,2}}$ is equivalent to $||\dot{u}_1||_{L^2}$. Then, $r_1 < 4\pi^2/T^2$, $\alpha_j < 2(j \in B)$ and (3.21) imply that

$$\varphi_1(u_1) \longrightarrow +\infty, \quad \text{as } \|u_1\|_{W_T^{1,2}} \longrightarrow \infty, \quad u_1 \in \widetilde{W}_T^{1,2}.$$
 (3.22)

It follows from (3.20) and (3.22) that φ_1 satisfies (φ_1) and (φ_2) in Lemma 2.5. Combining with Lemma 3.2, Lemma 2.5 shows that φ_1 has at least one critical point. Thus, we complete the proof.

4. Examples

Example 4.1. Let q = 4, p = 2, $T = \pi$, $t_1 = 1$, and $s_1 = 2$. Consider the following system:

$$\begin{aligned} \frac{d}{dt} \Phi_4(\dot{u}_1(t)) &= \nabla_{u_1} F(u_1(t) , u_2(t)), \quad \text{a.e. } t \in [0, \pi], \\ \frac{d}{dt} \Phi_2(\dot{u}_2(t)) &= \nabla_{u_2} F(u_1(t), u_2(t)), \quad \text{a.e. } t \in [0, \pi], \\ u_1(0) - u_1(\pi) &= \dot{u}_1(0) - \dot{u}_1(\pi) = 0, \\ u_2(0) - u_2(\pi) &= \dot{u}_2(0) - \dot{u}_2(\pi) = 0, \\ \Delta \Phi_4(\dot{u}_1(1)) &= \Phi_q(\dot{u}_1(1^+)) - \Phi_q(\dot{u}_1(1^-)) = \nabla I_1(u_1(1)), \\ \Delta \Phi_2(\dot{u}_2(2)) &= \Phi_p(\dot{u}_2(2^+)) - \Phi_p(\dot{u}_2(2^-)) = \nabla K_1(u_2(2)), \end{aligned}$$
(4.1)

where $F(x_1, x_2) = x_{11}^4 + x_{12}^4 + \dots + x_{1N}^4 + (1/\pi^2)(x_{21}^4 + x_{22}^2 + \dots + x_{2N}^2) - (1/2\pi^2)|x_2|^2$, $x_1 = (x_{11}, x_{12}, \dots, x_{1N}), x_2 = (x_{21}, x_{22}, \dots, x_{2N}), I_1(x) = e^{|x|^2}, K_1(x) = e^{|x|^2}, x \in \mathbb{R}^N$. It is easy to verify that all conditions of Theorem 1.1 hold so that system (4.1) has at least one weak solution. Moreover, if $F(x_1, x_2) = (1/\pi^2)(x_{21}^4 + x_{22}^4 + \dots + x_{2N}^4) - 1/2\pi^2|x_2|^2$, $x_2 = (x_{21}, x_{22}, \dots, x_{2N}), I_1(x) = 0$ and $K_1(x) = 0, x \in \mathbb{R}^N$, then system (4.1) has at least two nonzero solutions. *Example 4.2.* Let T = 2, $t_1 = 1$. Consider the following autonomous second-order Hamiltonian system with impulsive effects:

$$\begin{aligned} \ddot{u}(t) &= \nabla_{u} F(u(t)), \quad \text{a.e. } t \in [0, 2], \\ u(0) - u(2) &= \dot{u}(0) - \dot{u}(2) = 0, \\ \dot{u}(1^{+}) - \dot{u}(1^{-}) &= \nabla I_{1}(u(1)), \end{aligned}$$

$$(4.2)$$

where $F(z) = z_1^4 + z_2^2 + \dots + z_N^2 - 1/2|z|^2$, $I_1(z) = e^{|z|^2}$, $z = (z_1, \dots, z_N)^T \in \mathbb{R}^N$. It is easy to verify that all conditions of Theorem 1.3 hold so that system (4.2) has at least one weak solution. Moreover, if $F(z) = z_1^4 + z_2^4 + \dots + z_N^4 - 1/2|z|^2$ and $I_1(z) = 0$, $z \in \mathbb{R}^N$, then system (4.2) has at least two nonzero solutions.

Example 4.3. Let $T = \pi$, $t_1 = 2$. Consider the following autonomous second-order Hamiltonian system with impulsive effects:

$$\begin{aligned} \ddot{u}(t) &= \nabla_{u} F(u(t)), \quad \text{a.e. } t \in [0, \pi], \\ u(0) - u(\pi) &= \dot{u}(0) - \dot{u}(\pi) = 0, \\ \dot{u}(2^{+}) - \dot{u}(2^{-}) &= \nabla I_{1}(u(2)), \end{aligned}$$
(4.3)

where $F(z) = -|z|^2$, $I_1(z) = 2 \sin z_1$, $z = (z_1, ..., z_N)^{\tau} \in \mathbb{R}^N$. It is easy to verify that all conditions of Theorem 1.4 hold so that system (4.3) has at least one weak solution.

References

- M. S. Berger and M. Schechter, "On the solvability of semilinear gradient operator equations," Advances in Mathematics, vol. 25, no. 2, pp. 97–132, 1977.
- J. Mawhin, "Semicoercive monotone variational problems," Académie Royale de Belgique, vol. 73, no. 3-4, pp. 118–130, 1987.
- [3] J. Mawhin and M. Willem, Critical Point Theory and Hamiltonian Systems, vol. 74 of Applied Mathematical Sciences, Springer, New York, NY, USA, 1989.
- [4] J. Mawhin and M. Willem, "Critical points of convex perturbations of some indefinite quadratic forms and semilinear boundary value problems at resonance," *Annales de l'Institut Henri Poincaré*, vol. 3, no. 6, pp. 431–453, 1986.
- [5] C. Tang, "Periodic solutions of non-autonomous second order systems with γ-quasisubadditive potential," *Journal of Mathematical Analysis and Applications*, vol. 189, no. 3, pp. 671–675, 1995.
- [6] C. L. Tang, "Periodic solutions of nonautonomous second order systems," Journal of Mathematical Analysis and Applications, vol. 202, pp. 465–469, 1996.
- [7] C.-L. Tang, "Periodic solutions for nonautonomous second order systems with sublinear nonlinearity," *Proceedings of the American Mathematical Society*, vol. 126, no. 11, pp. 3263–3270, 1998.
- [8] C.-L. Tang and X.-P. Wu, "Periodic solutions for second order systems with not uniformly coercive potential," *Journal of Mathematical Analysis and Applications*, vol. 259, no. 2, pp. 386–397, 2001.
- [9] M. Willem, "Oscillations forces de systmes hamiltoniens," in Public. Smin. Analyse Non Linaire, Université de Besançon, 1981.
- [10] X. Wu, "Saddle point characterization and multiplicity of periodic solutions of non-autonomous second-order systems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 58, no. 7-8, pp. 899– 907, 2004.
- [11] X.-P. Wu and C.-L. Tang, "Periodic solutions of a class of non-autonomous second-order systems," *Journal of Mathematical Analysis and Applications*, vol. 236, no. 2, pp. 227–235, 1999.

- [12] F. Zhao and X. Wu, "Periodic solutions for a class of non-autonomous second order systems," *Journal of Mathematical Analysis and Applications*, vol. 296, no. 2, pp. 422–434, 2004.
- [13] F. Zhao and X. Wu, "Existence and multiplicity of periodic solution for non-autonomous second-order systems with linear nonlinearity," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 60, no. 2, pp. 325–335, 2005.
- [14] C.-L. Tang and X.-P. Wu, "A note on periodic solutions of nonautonomous second-order systems," Proceedings of the American Mathematical Society, vol. 132, no. 5, pp. 1295–1303, 2004.
- [15] X. H. Tang and Q. Meng, "Solutions of a second-order Hamiltonian system with periodic boundary conditions," *Nonlinear Analysis: Real World Applications*, vol. 11, no. 5, pp. 3722–3733, 2010.
- [16] Z. Wang and J. Zhang, "Periodic solutions of a class of second order non-autonomous Hamiltonian systems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 12, pp. 4480–4487, 2010.
- [17] R. Yang, "Periodic solutions of some autonomous second order Hamiltonian systems," Journal of Applied Mathematics and Computing, vol. 28, no. 1-2, pp. 51–58, 2008.
- [18] M. Schechter, "Periodic non-autonomous second-order dynamical systems," Journal of Differential Equations, vol. 223, no. 2, pp. 290–302, 2006.
- [19] M. Schechter, "Periodic solutions of second-order nonautonomous dynamical systems," Boundary Value Problems, Article ID 25104, 9 pages, 2006.
- [20] J. Ma and C.-L. Tang, "Periodic solutions for some nonautonomous second-order systems," Journal of Mathematical Analysis and Applications, vol. 275, no. 2, pp. 482–494, 2002.
- [21] F. Zhao and X. Wu, "Saddle point reduction method for some non-autonomous second order systems," *Journal of Mathematical Analysis and Applications*, vol. 291, no. 2, pp. 653–665, 2004.
- [22] J. J. Nieto and D. O'Regan, "Variational approach to impulsive differential equations," Nonlinear Analysis: Real World Applications, vol. 10, no. 2, pp. 680–690, 2009.
- [23] Y. Tian and W. Ge, "Applications of variational methods to boundary-value problem for impulsive differential equations," *Proceedings of the Edinburgh Mathematical Society. Series II*, vol. 51, no. 2, pp. 509–527, 2008.
- [24] J. Zhou and Y. Li, "Existence and multiplicity of solutions for some Dirichlet problems with impulsive effects," Nonlinear Analysis: Theory, Methods & Applications, vol. 71, no. 7-8, pp. 2856–2865, 2009.
- [25] J. Zhou and Y. Li, "Existence of solutions for a class of second-order Hamiltonian systems with impulsive effects," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 3-4, pp. 1594–1603, 2010.
- [26] W. Ding and D. Qian, "Periodic solutions for sublinear systems via variational approach," Nonlinear Analysis: Real World Applications, vol. 11, no. 4, pp. 2603–2609, 2010.
- [27] J. Sun, H. Chen, and J. J. Nieto, "Infinitely many solutions for second-order Hamiltonian system with impulsive effects," *Mathematical and Computer Modelling*, vol. 54, no. 1-2, pp. 544–555, 2011.
- [28] J. Sun, H. Chen, and T. Zhou, "Multiplicity of solutions for a fourth-order impulsive differential equation via variational methods," *Bulletin of the Australian Mathematical Society*, vol. 82, no. 3, pp. 446–458, 2010.
- [29] D. Paşca and C.-L. Tang, "Some existence results on periodic solutions of nonautonomous secondorder differential systems with (q, p)-Laplacian," *Applied Mathematics Letters*, vol. 23, no. 3, pp. 246– 251, 2010.
- [30] D. Paşca, "Periodic solutions of a class of nonautonomous second order differential systems with (q, p)-Laplacian," Bulletin of the Belgian Mathematical Society, vol. 17, no. 5, pp. 841–850, 2010.
- [31] X. Zhang and X. Tang, "Periodic solutions for an ordinary p-Laplacian system," Taiwanese Journal of Mathematics, vol. 15, no. 3, pp. 1369–1396, 2011.
- [32] X. Tang and X. Zhang, "Periodic solutions for second-order Hamiltonian systems with a p-Laplacian," Annales Universitatis Mariae Curie-Skłodowska. Sectio A, vol. 64, no. 1, pp. 93–113, 2010.
- [33] Y. Tian and W. Ge, "Periodic solutions of non-autonomous second-order systems with a p-Laplacian," Nonlinear Analysis: Theory, Methods & Applications, vol. 66, no. 1, pp. 192–203, 2007.
- [34] H. Brezis and L. Nirenberg, "Remarks on finding critical points," Communications on Pure and Applied Mathematics, vol. 44, no. 8-9, pp. 939–963, 1991.
- [35] P. H. Rabinowitz, Minimax Methods in Critical Point Theory with Applications to Differential Equations, vol. 65 of CBMS-NSF Regional Conference Series in Applied Mathematics, Vol. 65, American Mathematical Society, Providence, RI, USA, 1986.



Advances in **Operations Research**



The Scientific World Journal







Hindawi

Submit your manuscripts at http://www.hindawi.com



Algebra



Journal of Probability and Statistics



International Journal of Differential Equations





Complex Analysis





Mathematical Problems in Engineering



Abstract and Applied Analysis



Discrete Dynamics in Nature and Society



International Journal of Mathematics and Mathematical Sciences





Journal of **Function Spaces**



International Journal of Stochastic Analysis

