

Research Article

Necessary and Sufficient Condition for Stability of Generalized Expectation Value

Aziz El Kaabouchi¹ and Sumiyoshi Abe^{1,2}

¹ Institut Supérieur des Matériaux et Mécaniques Avancés, 44 F. A. Bartholdi, 72000 Le Mans, France

² Department of Physical Engineering, Mie University, Mie 514-8507, Japan

Correspondence should be addressed to Sumiyoshi Abe, suabe@sf6.so-net.ne.jp

Received 2 April 2011; Revised 21 June 2011; Accepted 23 June 2011

Academic Editor: Bernard Geurts

Copyright © 2011 A. El Kaabouchi and S. Abe. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A class of generalized definitions of expectation value is often employed in nonequilibrium statistical mechanics for complex systems. Here, the necessary and sufficient condition is presented for such a class to be stable under small deformations of a given arbitrary probability distribution.

Given a probability distribution $\{p_i\}_{i=1,2,\dots,W}$, that is, $0 \leq p_i \leq 1$ ($i = 1, 2, \dots, W$) and $\sum_{i=1}^W p_i = 1$, the ordinary expectation value of a quantity Q of a system under consideration is defined by $\sum_{i=1}^W p_i Q_i$, where W is the total number of accessible states and is enormously large in statistical mechanics, typically being $2^{10^{23}}$. In the field of generalized statistical mechanics for complex systems, on the other hand, discussions are often made about altering this definition. Among others, the so-called “escort average” is widely employed in the field of generalized statistical mechanics [1–3]. It is defined as follows:

$$\langle Q \rangle_{\phi} [p] = \sum_{i=1}^W P_i^{(\phi)} Q_i, \quad (1)$$

where $P_i^{(\phi)}$ stands for the escort probability distribution [4] given by

$$P_i^{(\phi)} = \frac{\phi(p_i)}{\sum_{j=1}^W \phi(p_j)}, \quad (2)$$

with a nonnegative function ϕ . In the special case when $\phi(x) = x$, $\langle Q \rangle_{\phi}$ is reduced to the ordinary expectation value mentioned above.

Consider measurements of a certain quantity of a system to obtain information about the probability distribution. Repeated measurements should be performed on the system, which is identically prepared each time. Suppose that two probability distributions, $\{p_i\}_{i=1,2,\dots,W}$ and $\{p'_i\}_{i=1,2,\dots,W'}$, are obtained through the measurements. They may slightly be different from each other, in general. If such measurements make sense, then the expectation values, $\langle Q \rangle[p]$ and $\langle Q \rangle[p']$, calculated from these two distributions should also be close to each other. This condition, which implies "experimental robustness," is represented as follows.

Definition (stability). An expectation value $\langle Q \rangle[p]$ is said to be stable, if the following predicate holds for any pair of probability distributions, $\{p_i\}_{i=1,2,\dots,W}$ and $\{p'_i\}_{i=1,2,\dots,W'}$:

$$(\forall \varepsilon > 0) (\exists \delta > 0) (\forall W) \quad (\|p - p'\|_1 < \delta \implies |\langle Q \rangle[p] - \langle Q \rangle[p']| < \varepsilon). \quad (3)$$

Here, $\|p - p'\|_1 = \sum_{i=1}^W |p_i - p'_i|$ is the l^1 -norm describing the distance between these two probability distributions. One might consider norms of other kinds, but what is physically relevant to discrete systems is the present l^1 -norm [5]. Equation (3) is analogous to Lesche's stability condition on entropic functionals [5], which has recently been revisited in the literature [6–11] (note that the discussion in [8] is corrected in [9]). This concept of stability is actually equivalent to that of uniform continuity.

In recent papers [12, 13], it has been shown that the generalized expectation value in (1) with a specific class, $\phi(x) = x^q$ ($q > 0$), (the associated expectation value being termed the q -expectation value), is not stable unless $q = 1$. This result needs the q -expectation-value formalism of nonextensive statistical mechanics [1, 2] be reconsidered. In addition, the result is supported by Boltzmann-like kinetic theory in an independent manner [14].

Here, it seems appropriate to make some comments on the latest situation of the problems concerning stabilities of entropic functionals and generalized expectation values. The authors of [15, 16] have presented discussions which aim to rescue the q -expectation values from the difficulties of their instability pointed out in [12]. Those authors insist that the q -expectation values can be stable in both the finite- W and continuous cases. Such possibilities are, however, fully refuted by the work in [13] both physically and mathematically, and the controversy seems to have been terminated with that work. The case of the continuous variables has further been carefully examined in a recent paper [17], where the so-called Tsallis q -entropies [1, 2] do not have the continuous limit in consistency with the physical principles such as the thermodynamic laws (see also [18, 19]). These controversies have led the researchers to give up the traditional form of nonextensive statistical mechanics based on the q -entropies and q -expectation values and to examine other entropic functionals combined with the ordinary definition of expectation values [20] (see also [21, 22]). Thus, it seems that nonextensive statistical mechanics has to be fully reexamined, theoretically.

In this paper, we present the necessary and sufficient condition for $\langle Q \rangle_\phi[p]$ in (1) to be stable.

Our main result is as follows.

Theorem. Let ϕ be nonnegative and continuous on $[0, 1]$, differentiable on $(0, 1)$, and satisfy the condition that $\phi(x) = 0 \Leftrightarrow x = 0$. And, let $Q = \{Q_i\}_{i=1,2,\dots,W}$ be a random variable. Then, $\langle Q \rangle_\phi[p]$ in (1) is stable, if and only if $\lim_{x \rightarrow +0} \phi(x)/x \in (0, \infty)$.

Proof. First, assume that $\lim_{x \rightarrow +0} \phi(x)/x = a > 0$. Then, there exists $\delta_1 > 0$ such that

$$a - \frac{a}{2} < \frac{\phi(x)}{x} < a + \frac{a}{2} \quad (\forall x \in (0, \delta_1]). \quad (4)$$

$\phi(x)/x$ does not vanish because of the condition $\phi(x) = 0 \Leftrightarrow x = 0$. Therefore, there exists $b > 0$ such that

$$\frac{\phi(x)}{x} \geq b \quad (\forall x \in (\delta_1, 1]). \quad (5)$$

Putting $c = \min\{a/2, b\}$ we have

$$cx \leq \phi(x) \quad (\forall x \in [0, 1]). \quad (6)$$

Consequently, for an arbitrarily large W and an arbitrary probability distribution $\{p_i\}_{i=1,2,\dots,W}$, we obtain

$$\frac{1}{\sum_{i=1}^W \phi(p_i)} \leq c. \quad (7)$$

From the mean value theorem, it follows that

$$|\phi(p_i) - \phi(p'_i)| \leq |p_i - p'_i| \cdot \sup_{x \in (0,1)} |\phi'(x)|, \quad (8)$$

where $\phi'(x)$ is the derivative of $\phi(x)$ with respect to x . For $\varepsilon > 0$, we put

$$\delta = \inf \left(\delta_1, \frac{c\varepsilon}{2|Q_{\max}| \cdot \left(\sup_{x \in (0,1)} |\phi'(x)| \right)} \right), \quad (9)$$

where $Q_{\max} = \max \{Q_i\}_{i=1,2,\dots,W}$. Now, for $\|p - p'\|_1 < \delta$, we have

$$\begin{aligned} & \left| \langle Q \rangle_{\phi} [p] - \langle Q \rangle_{\phi} [p'] \right| \\ &= \frac{1}{\sum_{i=1}^W \phi(p_i) \sum_{j=1}^W \phi(p'_j)} \left| \sum_{i=1}^W Q_i \left\{ \phi(p_i) \sum_{j=1}^W \phi(p'_j) - \phi(p'_i) \sum_{j=1}^W \phi(p_j) \right\} \right| \\ &\leq \frac{1}{\sum_{i=1}^W \phi(p_i) \sum_{j=1}^W \phi(p'_j)} \\ &\quad \times \left[\sum_{i=1}^W |Q_i| \left\{ |\phi(p_i) - \phi(p'_i)| \sum_{j=1}^W \phi(p'_j) + \phi(p'_i) \left| \sum_{j=1}^W \phi(p_j) - \sum_{j=1}^W \phi(p'_j) \right| \right\} \right] \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{\sum_{j=1}^W \phi(p_j)} \sum_{i=1}^W |Q_i| |\phi(p_i) - \phi(p'_i)| \\
&\quad + \frac{\sum_{j=1}^W |\phi(p_j) - \phi(p'_j)|}{\sum_{i=1}^W \phi(p_i) \sum_{j=1}^W \phi(p'_j)} \sum_{i=1}^W |Q_i| \phi(p'_i) \\
&\leq \frac{2|Q_{\max}|}{\sum_{j=1}^W \phi(p_j)} \sum_{i=1}^W |\phi(p_i) - \phi(p'_i)| \\
&\leq \frac{2|Q_{\max}|}{\sum_{j=1}^W \phi(p_j)} \|p - p'\|_1 \cdot \sup_{x \in (0,1)} |\phi'(x)| \\
&\leq \frac{2|Q_{\max}|}{c} \|p - p'\|_1 \cdot \sup_{x \in (0,1)} |\phi'(x)| \\
&< \varepsilon.
\end{aligned} \tag{10}$$

Therefore, $\langle Q \rangle_\phi [p]$ is stable.

On the other hand, suppose that $\lim_{x \rightarrow +0} \phi(x)/x \notin (0, \infty)$. That is, (i) $\lim_{x \rightarrow +0} \phi(x)/x = 0$ or (ii) $\lim_{x \rightarrow +0} \phi(x)/x = \infty$. Below, we will examine these cases separately.

(i) Consider the following deformation:

$$\begin{aligned}
p_i &= \frac{1}{W-1} (1 - \delta_{i1}), \\
p'_i &= \left(1 - \frac{\delta}{2}\right) p_i + \frac{\delta}{2} \delta_{i1},
\end{aligned} \tag{11}$$

which are normalized and satisfy $\|p - p'\|_1 = \delta$. We have

$$\begin{aligned}
\sum_{i=1}^W \phi(p_i) &= (W-1) \phi\left(\frac{1}{W-1}\right), \\
\sum_{i=1}^W \phi(p'_i) &= \phi\left(\frac{\delta}{2}\right) + (W-1) \phi\left(\frac{1}{W-1} \left(1 - \frac{\delta}{2}\right)\right).
\end{aligned} \tag{12}$$

Difference of the expectation values is calculated as follows:

$$\begin{aligned}
&\langle Q \rangle_\phi [p] - \langle Q \rangle_\phi [p'] \\
&= -\frac{Q_1 \phi(\delta/2)}{\phi(\delta/2) + (W-1) \phi((1/(W-1))(1 - \delta/2))} \\
&\quad + \left(\sum_{i=2}^W Q_i \right) \left\{ \frac{1}{W-1} - \frac{\phi((1/(W-1))(1 - \delta/2))}{\phi(\delta/2) + (W-1) \phi((1/(W-1))(1 - \delta/2))} \right\}
\end{aligned}$$

$$\begin{aligned}
&= \frac{W}{W-1} (\bar{Q} - Q_1) \\
&\quad \times \frac{\phi(\delta/2)/(1-\delta/2)}{\phi(\delta/2)/(1-\delta/2) + \phi((1/(W-1))(1-\delta/2))/[(1/(W-1))(1-\delta/2)]} \\
&\xrightarrow{W \rightarrow \infty} \bar{Q} - Q_1,
\end{aligned} \tag{13}$$

since $\lim_{x \rightarrow +0} \phi(x)/x = 0$, where \bar{Q} is the arithmetic mean, $\bar{Q} = \sum_{i=1}^W Q_i/W$. Therefore, $\langle Q \rangle_\phi [p]$ is not stable.

(ii) Consider the following deformation:

$$\begin{aligned}
p_i &= \delta_{i1}, \\
p'_i &= \left(1 - \frac{\delta}{2} \frac{W}{W-1}\right) p_i + \frac{\delta}{2} \frac{1}{W-1},
\end{aligned} \tag{14}$$

which are also normalized and satisfy $\|p - p'\|_1 = \delta$. We have

$$\begin{aligned}
\sum_{i=1}^W \phi(p_i) &= \phi(1), \\
\sum_{i=1}^W \phi(p'_i) &= \phi\left(1 - \frac{\delta}{2}\right) + (W-1)\phi\left(\frac{\delta}{2} \frac{1}{W-1}\right).
\end{aligned} \tag{15}$$

Difference of the expectation values is calculated as follows:

$$\begin{aligned}
\langle Q \rangle_\phi [p] - \langle Q \rangle_\phi [p'] &= Q_1 \left\{ 1 - \frac{\phi(1-\delta/2)}{\phi(1-\delta/2) + (W-1)\phi((\delta/2)(1/(W-1)))} \right\} \\
&\quad - \left(\sum_{i=2}^W Q_i \right) \frac{\phi((\delta/2)(1/(W-1)))}{\phi(1-\delta/2) + (W-1)\phi((\delta/2)(1/(W-1)))} \\
&= \frac{W}{W-1} (Q_1 - \bar{Q}) \\
&\quad \times \frac{\phi((\delta/2)(1/(W-1)))/[(\delta/2)(1/(W-1))]}{\phi(1-\delta/2)/(\delta/2) + \phi((\delta/2)(1/(W-1)))/[(\delta/2)(1/(W-1))]} \\
&\xrightarrow{W \rightarrow \infty} Q_1 - \bar{Q},
\end{aligned} \tag{16}$$

since $\lim_{x \rightarrow +0} \phi(x)/x = \infty$. Therefore, $\langle Q \rangle_\phi [p]$ is not stable. \square

In the above proof, we have employed the specific deformations of the probability distributions as the counterexamples, which are considered in [5]. It is pointed out in [13] that these deformed distributions may experimentally be generated.

Finally, we mention a couple of simple stable examples.

Example 1.

$$\phi(x) = e^x - 1. \quad (17)$$

Example 2.

$$\phi(x) = \ln(1 + x^\alpha), \quad (18)$$

which yields a stable generalized expectation value, if and only if $\alpha = 1$.

On the other hand, as mentioned earlier, the q -expectation value is not stable, since $\phi(x) = x^q$ ($q > 0$, $q \neq 1$) does not satisfy the condition $\lim_{x \rightarrow +0} \phi(x)/x \in (0, \infty)$.

In conclusion, we have considered a class of generalized definitions of expectation value that are often employed in nonequilibrium statistical mechanics for complex systems, and have presented the necessary and sufficient condition for such a class to be stable under small deformations of a given arbitrary probability distribution.

Acknowledgment

The work of S. Abe was supported in part by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science.

References

- [1] S. Abe and Y. Okamoto, Eds., *Nonextensive Statistical Mechanics and Its Applications*, vol. 560, Springer, Berlin, Germany, 2001.
- [2] C. Tsallis, *Introduction to Nonextensive Statistical Mechanics: Approaching a Complex World*, Springer, New York, NY, USA, 2009.
- [3] C. Tsallis and A. M. C. Souza, "Constructing a statistical mechanics for Beck-Cohen superstatistics," *Physical Review E*, vol. 67, no. 2, Article ID 026106, pp. 1–5, 2003.
- [4] C. Beck and F. Schlögl, *Thermodynamics of Chaotic Systems*, vol. 4, Cambridge University Press, Cambridge, UK, 1993, An Introduction.
- [5] B. Lesche, "Instabilities of Rényi entropies," *Journal of Statistical Physics*, vol. 27, no. 2, pp. 419–422, 1982.
- [6] S. Abe, "Stability of Tsallis entropy and instabilities of Rényi and normalized Tsallis entropies: a basis for q -exponential distributions," *Physical Review E*, vol. 66, no. 4, Article ID 046134, 2002.
- [7] S. Abe, G. Kaniadakis, and A. M. Scarfone, "Stabilities of generalized entropies," *Journal of Physics. A. Mathematical and General*, vol. 37, no. 44, pp. 10513–10519, 2004.
- [8] E. M. F. Curado and F. D. Nobre, "On the stability of analytic entropic forms," *Physica A*, vol. 335, no. 1-2, pp. 94–106, 2004.
- [9] A. El Kaabouchi, C. J. Ou, J. C. Chen, G. Z. Su, and Q. A. Wang, "A counterexample against the Lesche stability of a generic entropy functional," *Journal of Mathematical Physics*, vol. 52, Article ID 063302, 2011.
- [10] J. Naudts, "Continuity of a class of entropies and relative entropies," *Reviews in Mathematical Physics*, vol. 16, no. 6, pp. 809–822, 2004.
- [11] A. E. Rastegin, "Continuity and stability of partial entropic sums," *Letters in Mathematical Physics*, vol. 94, no. 3, pp. 229–242, 2010.
- [12] S. Abe, "Instability of q -averages in nonextensive statistical mechanics," *Europhysics Letters*, vol. 84, no. 6, Article ID 60006, 2008.

- [13] S. Abe, "Anomalous behavior of q -averages in nonextensive statistical mechanics," *Journal of Statistical Mechanics: Theory and Experiment*, vol. 2009, no. 7, Article ID P07027, 2009.
- [14] S. Abe, "Generalized molecular chaos hypothesis and the H theorem: problem of constraints and amendment of nonextensive statistical mechanics," *Physical Review E*, vol. 79, no. 4, Article ID 041116, 2009.
- [15] R. Hanel and S. Thurner, "Stability criteria for q -expectation values," *Physics Letters. A*, vol. 373, no. 16, pp. 1415–1420, 2009.
- [16] R. Hanel, S. Thurner, and C. Tsallis, "On the robustness of q -expectation values and Rényi entropy," *Europhysics Letters*, vol. 85, no. 2, Article ID 20005, 2009.
- [17] S. Abe, "Essential discreteness in generalized thermostatics with non-logarithmic entropy," *Europhysics Letters*, vol. 90, no. 4, Article ID 50004, 2010.
- [18] B. Andresen, "Comment on "Essential discreteness in generalized thermostatics with non-logarithmic entropy" by Abe Sumiyoshi," *Europhysics Letters*, vol. 92, no. 4, Article ID 40005, 2010.
- [19] S. Abe, "Reply to the comment by B. Andresen," *Europhysics Letters*, vol. 92, no. 4, Article ID 40006, 2010.
- [20] J. F. Lutsko, J. P. Boon, and P. Grosfils, "Is the Tsallis entropy stable?" *Europhysics Letters*, vol. 86, no. 4, Article ID 40005, 2009.
- [21] J. P. Boon and J. F. Lutsko, "Nonextensive formalism and continuous Hamiltonian systems," *Physics Letters A*, vol. 375, no. 3, pp. 329–334, 2011.
- [22] J. F. Lutsko and J. P. Boon, "Questioning the validity of non-extensive thermodynamics for classical Hamiltonian systems," *Europhysics Letters*, vol. 95, no. 2, Article ID 20006, 2011.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

