

Research Article

On the Laplacian Coefficients and Laplacian-Like Energy of Unicyclic Graphs with n Vertices and m Pendant Vertices

Xinying Pai^{1,2} and Sanyang Liu¹

¹ Department of Mathematics, Xidian University, Shanxi Xi'an 710071, China

² College of Science, China University of Petroleum, Shandong Qingdao 266580, China

Correspondence should be addressed to Xinying Pai, paixinying@upc.edu.cn

Received 12 June 2012; Accepted 13 September 2012

Academic Editor: Alvaro Valencia

Copyright © 2012 X. Pai and S. Liu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let $\Phi(G, \lambda) = \det(\lambda I_n - L(G)) = \sum_{k=0}^n (-1)^k c_k(G) \lambda^{n-k}$ be the characteristic polynomial of the Laplacian matrix of a graph G of order n . In this paper, we give four transforms on graphs that decrease all Laplacian coefficients $c_k(G)$ and investigate a conjecture A. Ilić and M. Ilić (2009) about the Laplacian coefficients of unicyclic graphs with n vertices and m pendant vertices. Finally, we determine the graph with the smallest Laplacian-like energy among all the unicyclic graphs with n vertices and m pendant vertices.

1. Introduction

Let $G = (V, E)$ be a simple undirected graph with n vertices and $|E|$ edges and, let $L(G) = D(G) - A(G)$ be its Laplacian matrix. The Laplacian polynomial of G is the characteristic polynomial of its Laplacian matrix. That is

$$\Phi(G, \lambda) = \det(\lambda I_n - L(G)) = \sum_{k=0}^n (-1)^k c_k(G) \lambda^{n-k}. \quad (1.1)$$

The Laplacian matrix $L(G)$ has nonnegative eigenvalues $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1} \geq \mu_n = 0$ [1]. From Viète's formulas,

$$c_k(G) = \sigma_k(\mu_1, \mu_2, \dots, \mu_{n-1}) = \sum_{I \subseteq \{1, 2, \dots, n-1\}, |I|=k} \prod_{i \in I} \mu_i \quad (1.2)$$

is a symmetric polynomial of order $n - 1$. In particular, we have $c_0(G) = 1, c_1(G) = 2|E(G)|, c_n(G) = 0$ and $c_{n-1}(G) = n\tau(G)$, where $\tau(G)$ is the number of spanning trees of G . If G is a tree, coefficient $c_{n-2}(G)$ is equal to its Wiener index, which is a sum of distance between all pairs of vertices:

$$c_{n-2}(G) = W(G) = \sum_{u,v \in V} d(u,v). \quad (1.3)$$

The Wiener index is considered as one of the most used topological indices with high correlation with many physical and chemical properties of molecular compounds.

A unicyclic graph is a connected graph in which the number of vertices equals the number of edges. Recently, the study on the Laplacian coefficients attracts much attention.

Mohar [2] proved that among all trees of order n , the k th Laplacian coefficients $c_k(G)$ are largest when the tree is a path and are smallest for stars. Stevanović and Ilić [3] showed that among all connected unicyclic graphs of order n , the k th Laplacian coefficients $c_k(G)$ are largest when the graph is a cycle C_n and smallest when the graph is an S_n with an additional edge between two of its pendent vertices, where S_n is a star of order n . He and Shan [4] proved that among all bicyclic graphs of order n , the k th Laplacian coefficients $c_k(G)$ is smallest when the graph is obtained from C_4 by adding one edge connecting two non-adjacent vertices and adding $n - 4$ pendent vertices attached to the vertex of degree 3. A. Ilić and M. Ilić [5] verified that among trees on n vertices and m leaves, the balanced starlike tree $S(n, m)$ (see Definition 2.2) has minimal Laplacian coefficients. Some other works on Laplacian coefficients can be found in [6–8].

In this paper, we determine the smallest k th Laplacian coefficients $c_k(G)$ among all unicyclic graphs with n vertices and m pendent vertices. Thus we completely solve a conjecture on the minimal Laplacian coefficients of unicyclic graphs with n vertices and m pendent vertices (see [5]).

Motivated by the results in [3, 4, 9–12] concerning the minimal Laplacian coefficients and Laplacian-like energy of some graphs and the minimal molecular graph energy of unicyclic graphs with n vertices and m pendent vertices, this paper will characterize the unicyclic graphs with n vertices and m pendent vertices, which minimize Laplacian-like energy.

2. Transformations and Lemmas

In this section, we introduce some graphic transformations and lemmas, which can be used to prove our main results. The Laplacian coefficients $c_k(G)$ of a graph G can be expressed in terms of subtree structures of G by the following result of Kelmans and Chelnokov [13]. Let F be a spanning forest of G with components $T_i, i = 1, 2, \dots, k$ having n_i vertices each, and let $\gamma(F) = \prod_{i=1}^k n_i$.

Lemma 2.1 (see [13]). *The Laplacian coefficient $c_{n-k}(G)$ of a graph G is given by*

$$c_{n-k}(G) = \sum_{F \in \mathcal{F}_k} \gamma(F), \quad (2.1)$$

where \mathcal{F}_k is the set of all spanning forests of G with exactly k components.

For a real number x , we use $\lfloor x \rfloor$ to represent the largest integer not greater than x and $\lceil x \rceil$ to represent the smallest integer not less than x .

Definition 2.2 (see [5]). The balanced starlike tree $S(n, m)$, $3 \leq m \leq n - 1$, is a tree of order n with just one center vertex v , and each of the m branches of T at v is a path of length $\lfloor (n - 1)/m \rfloor$ or $\lceil (n - 1)/m \rceil$.

Let P_n be the path with n vertices. A path $P : vv_1v_2 \cdots v_k$ in G is called a pendent path if $d(v_1) = d(v_2) = \cdots = d(v_{k-1}) = 2$ and $d(v_k) = 1$. If $k = 1$, then we say vv_1 is a pendent edge of the graph G . A leaf or pendent vertex is a vertex of degree one. A branching vertex is a vertex of degree greater than two. The k paths $P_{l_1}, P_{l_2}, \dots, P_{l_k}$ are said to have almost equal lengths if l_1, l_2, \dots, l_k satisfy $|l_i - l_j| \leq 1$ for $1 \leq i, j \leq k$.

Definition 2.3 (see [5]). The dumbbell $D(n, a, b)$ consists of the path P_{n-a-b} together with a independent vertices adjacent to one leaf of P_{n-a-b} and b independent vertices adjacent to the other leaf.

The union $G = G_1 \cup G_2$ of graph G_1 and G_2 with disjoint vertex sets V_1 and V_2 and edge sets E_1 and E_2 is the graph $G = (V, E)$ with $V = V_1 \cup V_2$ and $E = E_1 \cup E_2$. If G is a union of two paths of lengths a and b , then G is disconnected and has $a + b$ vertices and $a + b - 2$ edges. Let $m_k(G)$ be the number of matchings of G containing exactly k independent edges. Especially, let $m_k(a, b)$ be the number of k matchings in $G = P_a \cup P_b$.

Lemma 2.4 (see [5]). *Let v be a vertex of nontrivial connected graph G , and let $G(p, q)$ denote the graph obtained from G by adding pendent paths $P = vv_1v_2 \cdots v_p$ and $Q = vu_1u_2 \cdots u_q$, at vertex v . Assume that both numbers p and q are even. If $p - 2 \geq q + 2 \geq 4$, then for every k we have*

$$m_k(G(p, q)) \leq m_k(G(p - 2, q + 2)). \quad (2.2)$$

Lemma 2.5 (see [12]). *Let $m_k(a, b)$ be the number of k -matchings in $G = P_a \cup P_b$ and $n = 4s + r$ with $0 \leq r \leq 3$. Then the following inequality holds:*

$$m_k(n, 0) \geq m_k(n - 2, 2) \geq m_k(n - 4, 4) \geq \cdots \geq m_k(2s + r, 2s). \quad (2.3)$$

Lemma 2.6 (see [5]). *Among trees on n vertices and $2 \leq m \leq n - 2$ leaves, the balanced starlike tree $S(n, m)$ has minimal Laplacian coefficient $c_k(G)$, for every $k = 0, 1, \dots, n$.*

Definition 2.7 (see [5]). Let v be a vertex of a tree T of degree $m + 1$. Suppose that P_1, P_2, \dots, P_m are pendent paths incident with v , with lengths $n_i \geq 1, i = 1, 2, \dots, m$. Let w be the neighbor of v distinct from the starting vertices of paths v_1, v_2, \dots, v_m , respectively. We form a tree $T' = \delta(T, v)$ by removing the edges $vv_1, vv_2, \dots, vv_{m-1}$ from T and adding $m - 1$ new edges $wv_1, wv_2, \dots, wv_{m-1}$ incident with w . We say that T' is a δ -transformation of T .

Lemma 2.8 (see [5]). *Let T be an arbitrary tree, rooted at the center vertex. Let vertex v be on the deepest level of tree T among all branching vertices with degree at least three. Then for the δ -transformation tree $T' = \delta(T, v)$ and $0 \leq k \leq n$ holds:*

$$c_k(T) \geq c_k(T'). \quad (2.4)$$




Figure 1: Example of π_1 -transformation.

Lemma 2.9 (see [14]). For every acyclic graph T with n vertices,

$$c_k(T) = m_k(S(T)), \quad 0 \leq k \leq n, \quad (2.5)$$

where $S(T)$ means the subdivision graph of T .

3. Main Results

In this section, we present four new graphic transformations that decrease the Laplacian coefficients.

Definition 3.1. Let u be a vertex in the cycle C of a unicyclic graph G , such that u has degree $p + 2$ and p pendent paths named P_1, P_2, \dots, P_p , where $P_i: u_{i,1}, u_{i,2}, \dots, u_{i,l_i}$, $1 \leq i \leq p$. If $l_i \geq l_j + 2$, and let

$$G_1 = G - u_{i,l_i-1}u_{i,l_i} + u_{j,l_j}u_{i,l_i} \triangleq \pi_1(G). \quad (3.1)$$

We say that G_1 is a π_1 -transformation of G .

It is easy to see that π_1 -transformation preserves the size of a cycle of G and the number of pendent vertices.

Theorem 3.2. Let G be a connected unicyclic graph with n vertices and m pendent vertices, $G_1 = \pi_1(G)$. Then for every $k = 0, 1, \dots, n$,


$$c_k(G) \geq c_k(G_1), \quad (3.2)$$

with equality if and only if $k \in \{0, 1, n - 1, n\}$.

Proof. It is easy to see that $c_0(G_1) = c_0(G) = 1$, $c_1(G_1) = 2|E(G_1)| = 2|E(G)| = c_1(G)$, $c_n(G_1) = c_n(G) = 0$, $c_{n-1}(G_1) = n\tau(G_1) = n|E(C)| = n\tau(G) = c_{n-1}(G)$.

Now, consider the coefficients c_{n-k} ($k \neq 0, 1, n - 1, n$). Let \mathcal{F}_k and \mathcal{F}_{k_1} be the sets of spanning forests of G and G_1 with exactly k components, respectively.

Without loss of generality, we assume that $l_1 \geq l_2 + 2$. Let $G_1 = \pi_1(G) = G - u_{1,l_1-1}u_{1,l_1} + u_{2,l_2}u_{1,l_1}$ (see Figure 1).

Figure 2: Example of π_2 -transformation.

Obviously, by the definition of the spanning forest, the cycle C in the unicyclic graph satisfies that $C \notin F \in \mathcal{F}_k$ and $C \notin F_1 \in \mathcal{F}_{k_1}$, where F and F_1 are the arbitrary forests in \mathcal{F}_k and \mathcal{F}_{k_1} , respectively. Without loss of generality, we remove one of the edges in the cycle C , say uv , so we get T and T' , respectively. By Lemmas 2.4 and 2.9, we have that for every $k = 0, 1, \dots, n$,

$$c_k(T) \geq c_k(T'), \quad (3.3)$$

with equality if and only if $k \in \{0, 1, n-1, n\}$. If we remove the other edge, say xy , we get S and S' , respectively. By Lemmas 2.4 and 2.9, we have that for every $k = 0, 1, \dots, n$,

$$c_k(S) \geq c_k(S'), \quad (3.4)$$

with equality if and only if $k \in \{0, 1, n-1, n\}$.

It is easy to see that $T - xy = S - uv$ and $T' - xy = S' - uv$. We know that the numbers of the same tree of spanning forests of $T - xy$ and $T' - xy$ with exactly k components are equal to the numbers of the same tree of spanning forests of $S - uv$ and $S' - uv$ with exactly k components, respectively.

Applying to Definition 3.1 and Lemma 2.1, we can show that for every $k = 0, 1, \dots, n$,

$$c_k(G) \geq c_k(G_1), \quad (3.5)$$

with equality if and only if $k \in \{0, 1, n-1, n\}$. □

Definition 3.3. Let v be a vertex in a cycle C of a connected unicyclic graph G , where $d(v) \geq 3$. Suppose that u is one of two neighbors adjacent to v in C , such that u has degree $p+2$ and p pendent paths incident with u and v has degree $q+2$ and q pendent paths incident with v . Let

$$G_2 = G - vv_{q+1} + uv_{q+1} \triangleq \pi_2(G), \quad (3.6)$$

where v_{q+1} is one of the other neighbors adjacent to v in C . We say that G_2 is a π_2 -transformation of G (see Figure 2).

Obviously, π_2 -transformation decreases the size of a cycle of G and preserves the number of pendent vertices.

Theorem 3.4. *Let G be a connected unicyclic graph with n vertices and m pendent vertices, $G_2 = \pi_2(G)$. Then for every $k = 0, 1, \dots, n$,*

$$c_k(G) \geq c_k(G_2), \quad (3.7)$$

with equality if and only if $k \in \{0, 1, n\}$.

Proof. Obviously, $c_0(G_2) = c_0(G) = 1$, $c_1(G_2) = 2|E(G_2)| = 2|E(G)| = c_1(G)$, $c_n(G_2) = c_n(G) = 0$. For $k = n - 1$, the length of a cycle in G is greater than the length of a cycle in G_2 . Therefore, $c_{n-1}(G) > c_{n-1}(G_2)$.

Now, consider the coefficients c_{n-k} ($k \neq 0, 1, n - 1, n$). Let \mathcal{F}_k and \mathcal{F}_{k_2} be the sets of spanning forests of G and G_2 with exactly k components, respectively. Let $F_2 \in \mathcal{F}_{k_2}$ and T' be the component of F_2 and $u \in V(T')$. If $v_{q+1} \in V(T')$, we define F with $V(F) = V(G)$ and

$$E(F) = E(F_2) - uv_{q+1} + vv_{q+1}. \quad (3.8)$$

Now, we distinguish F_2 as the following two cases.

Case 1 ($v \in V(T')$). We have trees of equal sizes in both spanning forests thus $\gamma(F) = \gamma(F_2)$.

Case 2 ($v \notin V(T')$). Let vertex v be in the tree S' , that is, $v \in V(S')$.

Note the fact that uv is a cut edge of G_2 . It is easy to see that F is a spanning forest of G , and the number of components of F is $k - 1$ or k . We claim that $F \in \mathcal{F}_k$. Otherwise, u, v belong to one tree of F ; then there exists a path P joining v_{q+1} to u in F ; then $uPv_{q+1}u$ is a cycle of F_2 , which contradicts the fact that F_2 is a forest.


Suppose that $T' - v_{q+1}$ contains $a \geq 1$ vertices in the cycle C (including u) and $b \geq 0$ vertices in the paths P_1, \dots, P_p , and $T' - u$ contains $c \geq 1$ vertices in the cycle C . Let S' contain $d \geq 1$ in the paths P_{p+1}, \dots, P_{p+q} . Assume the orders of the components of F_2 different from T' and S' are n_1, n_2, \dots, n_{k-2} . We have

$$\begin{aligned} \gamma(F) - \gamma(F_2) &= [(a+b)(c+d) - (a+b+c)d] \prod_{i=1}^{k-2} n_i \\ &= c(a+b-d) \prod_{i=1}^{k-2} n_i = c(a+b-d)N, \end{aligned} \quad (3.9)$$

where $N = \prod_{i=1}^{k-2} n_i$.

If we sum all differences for such forest, having fixed values a, c and $b+d = M$, we get

$$\begin{aligned} \sum_{F \in \mathcal{F}^*} \gamma(F) - \gamma(F_2) &= \sum_{F \in \mathcal{F}^*} c(a+b-d)N \\ &= cN \sum_{b=0}^{M-1} (a+2b-M) = (a-1)cNM. \end{aligned} \quad (3.10)$$

Figure 3: Example of π_3 -transformation.

It is easy to see that $a \geq 1$ and $c \geq 1$, so $(a-1)cNM \geq 0$. Since at least one vertex is in $C - u - v_{q+1}$, there exists one forest F_2 such that $a > 1$ and $c \geq 1$, and then $(a-1)cNM > 0$.

If $v_{q+1} \notin V(T')$, thus $\gamma(F) = \gamma(F_2)$.

Therefore, by using Lemma 2.1, we get

$$c_k(G) = \sum_{F \in \mathcal{F}_k} \gamma(F) > \sum_{F_2 \in \mathcal{F}_{k_2}} \gamma(F_2) = c_k(G_2). \quad (3.11)$$

This completes the proof of Theorem 3.4. \square

Definition 3.5. Let v (not in the cycle C) be a vertex of degree $q+1$ in a connected unicyclic graph G . Suppose that P_{p+1}, \dots, P_{p+q} are pendent paths incident with v . Let u be the neighbor of v distinct from the starting vertices of paths v_1, v_2, \dots, v_q , respectively. Let

$$G_3 = \pi_3(G) = G - vv_2 - vv_3 - \dots - vv_q + uv_2 + uv_3 + \dots + uv_q. \quad (3.12)$$

We say that G_3 is a π_3 -transformation of G (see Figure 3).

It is not difficult to see that π_3 -transformation preserves the size of a cycle of G and the number of pendent vertices.

Theorem 3.6. Let G be a connected unicyclic graph with n vertices and m pendent vertices, $G_3 = \pi_3(G)$. Then for every $k = 0, 1, \dots, n$,

$$c_k(G) \geq c_k(G_3), \quad (3.13)$$

with equality if and only if $k \in \{0, 1, n-1, n\}$.

Proof. Obviously, $c_0(G_3) = c_0(G) = 1$, $c_1(G_3) = 2|E(G_3)| = 2|E(G)| = c_1(G)$, $c_n(G_3) = c_n(G) = 0$, $c_{n-1}(G_3) = n\tau(G_3) = n|E(C)| = n\tau(G) = c_{n-1}(G)$.

Now, consider the coefficients c_{n-k} ($k \neq 0, 1, n-1, n$). Let \mathcal{F}_k and \mathcal{F}_{k_3} be the sets of spanning forests of G and G_3 with exactly k components, respectively. Obviously, by the definition of the spanning forest, the cycle C in the unicyclic graph satisfies that $C \notin F \in \mathcal{F}_k$ and $C \notin F_3 \in \mathcal{F}_{k_3}$, where F and F_3 are the arbitrary forests in \mathcal{F}_k and \mathcal{F}_{k_3} , respectively. Without loss of generality, we remove one of the edges on the cycle, say wu , so we get two trees T and




Figure 4: Example of π_4 -transformation.

T' , respectively. Applying to Definition 2.7, we know that $T' = \delta(T)$. Then using Lemma 2.8, we can get that for every $k = 0, 1, \dots, n$,

$$c_k(T) \geq c_k(T'), \quad (3.14)$$

with equality if and only if $k \in \{0, 1, n-1, n\}$. If we remove another edge, say xy , we get S and S' , respectively. By Definition 2.7, we know that $S' = \delta(S)$. Then applying to Lemma 2.8, we get that for every $k = 0, 1, \dots, n$,

$$c_k(S) \geq c_k(S'), \quad (3.15)$$

with equality if and only if $k \in \{0, 1, n-1, n\}$.

It is easy to see that $T - xy = S - uv$ and $T' - xy = S' - uv$. We know that the numbers of the same tree of spanning forests of $T - xy$ and $T' - xy$ with exactly k components are equal to the numbers of the same tree of spanning forests of $S - uv$ and $S' - uv$ with exactly k components, respectively.

By Definition 3.5 and Lemma 2.1, we have that for every $k = 0, 1, \dots, n$,

$$c_k(G) \geq c_k(G_3), \quad (3.16)$$

with equality if and only if $k \in \{0, 1, n-1, n\}$. □

Definition 3.7. Let u, v , and w be three vertices on the triangle in a unicyclic graph G . Suppose that P_1, \dots, P_p are pendent paths incident with u , P_{p+1}, \dots, P_{p+q} are pendent paths incident with v , and $P_{p+q+1}, \dots, P_{p+q+l}$ are pendent paths incident with w ($p+q+l = m$). Let

$$G_4 = G - vv_1 - \dots - vv_q + uv_1 + \dots + uv_q \triangleq \pi_4(G). \quad (3.17)$$

We say that G_4 is a π_4 -transformation of G (see Figure 4).

Theorem 3.8. Let u, v , and w be three vertices on the triangle in a unicyclic graph G , $G_4 = \pi_4(G)$. Then for every $k = 0, 1, \dots, n$,

$$c_k(G) \geq c_k(G_4), \quad (3.18)$$

with equality if and only if $k \in \{0, 1, n-1, n\}$.




Figure 5: Obtained trees from Figure 4.

Proof. It is obvious to see that $c_0(G_4) = c_0(G) = 1$, $c_1(G_4) = 2|E(G_4)| = 2|E(G)| = c_1(G)$, $c_n(G_4) = c_n(G) = 0$. For $k = n - 1$, the length of a cycle in G_4 is equal to the length of a cycle in G . Therefore, $c_{n-1}(G) = c_{n-1}(G_4)$.

Now, consider the coefficient c_{n-k} ($k \neq 0, 1, n - 1, n$). Let \mathcal{F}_k and \mathcal{F}_{k_4} be the sets of spanning forests of G and G_4 with exactly k components, respectively.

Similarly to the proof of Theorem 3.2, we can get 6 trees as shown in Figure 5. Obviously, by Definition 2.7, we know that $T'_i = \delta(T_i)$ ($i = 1, 2, 3$). And according to Lemma 2.8, we can verify that

$$\begin{aligned} c_k(T_1) &\geq c_k(T'_1), \\ c_k(T_2) &\geq c_k(T'_2), \\ c_k(T_3) &\geq c_k(T'_3). \end{aligned} \tag{3.19}$$


By (3.19), Definition 3.7, and Lemma 2.1, it is easy to see that for every $k = 0, 1, \dots, n$,

$$c_k(G) \geq c_k(G_4), \tag{3.20}$$

with equality if and only if $k \in \{0, 1, n - 1, n\}$. This completes the proof of Theorem 3.8. \square

Theorem 3.9. Let G be a connected unicyclic graph with n vertices and m pendent vertices. Then for $0 \leq k \leq n$,

$$c_k(G) \geq c_k(S'(n, m)), \tag{3.21}$$

Figure 6: $S'(n, m)$.

with equality if and only if $k \in \{0, 1, n\}$, where $S'(n, m)$ is as shown in Figure 6, and each of the m branches at v is a path of length $\lfloor (n-3)/m \rfloor$ or $\lceil (n-3)/m \rceil$.

Proof. Let $C = w_1w_2 \cdots w_t w_1$ be a cycle of connected unicyclic graph G , and let T_i be a tree attached at $w_i, i = 1, 2, \dots, t$. We can apply π_3 -transformation to T_i , such that the tree contains one branch vertex w_i with pendent path attached to it. Next, we can apply π_2 -transformation to decrease the size of the cycle C as long as the length of C is not 3. Then we can apply π_1 -transformation at the longest and the shortest path repeatedly, the Laplacian coefficients do not increase while the attached paths become more balanced. Finally, we can apply π_4 -transformation as long as it is not $S'(n, m)$.

According to Theorems 3.2, 3.4, 3.6, and 3.8, we know that π_i -transformation ($i = 1, 2, 3, 4$) cannot increase the Laplacian coefficients. So, for an arbitrary unicyclic graph G with n vertices and m pendent vertices, we verify that

$$c_k(G) \geq c_k(S'(n, m)), \quad (3.22)$$

where $0 \leq k \leq n$ and with equality if and only if $k = 0, 1, n$. This completes the proof of Theorem 3.9. \square

4. Laplacian-Like Energy of Unicyclic Graphs with m Pendent Vertices

Let G be a graph. The Laplacian-like energy of graph G , LEL for short, is defined as follows:

$$\text{LEL}(G) = \sum_{k=1}^{n-1} \sqrt{\mu_k}, \quad (4.1)$$

where $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_n = 0$ are the Laplacian eigenvalues of G . This concept was introduced by J. Liu and B. Liu [9], where it was demonstrated it has similar feature as molecular graph energy (for more details see [15]). Stevanović in [10] presented a connection between LEL and Laplacian coefficients.

Theorem 4.1. *Let G and H be two graphs with n vertices. If $c_k(G) \leq c_k(H)$ for $k = 1, 2, \dots, n-1$, then $\text{LEL}(G) \leq \text{LEL}(H)$. Furthermore, if a strict inequality $c_k(G) < c_k(H)$ holds for some $1 \leq k \leq n-1$, then $\text{LEL}(G) < \text{LEL}(H)$.*

Using this result, we can conclude the following.

Corollary 4.2. *Let G be a connected unicyclic graph with n vertices and m pendent vertices. Then if $G \not\cong S'(n, m)$*

$$\text{LEL}(S'(n, m)) < \text{LEL}(G), \quad (4.2)$$

where $S'(n, m)$ is shown in Figure 6, and each of the m branches at v is a path of length $\lfloor (n-3)/m \rfloor$ or $\lceil (n-3)/m \rceil$.

Acknowledgment

The authors would like to express their sincere gratitude to the anonymous referees whose constructive comments, valuable suggestions, and careful reading improved the final form of this paper.

References

- [1] D. M. Cvetković, M. Doob, and H. Sachs, *Spectra of Graphs-Theory and Applications*, Johann Ambrosius Barth, Heidelberg, Germany, 3rd edition, 1995.
- [2] B. Mohar, "On the Laplacian coefficients of acyclic graphs," *Linear Algebra and its Applications*, vol. 422, no. 2-3, pp. 736–741, 2007.
- [3] D. Stevanović and A. Ilić, "On the Laplacian coefficients of unicyclic graphs," *Linear Algebra and its Applications*, vol. 430, no. 8-9, pp. 2290–2300, 2009.
- [4] C.-X. He and H.-Y. Shan, "On the Laplacian coefficients of bicyclic graphs," *Discrete Mathematics*, vol. 310, no. 23, pp. 3404–3412, 2010.
- [5] A. Ilić and M. Ilić, "Laplacian coefficients of trees with given number of leaves or vertices of degree two," *Linear Algebra and its Applications*, vol. 431, no. 11, pp. 2195–2202, 2009.
- [6] A. Ilić, "On the ordering of trees by the Laplacian coefficients," *Linear Algebra and its Applications*, vol. 431, no. 11, pp. 2203–2212, 2009.
- [7] X.-D. Zhang, X.-P. Lv, and Y.-H. Chen, "Ordering trees by the Laplacian coefficients," *Linear Algebra and its Applications*, vol. 431, no. 12, pp. 2414–2424, 2009.
- [8] W. Lin and W. Yan, "Laplacian coefficients of trees with a given bipartition," *Linear Algebra and its Applications*, vol. 435, no. 1, pp. 152–162, 2011.
- [9] J. Liu and B. Liu, "A Laplacian-energy-like invariant of a graph," *MATCH. Communications in Mathematical and in Computer Chemistry*, vol. 59, no. 2, pp. 397–419, 2008.
- [10] D. Stevanović, "Laplacian-like energy of trees," *MATCH. Communications in Mathematical and in Computer Chemistry*, vol. 61, no. 2, pp. 407–417, 2009.
- [11] S.-W. Tan, "On the Laplacian coefficients of unicyclic graphs with prescribed matching number," *Discrete Mathematics*, vol. 311, no. 8-9, pp. 582–594, 2011.
- [12] A. Ilić, A. Ilić, and D. Stevanović, "On the Wiener index and Laplacian coefficients of graphs with given diameter or radius," *MATCH. Communications in Mathematical and in Computer Chemistry*, vol. 63, no. 1, pp. 91–100, 2010.
- [13] A. K. Kelmans and V. M. Chelnokov, "A certain polynomial of a graph and graphs with an extremal number of trees," *Journal of Combinatorial Theory B*, vol. 16, pp. 197–214, 1974.
- [14] B. Zhou and I. Gutman, "A connection between ordinary and Laplacian spectra of bipartite graphs," *Linear and Multilinear Algebra*, vol. 56, no. 3, pp. 305–310, 2008.
- [15] I. Gutman, "The energy of a graph," *Berichte der Mathematisch-Statistischen Sektion im Forschungszentrum Graz*, vol. 103, no. 100-105, pp. 1–22, 1978.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

