

Research Article

The Generalized Order- k Lucas Sequences in Finite Groups

Ömür Deveci^{1,2} and Erdal Karaduman^{1,2}

¹ Department of Mathematics, Faculty of Arts and Science, Kafkas University, 36100 Kars, Turkey

² Department of Mathematics, Faculty of Science, Atatürk University, 25240 Erzurum, Turkey

Correspondence should be addressed to Ömür Deveci, odeveci36@yahoo.com.tr

Received 18 October 2011; Accepted 11 December 2011

Academic Editor: Reinaldo Martinez Palhares

Copyright © 2012 Ö. Deveci and E. Karaduman. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the generalized order- k Lucas sequences modulo m . Also, we define the i th generalized order- k Lucas orbit $l_A^{i, \{\alpha_1, \alpha_2, \dots, \alpha_{k-1}\}}(G)$ with respect to the generating set A and the constants $\alpha_1, \alpha_2, \dots, \alpha_{k-1}$ for a finite group $G = \langle A \rangle$. Then, we obtain the lengths of the periods of the i th generalized order- k Lucas orbits of the binary polyhedral groups $\langle n, 2, 2 \rangle, \langle 2, n, 2 \rangle, \langle 2, 2, n \rangle$ and the polyhedral groups $(n, 2, 2), (2, n, 2), (2, 2, n)$ for $1 \leq i \leq k$.

1. Introduction

The well-known Fibonacci sequence $\{F_n\}$ is defined as

$$F_1 = F_2 = 1, \quad \text{for } n > 2, \quad F_n = F_{n-1} + F_{n-2}. \quad (1.1)$$

We call F_n the n th Fibonacci number. The Fibonacci sequence is

$$1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots \quad (1.2)$$

Definition 1.1. Let $f_n^{(k)}$ denote the n th member of the k -step Fibonacci sequence defined as

$$f_n^{(k)} = \sum_{j=1}^k f_{n-j}^{(k)} \quad \text{for } n > k \quad (1.3)$$

with boundary conditions $f_i^{(k)} = 0$ for $1 \leq i < k$ and $f_k^{(k)} = 1$. Reducing this sequence by modulus m , we can get a repeating sequence, which we denote by

$$f(k, m) = \left(f_1^{(k,m)}, f_2^{(k,m)}, \dots, f_n^{(k,m)} \dots \right), \quad (1.4)$$

where $f_i^{(k,m)} = f_i^{(k)} \pmod{m}$. We then have that $(f_1^{(k,m)}, f_2^{(k,m)}, \dots, f_k^{(k,m)}) = (0, 0, \dots, 0, 1)$ and it has the same recurrence relation as in (1.3) [1].

Theorem 1.2. $f(k, m)$ is a periodic sequence [1].

Let $h_k(m)$ denote the smallest period of $f(k, m)$, called the period of $f(k, m)$ or the Wall number of the k -step Fibonacci sequence modulo m . For more information see [1].

Definition 1.3. Let $h_{k(a_1, a_2, \dots, a_k)}(m)$ denote the smallest period of the integer-valued recurrence relation $u_n = u_{n-1} + u_{n-2} + \dots + u_{n-k}$, $u_1 = a_1$, $u_2 = a_2, \dots$, $u_k = a_k$ when each entry is reduced modulo m [2].

Lemma 1.4. For $a_1, a_2, \dots, a_k, x_1, x_2, \dots, x_k \in \mathbb{Z}$ with $m > 0$, a_1, a_2, \dots, a_k not all congruent to zero modulo m and x_1, x_2, \dots, x_k not all congruent to zero modulo m ,

$$h_{k(a_1, a_2, \dots, a_k)}(m) = h_{k(x_1, x_2, \dots, x_k)}(m), \quad (1.5)$$

see [2].

In [3], Taşçı and Kılıç defined the k sequences of the generalized order- k Lucas numbers as follows:

$$l_n^i = \sum_{j=1}^k l_{n-j}^i, \quad (1.6)$$

for $n > 0$ and $1 \leq i \leq k$, with boundary (initial) conditions

$$l_n^i = \begin{cases} 2 & \text{if } i = 2 - n, \\ -1 & \text{if } i = 1 - n, \\ 0 & \text{otherwise,} \end{cases} \quad (1.7)$$

for $1 - k \leq n \leq 0$, where l_n^i is the n th term of the i th sequence. When $i = 1$ and $k = 2$, the generalized order- k Lucas sequence reduces to the usual negative Fibonacci sequence, that is, $l_n^1 = -F_{n+1}$ for all $n \in \mathbb{Z}^+$.

In [3], it is obtained that

$$\begin{bmatrix} l_{n+1}^i \\ l_n^i \\ l_{n-1}^i \\ \vdots \\ l_{n-k+2}^i \end{bmatrix} = A \begin{bmatrix} l_n^i \\ l_{n-1}^i \\ l_{n-2}^i \\ \vdots \\ l_{n-k+1}^i \end{bmatrix}, \quad (1.8)$$

where

$$A = \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{bmatrix}. \quad (1.9)$$

The Lucas sequence, the generalized Lucas sequence, and their properties have been studied by several authors; see for example, [4–9]. The study of the Fibonacci sequences in groups began with the earlier work of Wall [10]. Knox examined the k -nacci (k -step Fibonacci) sequences in finite groups [11]. Karaduman and Aydin examined the periods of the 2-step general Fibonacci sequences in dihedral groups D_n [12]. Lü and Wang contributed to the study of the Wall number for the k -step Fibonacci sequence [1]. C. M. Campbell and P. P. Campbell examined the behaviour of the Fibonacci lengths of finite binary polyhedral groups [13]. Also, Devenci et al. obtained the periods of the k -nacci sequences in finite binary polyhedral groups [14]. Now, we extend the concept to k sequences of the generalized order- k Lucas numbers and we examine the periods of the i th generalized order- k Lucas orbits of the binary polyhedral groups $\langle n, 2, 2 \rangle$, $\langle 2, n, 2 \rangle$, $\langle 2, 2, n \rangle$ and the polyhedral groups $(n, 2, 2)$, $(2, n, 2)$, $(2, 2, n)$ for $1 \leq i \leq k$.

In this paper, the usual notation p is used for a prime number.

2. Main Results and Proofs

Reducing the k sequences of the generalized order- k Lucas numbers by modulus m , we can get a repeating sequence denoted by

$$l(i, m) = (\dots, l_1^{(i, m)}, l_2^{(i, m)}, \dots, l_n^{(i, m)}, \dots) \quad \text{for } n > 0, 1 \leq i \leq k, \quad (2.1)$$

where $l_n^{(i, m)} = l_n^i \pmod{m}$. It has the same recurrence relation as that in (1.6).

Let the notation $hl_k^i(m)$ denote the smallest period of $l(i, m)$. It is easy to see from Lemma 1.4 that $h_k(m) = hl_k^i(m)$.

For a given matrix $M = [b_{ij}]$ with b_{ij} 's being integers, $M \pmod{m}$ means that every entry of M is reduced modulo m , that is, $M \pmod{m} = (b_{ij} \pmod{m})$.

Let $\langle A \rangle_{p^\alpha} = \{A^i \pmod{p^\alpha} \mid i \geq 0\}$ be a cyclic group, and let $|\langle A \rangle_{p^\alpha}|$ denote the order of $\langle A \rangle_{p^\alpha}$. Then, we have the following.

Theorem 2.1. $h_k(p^\alpha) = |\langle A \rangle_{p^\alpha}|$.

Proof. It is clear that $|\langle A \rangle_{p^\alpha}|$ is divisible by $h_k(p^\alpha)$. Then we need only to prove that $h_k(p^\alpha)$ is divisible by $|\langle A \rangle_{p^\alpha}|$. Let $h_k(p^\alpha) = \lambda$. Then we have

$$A^\lambda = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1k} \\ a_{21} & a_{22} & \cdots & a_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kk} \end{bmatrix}. \quad (2.2)$$

By mathematical induction it is easy to prove that the elements of the matrix A^λ are in the following forms:

$$\begin{aligned} a_{11} &= f_{\lambda+k}^{(k)}, & a_{12} &= f_{\lambda+k-1}^{(k)} + \cdots + f_{\lambda+1}^{(k)}, & a_{13} &= f_{\lambda+k-1}^{(k)} + \cdots + f_{\lambda+2}^{(k)}, \dots, & a_{1k-1} &= f_{\lambda+k-1}^{(k)} + f_{\lambda+k-2}^{(k)}, & a_{1k} &= f_{\lambda+k-1}^{(k)}, \\ a_{21} &= f_{\lambda+k-1}^{(k)}, & a_{22} &= f_{\lambda+k-2}^{(k)} + \cdots + f_{\lambda}^{(k)}, & a_{23} &= f_{\lambda+k-2}^{(k)} + \cdots + f_{\lambda+1}^{(k)}, \dots, & a_{2k-1} &= f_{\lambda+k-2}^{(k)} + f_{\lambda+k-3}^{(k)}, & a_{2k} &= f_{\lambda+k-2}^{(k)}, \\ & \vdots & & \vdots & & \vdots & & \vdots & & \vdots \\ a_{k1} &= f_{\lambda+1}^{(k)}, & a_{k2} &= f_{\lambda}^{(k)} + \cdots + f_{\lambda-k+2}^{(k)}, & a_{k3} &= f_{\lambda}^{(k)} + \cdots + f_{\lambda-k+3}^{(k)}, \dots, & a_{kk-1} &= f_{\lambda}^{(k)} + f_{\lambda-1}^{(k)}, & a_{kk} &= f_{\lambda}^{(k)}. \end{aligned} \quad (2.3)$$

We thus obtain that

$$\begin{aligned} a_{ii} &\equiv 1 \pmod{p^\alpha} \quad \text{for } 1 \leq i \leq k, \\ a_{ij} &\equiv 0 \pmod{p^\alpha} \quad \text{for } 1 \leq i, j \leq k \text{ such that } i \neq j. \end{aligned} \quad (2.4)$$

So we get that $A^\lambda \equiv I \pmod{p^\alpha}$, which yields that n is divisible by $|\langle A \rangle_{p^\alpha}|$. We are done. \square

Definition 2.2. Let G be a finitely generated group $G = \langle A \rangle$, where $A = \{a_1, a_2, \dots, a_k\}$ and $1 \leq i \leq k$. The sequence

$$x_0 = (a_1)_{\alpha_1}, \quad x_1 = (a_2)_{\alpha_2}, \dots, \quad x_{k-2} = (a_{k-1})_{\alpha_{k-1}}, \quad (2.5)$$

where

$$(a_u)_{\alpha_u} = \begin{cases} a_u a_k^{i_u-k} & \text{if } \alpha_u = 1, \\ a_k^{i_u-k} a_u & \text{if } \alpha_u = 2 \end{cases} \quad (2.6)$$

such that $1 \leq u \leq k-1$ and $1 \leq \alpha_u \leq 2$, $x_{k-1} = a_k^{\alpha_k}$, $x_{k+\beta} = \prod_{j=1}^k x_{\beta+j-1}$ for $\beta \geq 0$, is called the i th generalized order- k Lucas orbit of G with respect to the generating set A and the $\alpha_1, \alpha_2, \dots, \alpha_{k-1}$ constants, denoted by $l_A^{i, \{\alpha_1, \alpha_2, \dots, \alpha_{k-1}\}}(G)$.

Example 2.3. The 3rd generalized order-4 Lucas orbits $l_{\{a_1, a_2, a_3, a_4\}}^{3, \{1, 1, 1\}}(G)$, $l_{\{a_1, a_2, a_3, a_4\}}^{3, \{1, 2, 1\}}(G)$, $l_{\{a_1, a_2, a_3, a_4\}}^{3, \{1, 1, 2\}}(G)$, $l_{\{a_1, a_2, a_3, a_4\}}^{3, \{1, 2, 2\}}(G)$, $l_{\{a_1, a_2, a_3, a_4\}}^{3, \{2, 1, 1\}}(G)$, $l_{\{a_1, a_2, a_3, a_4\}}^{3, \{2, 2, 1\}}(G)$, $l_{\{a_1, a_2, a_3, a_4\}}^{3, \{2, 1, 2\}}(G)$, and $l_{\{a_1, a_2, a_3, a_4\}}^{3, \{2, 2, 2\}}(G)$ of the finitely generated group $G = \langle A \rangle$, where $A = \{a_1, a_2, a_3, a_4\}$, respectively, are as follows:

$$\begin{aligned}
 x_0 &= a_1 a_4^{\alpha_3} = a_1, \quad x_1 = a_2 a_4^{\alpha_2} = a_2 a_4^{-1}, \quad x_2 = a_3 a_4^{\alpha_1} = a_3 a_4^2, \quad x_3 = a_4^{\alpha_0} = e, \quad x_{4+\beta} = \prod_{j=1}^4 x_{\beta+j-1} \\
 & \qquad \qquad \qquad (\beta \geq 0), \\
 x_0 &= a_1 a_4^{\alpha_3} = a_1, \quad x_1 = a_4^{\alpha_2} a_2 = a_4^{-1} a_2, \quad x_2 = a_3 a_4^{\alpha_1} = a_3 a_4^2, \quad x_3 = a_4^{\alpha_0} = e, \quad x_{4+\beta} = \prod_{j=1}^4 x_{\beta+j-1} \\
 & \qquad \qquad \qquad (\beta \geq 0), \\
 x_0 &= a_1 a_4^{\alpha_3} = a_1, \quad x_1 = a_2 a_4^{\alpha_2} = a_2 a_4^{-1}, \quad x_2 = a_4^{\alpha_1} a_3 = a_4^2 a_3, \quad x_3 = a_4^{\alpha_0} = e, \quad x_{4+\beta} = \prod_{j=1}^4 x_{\beta+j-1} \\
 & \qquad \qquad \qquad (\beta \geq 0), \\
 x_0 &= a_1 a_4^{\alpha_3} = a_1, \quad x_1 = a_4^{\alpha_2} a_2 = a_4^{-1} a_2, \quad x_2 = a_4^{\alpha_1} a_3 = a_4^2 a_3, \quad x_3 = a_4^{\alpha_0} = e, \quad x_{4+\beta} = \prod_{j=1}^4 x_{\beta+j-1} \\
 & \qquad \qquad \qquad (\beta \geq 0), \\
 x_0 &= a_4^{\alpha_3} a_1 = a_1, \quad x_1 = a_2 a_4^{\alpha_2} = a_2 a_4^{-1}, \quad x_2 = a_3 a_4^{\alpha_1} = a_3 a_4^2, \quad x_3 = a_4^{\alpha_0} = e, \quad x_{4+\beta} = \prod_{j=1}^4 x_{\beta+j-1} \\
 & \qquad \qquad \qquad (\beta \geq 0), \\
 x_0 &= a_4^{\alpha_3} a_1 = a_1, \quad x_1 = a_4^{\alpha_2} a_2 = a_4^{-1} a_2, \quad x_2 = a_3 a_4^{\alpha_1} = a_3 a_4^2, \quad x_3 = a_4^{\alpha_0} = e, \quad x_{4+\beta} = \prod_{j=1}^4 x_{\beta+j-1} \\
 & \qquad \qquad \qquad (\beta \geq 0), \\
 x_0 &= a_4^{\alpha_3} a_1 = a_1, \quad x_1 = a_2 a_4^{\alpha_2} = a_2 a_4^{-1}, \quad x_2 = a_4^{\alpha_1} a_3 = a_4^2 a_3, \quad x_3 = a_4^{\alpha_0} = e, \quad x_{4+\beta} = \prod_{j=1}^4 x_{\beta+j-1} \\
 & \qquad \qquad \qquad (\beta \geq 0), \\
 x_0 &= a_4^{\alpha_3} a_1 = a_1, \quad x_1 = a_4^{\alpha_2} a_2 = a_4^{-1} a_2, \quad x_2 = a_4^{\alpha_1} a_3 = a_4^2 a_3, \quad x_3 = a_4^{\alpha_0} = e, \quad x_{4+\beta} = \prod_{j=1}^4 x_{\beta+j-1} \\
 & \qquad \qquad \qquad (\beta \geq 0).
 \end{aligned} \tag{2.7}$$

It is well known that a sequence of group elements is periodic if, after a certain point, it consists only of repetitions of a fixed subsequence. The number of elements in the repeating subsequence is the period of the sequence.

Theorem 2.4. *The i th generalized order- k Lucas orbits in a finite group are periodic.*

Proof. The proof is similar to the proof of Theorem 1 in [10] and is omitted.

We denote the length of the period of the sequence $l_A^{i, \{\alpha_1, \alpha_2, \dots, \alpha_{k-1}\}}(G)$ by $\text{LEN}_A l^{i, \{\alpha_1, \alpha_2, \dots, \alpha_{k-1}\}}(G)$ and call it the i th generalized order- k Lucas length of G with respect to the generating set A and the constants $\alpha_1, \alpha_2, \dots, \alpha_{k-1}$.

From the definition it is clear that the i th generalized order- k Lucas length of a group depends on the chosen generating set and the order in which the assignments of x_0, x_1, \dots, x_{n-1} are made.

We will now address the i th generalized order- k Lucas lengths in specific classes of groups.

The binary polyhedral group $\langle l, m, n \rangle$, for $l, m, n > 1$, is defined by the presentation

$$\langle x, y, z \mid x^l = y^m = z^n = xyz \rangle \quad (2.8)$$

or

$$\langle x, y \mid x^l = y^m = (xy)^n \rangle. \quad (2.9)$$

The binary polyhedral group $\langle l, m, n \rangle$ is finite if, and only if, the number $k = lmn(1/l + 1/m + 1/n - 1) = mn + nl + lm - lmn$ is positive. Its order is $4lmn/k$.

For more information on these groups, see [15, pages 68–71].

The polyhedral group (l, m, n) , for $l, m, n > 1$, is defined by the presentation

$$\langle x, y, z \mid x^l = y^m = z^n = xyz = e \rangle \quad (2.10)$$

or

$$\langle x, y \mid x^l = y^m = (xy)^n = e \rangle. \quad (2.11)$$

The polyhedral group (l, m, n) is finite if, and only if, the number $k = lmn(1/l + 1/m + 1/n - 1) = mn + nl + lm - lmn$ is positive. Its order is $2lmn/k$.

For more information on these groups, see [15, pages 67–68]. \square

Theorem 2.5. *The i th generalized order-3 Lucas lengths of the binary polyhedral group $\langle n, 2, 2 \rangle$ for every i integer such that $1 \leq i \leq 3$ and the generating triple $\{x, y, z\}$ are as follows:*

- (i) $\text{LEN}_{\{x, y, z\}} l^{1, \{\alpha_1, \alpha_2\}}(\langle n, 2, 2 \rangle) = 8$ for $1 \leq \alpha_1, \alpha_2 \leq 2$,
- (ii) $\text{LEN}_{\{x, y, z\}} l^{2, \{\alpha_1, \alpha_2\}}(\langle n, 2, 2 \rangle) = h_3(2n)$ for $1 \leq \alpha_1, \alpha_2 \leq 2$,

- (iii) (1) $\text{LEN}_{\{x,y,z\}}^{3,\{1,1\}}(\langle n, 2, 2 \rangle) = \text{LEN}_{\{x,y,z\}}^{3,\{1,2\}}(\langle n, 2, 2 \rangle) = 8,$
 (2) $\text{LEN}_{\{x,y,z\}}^{3,\{2,1\}}(\langle n, 2, 2 \rangle) = \text{LEN}_{\{x,y,z\}}^{3,\{2,2\}}(\langle n, 2, 2 \rangle) = 4n$ if n is even, $8n$ if n is odd.

Proof. We prove the result by direct calculation. We first note that in the group defined by $\langle x, y, z \mid x^n = y^2 = z^2 = xyz, |x| = 2n$ (where $|x|$ means the order of x), $|y| = 4, |z| = 4, x = zy^3, y = x^{-1}z,$ and $z = xy$.

(i) The 1st generalized order-3 Lucas orbits of the group $\langle n, 2, 2 \rangle$ for generating triple $\{x, y, z\}$ and every constant α_1, α_2 such that $1 \leq \alpha_1, \alpha_2 \leq 2$ are the same and are as follows:

$$\begin{aligned} x_0 = x, x_1 = y, x_2 = z^{-1}, x_3 = e, x_4 = yz^{-1}, x_5 = xz^{-1}, x_6 = z^{-1}, \\ x_7 = x^n, x_8 = x, x_9 = y, x_{10} = z^{-1}, \dots \end{aligned} \quad (2.12)$$

Since the elements succeeding $x_8, x_9,$ and x_{10} depend on $x, y,$ and z^{-1} for their values, the cycle is again the 8th element; that is, $x_0 = x_8, x_1 = x_9, x_2 = x_{10}, \dots$. Thus, $\text{LEN}_{\{x,y,z\}}^{1,\{\alpha_1, \alpha_2\}}(\langle n, 2, 2 \rangle) = 8$ for $1 \leq \alpha_1, \alpha_2 \leq 2$.

(ii) Firstly, let us consider the orbits $l_{\{x,y,z\}}^{2,\{1,1\}}(\langle n, 2, 2 \rangle)$ and $l_{\{x,y,z\}}^{2,\{2,1\}}(\langle n, 2, 2 \rangle)$. The orbits $l_{\{x,y,z\}}^{2,\{1,1\}}(\langle n, 2, 2 \rangle)$ and $l_{\{x,y,z\}}^{2,\{2,1\}}(\langle n, 2, 2 \rangle)$ are the same and are as follows:

$$\begin{aligned} x_0 = x, x_1 = x^{-1}, x_2 = z^2, x_3 = z^2, x_4 = x^{-1}, x_5 = x^{-1}, x_6 = x^{-2}z^2, \\ x_7 = x^{-4}z^2, x_8 = x^{-7}, x_9 = x^{-13}, x_{10} = x^{-24}z^2, x_{11} = x^{-44}z^2, \dots \end{aligned} \quad (2.13)$$

For $m > 3$ we can see that the sequence will separate into some natural layers and each layer will be of the form

$$x_m = \begin{cases} x^{u_m} & \text{if } m \equiv 0 \pmod{4}, \\ x^{u_m} & \text{if } m \equiv 1 \pmod{4}, \\ x^{u_m} z^2 & \text{if } m \equiv 2 \pmod{4}, \\ x^{u_m} z^2 & \text{if } m \equiv 3 \pmod{4}, \end{cases} \quad (2.14)$$

where

$$u_m = u_{m-3} + u_{m-2} + u_{m-1}, \quad u_0 = 1, \quad u_1 = -1, \quad u_2 = 0. \quad (2.15)$$

Now the proof is finished when we note that the sequence will repeat when $x_{h_3(2n)} = x, x_{h_3(2n)+1} = x^{-1},$ and $x_{h_3(2n)+2} = z^2,$ where $h_3(2n)$ is the 3-step Wall number of the positive integer $2n$ and $h_3(2n) = 4\mu$ ($\mu \in N$). Letting $L = \text{LEN}_{\{x,y,z\}}^{2,\{1,1\}}(\langle n, 2, 2 \rangle) = \text{LEN}_{\{x,y,z\}}^{2,\{2,1\}}(\langle n, 2, 2 \rangle),$ we have

$$x_L = x^{u_L}, x_{L+1} = x^{u_{L+1}}, x_{L+2} = x^{u_{L+2}} z^2. \quad (2.16)$$

Using Lemma 1.4, we obtain $u_L = u_0 = 1$, $u_{L+1} = u_1 = -1$, and $u_{L+2} = u_2 = 0$. In this case the above equalities give

$$x_L = x^{u_L} = x, x_{L+1} = x^{u_{L+1}} = x^{-1}, x_{L+2} = x^{u_{L+2}} z^2 = x^0 z^2 = z^2. \quad (2.17)$$

The smallest nontrivial integer satisfying the above conditions occurs when the period is $h_3(2n)$.

Secondly, let us consider the orbits $l_{\{x,y,z\}}^{2,\{1,2\}}(\langle n, 2, 2 \rangle)$ and $l_{\{x,y,z\}}^{2,\{2,2\}}(\langle n, 2, 2 \rangle)$. The orbits $l_{\{x,y,z\}}^{2,\{1,2\}}(\langle n, 2, 2 \rangle)$ and $l_{\{x,y,z\}}^{2,\{2,2\}}(\langle n, 2, 2 \rangle)$ are the same and are as follows:

$$\begin{aligned} x_0 = x, x_1 = x, x_2 = z^2, x_3 = x^2 z^2, x_4 = x^3, x_5 = x^5, x_6 = x^{10} z^2, \\ x_7 = x^{18} z^2, x_8 = x^{33}, x_9 = x^{61}, x_{10} = x^{112} z^2, x_{11} = x^{206} z^2, \dots \end{aligned} \quad (2.18)$$

For $m > 3$ we can see that the sequence will separate into some natural layers and each layer will be of the form

$$x_m = \begin{cases} x^{v_m} & \text{if } m \equiv 0 \pmod{4}, \\ x^{v_m} & \text{if } m \equiv 1 \pmod{4}, \\ x^{v_m} z^2 & \text{if } m \equiv 2 \pmod{4}, \\ x^{v_m} z^2 & \text{if } m \equiv 3 \pmod{4}, \end{cases} \quad (2.19)$$

where

$$v_m = v_{m-3} + v_{m-2} + v_{m-1}, \quad v_0 = 1, \quad v_1 = 1, \quad v_2 = 0. \quad (2.20)$$

Now the proof is finished when we note that the sequence will repeat when $x_{h_3(2n)} = x$, $x_{h_3(2n)+1} = x$ and $x_{h_3(2n)+2} = z^2$. Letting $L = \text{LEN}_{\{x,y,z\}} l_{\{x,y,z\}}^{2,\{1,2\}}(\langle n, 2, 2 \rangle) = \text{LEN}_{\{x,y,z\}} l_{\{x,y,z\}}^{2,\{2,2\}}(\langle n, 2, 2 \rangle)$, we have

$$x_L = x^{v_L}, x_{L+1} = x^{v_{L+1}}, x_{L+2} = x^{v_{L+2}} z^2. \quad (2.21)$$

Using Lemma 1.4, we obtain $v_L = v_0 = 1$, $v_{L+1} = v_1 = 1$, and $v_{L+2} = v_2 = 0$. In this case the above equalities give

$$x_L = x^{v_L} = x, x_{L+1} = x^{v_{L+1}} = x, x_{L+2} = x^{v_{L+2}} z^2 = x^0 z^2 = z^2. \quad (2.22)$$

The smallest nontrivial integer satisfying the above conditions occurs when the period is $h_3(2n)$.

(iii) (1) The orbits $l_{\{x,y,z\}}^{3,\{1,1\}}(\langle n, 2, 2 \rangle)$ and $l_{\{x,y,z\}}^{3,\{1,2\}}(\langle n, 2, 2 \rangle)$ are the same and are as follows:

$$\begin{aligned} x_0 &= xz^{-1}, x_1 = y^3, x_2 = e, x_3 = x^{n+2}, x_4 = yx^2, \\ x_5 &= y^3, x_6 = x^n, x_7 = x^{n-2}, x_8 = xz^{-1}, x_9 = y^3, x_{10} = e, \dots \end{aligned} \quad (2.23)$$

So, we get $\text{LEN}_{\{x,y,z\}} l_{\{x,y,z\}}^{3,\{1,1\}}(\langle n, 2, 2 \rangle) = \text{LEN}_{\{x,y,z\}} l_{\{x,y,z\}}^{3,\{1,2\}}(\langle n, 2, 2 \rangle) = 8$.

(2) The orbits $l_{\{x,y,z\}}^{3,\{2,1\}}(\langle n, 2, 2 \rangle)$ and $l_{\{x,y,z\}}^{3,\{2,2\}}(\langle n, 2, 2 \rangle)$ are the same and are as follows:

$$\begin{aligned} x_0 &= y^3, x_1 = x^{n+1}, x_2 = e, x_3 = yx, x_4 = x^{-1}, x_5 = x^{-2}, x_6 = y^3 x^{-3}, \\ x_7 &= x^{-3} y x^{-3}, x_8 = y^3, x_9 = x^{n+1}, x_{10} = x^4, x_{11} = yx^5, x_{12} = y^3, \\ x_{13} &= x^{-1}, x_{14} = x^{-6}, x_{15} = y^3 x^{-7}, x_{16} = y^3, x_{17} = x^{n+1}, x_{18} = x^8, \dots \end{aligned} \quad (2.24)$$

The sequence can be said to form layers of length eight. Using the above, the sequence becomes

$$\begin{aligned} x_0 &= y^3, x_1 = x^{n+1}, x_2 = e, \dots, \\ x_8 &= y^3, x_9 = x^{n+1}, x_{10} = x^4, \dots, \\ x_{16} &= y^3, x_{17} = x^{n+1}, x_{18} = x^8, \dots, \\ x_{8i} &= y^3, x_{8i+1} = x^{n+1}, x_{8i+2} = x^{4i}, \dots \end{aligned} \quad (2.25)$$

So we need the smallest $i \in \mathbb{N}$ such that $4i = 2nk$ for $k \in \mathbb{N}$.

If n is even, then $i = n/2$. Thus, $8i = 4n$ and $\text{LEN}_{\{x,y,z\}} l_{\{x,y,z\}}^{3,\{2,1\}}(\langle n, 2, 2 \rangle) = \text{LEN}_{\{x,y,z\}} l_{\{x,y,z\}}^{3,\{2,2\}}(\langle n, 2, 2 \rangle) = 4n$.

If n is odd, then $i = n$. Thus, $8i = 8n$ and $\text{LEN}_{\{x,y,z\}} l_{\{x,y,z\}}^{3,\{2,1\}}(\langle n, 2, 2 \rangle) = \text{LEN}_{\{x,y,z\}} l_{\{x,y,z\}}^{3,\{2,2\}}(\langle n, 2, 2 \rangle) = 8n$. \square

Theorem 2.6. *The i th generalized order-2 Lucas lengths of the binary polyhedral group $\langle n, 2, 2 \rangle$ for every i such that $1 \leq i \leq 2$ and the generating pair $\{x, y\}$ are 6.*

Proof. We prove the result by direct calculation. We first note that in the group defined by

$$\langle x, y \mid x^n = y^2 = (xy)^2 \rangle, \quad |x| = 2n, \quad |y| = 4, \quad xy = yx^{-1}, \quad yx = x^{-1}y. \quad (2.26)$$

Firstly, let us consider the orbits $l_{\{x,y\}}^{1,\{1\}}(\langle n, 2, 2 \rangle)$ and $l_{\{x,y\}}^{1,\{2\}}(\langle n, 2, 2 \rangle)$. The orbits $l_{\{x,y\}}^{1,\{1\}}(\langle n, 2, 2 \rangle)$ and $l_{\{x,y\}}^{1,\{2\}}(\langle n, 2, 2 \rangle)$ are the same and are as follows:

$$x_0 = x, x_1 = y^{-1}, x_2 = xy^{-1}, x_3 = x^{n-1}, x_4 = x^2y, x_5 = y^{-1}x^{-1}, x_6 = x, x_7 = y^{-1}, \dots \quad (2.27)$$

So, we get $\text{LEN}_{\{x,y\}} l_{\{x,y\}}^{1,\{1\}}(\langle n, 2, 2 \rangle) = \text{LEN}_{\{x,y\}} l_{\{x,y\}}^{1,\{2\}}(\langle n, 2, 2 \rangle) = 6$.

Secondly, let us consider the orbit $l_{\{x,y\}}^{2,\{1\}}(\langle n, 2, 2 \rangle)$. The orbit $l_{\{x,y\}}^{2,\{1\}}(\langle n, 2, 2 \rangle)$ is as follows:

$$x_0 = xy^{-1}, x_1 = x^n, x_2 = xy, x_3 = xy^{-1}, x_4 = e, x_5 = xy^{-1}, x_6 = xy^{-1}, x_7 = x^n, \dots \quad (2.28)$$

So, we get $\text{LEN}_{\{x,y\}} l_{\{x,y\}}^{2,\{1\}}(\langle n, 2, 2 \rangle) = 6$.

Thirdly, let us consider the orbit $l_{\{x,y\}}^{2,\{2\}}(\langle n, 2, 2 \rangle)$. The orbit $l_{\{x,y\}}^{2,\{2\}}(\langle n, 2, 2 \rangle)$ is as follows:

$$x_0 = y^{-1}x, x_1 = x^n, x_2 = yx, x_3 = y^{-1}x, x_4 = e, x_5 = y^{-1}x, x_6 = y^{-1}x, x_7 = x^n, \dots \quad (2.29)$$

So, we get $\text{LEN}_{\{x,y\}} l_{\{x,y\}}^{2,\{2\}}(\langle n, 2, 2 \rangle) = 6$. □

Theorem 2.7. *The i th generalized order-3 Lucas lengths of the binary polyhedral group $\langle 2, n, 2 \rangle$ for every i integer such that $1 \leq i \leq 3$ and the generating triple $\{x, y, z\}$ are as follows:*

- (i) $\text{LEN}_{\{x,y,z\}} l^{1,\{\alpha_1, \alpha_2\}} \langle 2, n, 2 \rangle = 8$ for $1 \leq \alpha_1, \alpha_2 \leq 2$,
- (ii) $\text{LEN}_{\{x,y,z\}} l^{2,\{\alpha_1, \alpha_2\}} \langle 2, n, 2 \rangle = 8$ for $1 \leq \alpha_1, \alpha_2 \leq 2$,
- (iii) $\text{LEN}_{\{x,y,z\}} l^{3,\{\alpha_1, \alpha_2\}} \langle 2, n, 2 \rangle = h_3(2n)$ for $1 \leq \alpha_1, \alpha_2 \leq 2$.

Proof. The proof is similar to the proof of Theorem 2.5 and is omitted. □

Theorem 2.8. *The i th generalized order-2 Lucas lengths of the binary polyhedral group $\langle 2, n, 2 \rangle$ for every i such that $1 \leq i \leq 2$ and the generating pair $\{x, y\}$ are 6.*

Proof. The proof is similar to the proof of Theorem 2.6 and is omitted. □

Theorem 2.9. *The i th generalized order-3 Lucas lengths of the binary polyhedral group $\langle 2, 2, n \rangle$ for every i integer such that $1 \leq i \leq 3$ and the generating triple $\{x, y, z\}$ are as follows:*

(i)

$$\text{LEN}_{\{x,y,z\}} l^{1,\{\alpha_1, \alpha_2\}}(\langle 2, 2, n \rangle) = \begin{cases} 4n & \text{if } n \text{ is even,} \\ 8n & \text{if } n \text{ is odd} \end{cases} \quad \text{for } 1 \leq \alpha_1, \alpha_2 \leq 2, \quad (2.30)$$

(ii)

$$\text{LEN}_{\{x,y,z\}} l^{2,\{\alpha_1, \alpha_2\}}(\langle 2, 2, n \rangle) = \begin{cases} 2n & \text{if } n \equiv 0 \pmod{4}, \\ 4n & \text{if } n \equiv 2 \pmod{4}, \\ 8n & \text{otherwise} \end{cases} \quad \text{for } 1 \leq \alpha_1, \alpha_2 \leq 2, \quad (2.31)$$

(iii)

$$\text{LEN}_{\{x,y,z\}} l^{3,\{\alpha_1, \alpha_2\}}(\langle 2, 2, n \rangle) = 8 \quad \text{for } 1 \leq \alpha_1, \alpha_2 \leq 2 \quad (2.32)$$

Proof. We prove the result by direct calculation. We first note that in the group defined by $\langle x, y, z \mid x^2 = y^2 = z^n = xyz \rangle$, $|x| = 4$, $|y| = 4$, $|z| = 2n$, $x = yz$, $y = xz^{-1}$ and $z = yx^{-1}$.

(i) the 1st generalized order-3 Lucas orbits of the group $\langle 2, 2, n \rangle$ for generating triple $\{x, y, z\}$ and every constant α_1, α_2 such that $1 \leq \alpha_1, \alpha_2 \leq 2$ are the same and are as follows:

$$\begin{aligned} x_0 = x, x_1 = y, x_2 = z^{-1}, x_3 = yz^2y, x_4 = y^2z^3y, x_5 = y, x_6 = y^2z, \\ x_7 = y^2z^4, x_8 = xz^4, x_9 = y, x_{10} = z^{-1}, x_{11} = yz^6y, x_{12} = y^2z^7y, \\ x_{13} = y, x_{14} = y^2z, x_{15} = y^2z^8, x_{16} = xz^8, x_{17} = y, x_{18} = z^{-1}, \dots \end{aligned} \quad (2.33)$$

The sequence can be said to form layers of length eight. Using the above, the sequence becomes

$$\begin{aligned} x_0 = x, x_1 = y, x_2 = z^{-1}, \dots, \\ x_8 = xz^4, x_9 = y, x_{10} = z^{-1}, \dots, \\ x_{16} = xz^8, x_{17} = y, x_{18} = z^{-1}, \dots, \\ x_{8i} = xz^{4i}, x_{8i+1} = y, x_{8i+2} = z^{-1}, \dots \end{aligned} \quad (2.34)$$

So, we need the smallest $i \in \mathbb{N}$ such that $4i = 2nk$ for $k \in \mathbb{N}$.

If n is even, then $i = n/2$. Thus, $8i = 4n$ and $\text{LEN}_{\{x,y,z\}}^{l^{1,(\alpha_1,\alpha_2)}}(\langle 2, 2, n \rangle) = 4n$ for $1 \leq \alpha_1, \alpha_2 \leq 2$.

If n is odd, then $i = n$. Thus, $8i = 8n$ and $\text{LEN}_{\{x,y,z\}}^{l^{1,(\alpha_1,\alpha_2)}}(\langle 2, 2, n \rangle) = 8n$ for $1 \leq \alpha_1, \alpha_2 \leq 2$.

(ii) The orbits $l_{\{x,y,z\}}^{2,\{1,1\}}(\langle 2, 2, n \rangle)$ and $l_{\{x,y,z\}}^{2,\{2,1\}}(\langle 2, 2, n \rangle)$ are the same and are as follows:

$$\begin{aligned} x_0 = x, x_1 = yz^{-1}, x_2 = z^2, x_3 = z^n, x_4 = xz^n, x_5 = z^2x, x_6 = xz^2x, \\ x_7 = xz^4x, x_8 = z^8x, x_9 = yz^{-1}, x_{10} = z^2, x_{11} = z^{n+8}, x_{12} = xz^{n+8}, \\ x_{13} = z^2x, x_{14} = xz^2x, x_{15} = xz^{12}x, x_{16} = z^{16}x, x_{17} = yz^{-1}, x_{18} = z^2, \dots \end{aligned} \quad (2.35)$$

The sequence can be said to form layers of length eight. Using the above, the sequence becomes

$$\begin{aligned} x_0 = x, x_1 = yz^{-1}, x_2 = z^2, \dots, \\ x_8 = z^8x, x_9 = yz^{-1}, x_{10} = z^2, \dots, \\ x_{16} = z^{16}x, x_{17} = yz^{-1}, x_{18} = z^2, \dots, \\ x_{8i} = z^{8i}x, x_{8i+1} = yz^{-1}, x_{8i+2} = z^2, \dots \end{aligned} \quad (2.36)$$

So, we need the smallest $i \in \mathbb{N}$ such that $4i = 2nk$ for $k \in \mathbb{N}$.

If $n \equiv 0 \pmod{4}$, then $i = n/4$. Thus, $8i = 2n$ and $\text{LEN}_{\{x,y,z\}}^{l^{2,\{1,1\}}}(\langle 2, 2, n \rangle) = \text{LEN}_{\{x,y,z\}}^{l^{2,\{1,1\}}}(\langle 2, 2, n \rangle) = 2n$.

If $n \equiv 2 \pmod{4}$, then $i = n/2$. Thus, $8i = 4n$ and $\text{LEN}_{\{x,y,z\}}^{l^{2,\{1,1\}}}(\langle 2, 2, n \rangle) = \text{LEN}_{\{x,y,z\}}^{l^{2,\{1,1\}}}(\langle 2, 2, n \rangle) = 4n$.

If $n \equiv 1 \pmod{4}$ or $n \equiv 3 \pmod{4}$, then $i = n$. Thus, $8i = 8n$ and $\text{LEN}_{\{x,y,z\}}^{l^{2,\{1,1\}}}(\langle 2, 2, n \rangle) = \text{LEN}_{\{x,y,z\}}^{l^{2,\{1,1\}}}(\langle 2, 2, n \rangle) = 8n$.

The orbits $l_{\{x,y,z\}}^{2,\{1,1\}}(\langle 2, 2, n \rangle)$ and $l_{\{x,y,z\}}^{2,\{2,1\}}(\langle 2, 2, n \rangle)$ are the same. The proofs for these orbits are similar to the above and are omitted.

(iii) The orbits $l_{\{x,y,z\}}^{3,\{1,1\}}(\langle 2, 2, n \rangle)$, $l_{\{x,y,z\}}^{3,\{1,2\}}(\langle 2, 2, n \rangle)$, $l_{\{x,y,z\}}^{3,\{2,1\}}(\langle 2, 2, n \rangle)$, and $l_{\{x,y,z\}}^{3,\{2,2\}}(\langle 2, 2, n \rangle)$, respectively, are as follows:

$$\begin{aligned}
 x_0 &= y, x_1 = xz, x_2 = e, x_3 = z^{n+2}, x_4 = xz^{n+3}, x_5 = xz, \\
 x_6 &= z^n, x_7 = xz^2x, x_8 = y, x_9 = xz, x_{10} = e, \dots, \\
 x_0 &= y, x_1 = z^2y, x_2 = e, x_3 = xzy, x_4 = z^4y^3, x_5 = z^2y, \\
 x_6 &= z^n, x_7 = z^{n+2}, x_8 = y, x_9 = z^2y, x_{10} = e, \dots, \\
 x_0 &= xz, x_1 = xz, x_2 = e, x_3 = z^n, x_4 = xz^{n+1}, x_5 = xz, \\
 x_6 &= z^n, x_7 = z^n, x_8 = xz, x_9 = xz, x_{10} = e, \dots, \\
 x_0 &= xz, x_1 = z^2y, x_2 = e, x_3 = yz^4y, x_4 = z^{n+6}y, x_5 = z^2y, \\
 x_6 &= z^n, x_7 = z^{n+4}, x_8 = xz, x_9 = z^2y, x_{10} = e, \dots,
 \end{aligned} \tag{2.37}$$

which have period 8. □

Theorem 2.10. *The i th generalized order-2 Lucas lengths of the binary polyhedral group $\langle 2, 2, n \rangle$ for every i integer such that $1 \leq i \leq 2$ and the generating triple $\{x, y\}$ are as follows:*

- (i) $\text{LEN}_{\{x,y\}}^{l^{1,\{1\}}}(\langle 2, 2, n \rangle) = \text{LEN}_{\{x,y\}}^{l^{1,\{2\}}}(\langle 2, 2, n \rangle) = 6$,
- (ii) $\text{LEN}_{\{x,y\}}^{l^{2,\{1\}}}(\langle 2, 2, n \rangle) = \text{LEN}_{\{x,y\}}^{l^{2,\{2\}}}(\langle 2, 2, n \rangle) = h_2(2n)$.

Proof. We prove the result by direct calculation. We first note that in the group defined by $\langle x, y \mid x^2 = y^2 = (xy)^n \rangle$, $|x| = 4$, $|y| = 4$, and $|xy| = 2n$.

- (i) The orbits $l_{\{x,y\}}^{1,\{1\}}(\langle 2, 2, n \rangle)$ and $l_{\{x,y\}}^{1,\{2\}}(\langle 2, 2, n \rangle)$ are the same and are as follows:

$$x_0 = x, x_1 = y^3, x_2 = xy^3, x_3 = yxy, x_4 = y^3, x_5 = yx, x_6 = x, x_7 = y^3, \dots, \tag{2.38}$$

which have period 6.

- (ii) The orbits $l_{\{x,y\}}^{2,\{1\}}(\langle 2, 2, n \rangle)$ and $l_{\{x,y\}}^{2,\{2\}}(\langle 2, 2, n \rangle)$ are the same and are as follows:

$$x_0 = (xy)^{n-1}, x_1 = (xy)^n, \dots \tag{2.39}$$

We consider the recurrence relation defined by the following:

$$u_m = u_{m-2} + u_{m-1}, \quad u_0 = n - 1, \quad u_1 = n. \tag{2.40}$$

Then a routine induction shows that $x_m = (xy)^{u_m}$. Using Lemma 1.4, we obtain $u_L = u_0 = n-1$ and $u_{L+1} = u_1 = n$. In this case the equalities $x_m = (xy)^{u_m}$ give

$$x_L = (xy)^{u_L} = (xy)^{n-1}, x_{L+1} = (xy)^{u_{L+1}} = (xy)^n. \quad (2.41)$$

The smallest nontrivial integer satisfying the above conditions occurs when the period is $h_2(2n)$. \square

Theorem 2.11. *The i th generalized order-3 Lucas lengths of the polyhedral group $(n, 2, 2)$ for every i integer such that $1 \leq i \leq 3$ and the generating triple $\{x, y, z\}$ are as follows:*

- (i) $\text{LEN}_{\{x,y,z\}}^{l^{1, \{\alpha_1, \alpha_2\}}}((n, 2, 2)) = 6$ for $1 \leq \alpha_1, \alpha_2 \leq 2$,
- (ii) $\text{LEN}_{\{x,y,z\}}^{l^{2, \{\alpha_1, \alpha_2\}}}((n, 2, 2)) = h_3(n)$ for $1 \leq \alpha_1, \alpha_2 \leq 2$,
- (iii) (1) $\text{LEN}_{\{x,y,z\}}^{l^{3, \{1, 1\}}}((n, 2, 2)) = \text{LEN}_{\{x,y,z\}}^{l^{3, \{1, 2\}}}((n, 2, 2)) = 8$,
 (2) $\text{LEN}_{\{x,y,z\}}^{l^{3, \{2, 1\}}}((n, 2, 2)) = \text{LEN}_{\{x,y,z\}}^{l^{3, \{2, 2\}}}((n, 2, 2)) = 4$.

Proof. (i) We follow the proof given in [13].

The proofs of (ii) and (iii) are similar to the proofs of Theorem 2.5(ii) and 2.5(iii) and are omitted. \square

Theorem 2.12. *The i th generalized order-2 Lucas lengths of the polyhedral group $(n, 2, 2)$ for every i integer such that $1 \leq i \leq 2$ and the generating triple $\{x, y\}$ are as follows:*

- (i) $\text{LEN}_{\{x,y\}}^{l^{1, \{1\}}}((n, 2, 2)) = \text{LEN}_{\{x,y\}}^{l^{1, \{2\}}}((n, 2, 2)) = 6$,
- (ii) $\text{LEN}_{\{x,y\}}^{l^{2, \{1\}}}((n, 2, 2)) = \text{LEN}_{\{x,y\}}^{l^{2, \{2\}}}((n, 2, 2)) = 3$.

Proof. (i) The orbits $l^{1, \{1\}}((n, 2, 2))$ and $l^{1, \{2\}}((n, 2, 2))$ are the natural extension of the result of dihedral groups given in [16].

(ii) The orbits $l_{\{x,y\}}^{2, \{1\}}((n, 2, 2))$ and $l_{\{x,y\}}^{2, \{2\}}((n, 2, 2))$, respectively, are as follows:

$$\begin{aligned} x_0 = xy, x_1 = e, x_2 = xy, x_3 = xy, x_4 = e, \dots, \\ x_0 = yx, x_1 = e, x_2 = yx, x_3 = yx, x_4 = e, \dots, \end{aligned} \quad (2.42)$$

which have period 3. \square

Theorem 2.13. *The i th generalized order-3 Lucas lengths of the polyhedral group $(2, n, 2)$ for every i integer such that $1 \leq i \leq 3$ and the generating triple $\{x, y, z\}$ are as follows:*

- (i) $\text{LEN}_{\{x,y,z\}}^{l^{1, \{\alpha_1, \alpha_2\}}}((2, n, 2)) = 6$ for $1 \leq \alpha_1, \alpha_2 \leq 2$,
- (ii) (1) $\text{LEN}_{\{x,y,z\}}^{l^{2, \{1, 1\}}}((2, n, 2)) = \text{LEN}_{\{x,y,z\}}^{l^{2, \{2, 1\}}}((2, n, 2)) = 4$,
 (2) $\text{LEN}_{\{x,y,z\}}^{l^{2, \{1, 2\}}}((2, n, 2)) = \text{LEN}_{\{x,y,z\}}^{l^{2, \{2, 2\}}}((2, n, 2)) = 8$,
- (iii) $\text{LEN}_{\{x,y,z\}}^{l^{3, \{\alpha_1, \alpha_2\}}}((2, n, 2)) = h_3(n)$ for $1 \leq \alpha_1, \alpha_2 \leq 2$.

Proof. (i) We follow the proof given in [13].

The proofs of (ii) and (iii) are similar to the proofs of Theorem 2.5(ii) and 2.5(iii) and are omitted. \square

Theorem 2.14. *The i th generalized order-2 Lucas lengths of the polyhedral group $(2, n, 2)$ for every i such that $1 \leq i \leq 2$ and the generating pair $\{x, y\}$ are 6.*

Proof. The proof is similar to the proof of Theorem 2.6 and is omitted. \square

Theorem 2.15. *The i th generalized order-3 Lucas lengths of the polyhedral group $(2, 2, n)$ for every i integer such that $1 \leq i \leq 3$ and the generating triple $\{x, y, z\}$ are as follows:*

(i)

$$\text{LEN}_{\{x,y,z\}}^{l^{1,(\alpha_1,\alpha_2)}}(2, 2, n) = \begin{cases} 2n & \text{if } n \equiv 0 \pmod{4}, \\ 4n & \text{if } n \equiv 2 \pmod{4}, \text{ for } 1 \leq \alpha_1, \alpha_2 \leq 2 \\ 8n & \text{otherwise,} \end{cases} \quad (2.43)$$

(ii)

$$\text{LEN}_{\{x,y,z\}}^{l^{2,(\alpha_1,\alpha_2)}}((2, 2, n)) = \begin{cases} n & \text{if } n \equiv 0 \pmod{8}, \\ 2n & \text{if } n \equiv 4 \pmod{8}, \\ 4n & \text{if } n \equiv 2 \pmod{8}, \\ 8n & \text{otherwise,} \end{cases} \text{ for } 1 \leq \alpha_1, \alpha_2 \leq 2 \quad (2.44)$$

(iii) (1)

$$\begin{aligned} \text{LEN}_{\{x,y,z\}}^{l^{3,(1,1)}}((2, 2, n)) &= \text{LEN}_{\{x,y,z\}}^{l^{3,(1,2)}}((2, 2, n)) \\ &= \text{LEN}_{\{x,y,z\}}^{l^{3,(2,2)}}((2, 2, n)) = 8, \end{aligned} \quad (2.45)$$

(2)

$$\text{LEN}_{\{x,y,z\}}^{l^{3,(2,1)}}((2, 2, n)) = 4. \quad (2.46)$$

Proof. The proof is similar to the proof of Theorem 2.9 and is omitted. \square

Theorem 2.16. *The i th generalized order-2 Lucas lengths of the polyhedral group $(2, 2, n)$ for every i integer such that $1 \leq i \leq 2$ and the generating triple $\{x, y\}$ are as follows:*

$$(i) \text{LEN}_{\{x,y\}}^{l^{1,(1)}}((2, 2, n)) = \text{LEN}_{\{x,y\}}^{l^{1,(2)}}((2, 2, n)) = 6,$$

$$(ii) \text{LEN}_{\{x,y\}}^{l^{2,(1)}}((2, 2, n)) = \text{LEN}_{\{x,y\}}^{l^{2,(2)}}((2, 2, n)) = h_2(n).$$

Proof. (i) The orbits $l^{1,(1)}((2, 2, n))$ and $l^{1,(2)}((2, 2, n))$ are the natural extension of the result of dihedral groups given in [16].

(ii) The proof is similar to the proof of Theorem 2.10(ii) and is omitted. \square

Acknowledgment

This Project was supported by the Commission for the Scientific Research Projects of Kafkas University. The Project number is 2010-FEF-61.

References

- [1] K. Lü and J. Wang, “ k -step Fibonacci sequence modulo m ,” *Utilitas Mathematica*, vol. 71, pp. 169–177, 2006.
- [2] Ö. Deveci, E. Karaduman, and C. M. Campbell, “The periods of k -nacci sequences in centropolyhedral groups and related groups,” *Ars Combinatoria*, vol. 97, pp. 193–210, 2010.
- [3] D. Taşçı and E. Kılıç, “On the order- k generalized Lucas numbers,” *Applied Mathematics and Computation*, vol. 155, no. 3, pp. 637–641, 2004.
- [4] A. Akbary and Q. Wang, “A generalized Lucas sequence and permutation binomials,” *Proceedings of the American Mathematical Society*, vol. 134, no. 1, pp. 15–22, 2006.
- [5] H. Doostie and K. Ahmadidelir, “A class of Z -metacyclic groups involving the Lucas numbers,” *Novi Sad Journal of Mathematics*, vol. 39, no. 1, pp. 21–29, 2009.
- [6] E. Kılıç and D. Taşçı, “On the generalized order- k Fibonacci and Lucas numbers,” *The Rocky Mountain Journal of Mathematics*, vol. 36, no. 6, pp. 1915–1926, 2006.
- [7] G.-Y. Lee, “ k -Lucas numbers and associated bipartite graphs,” *Linear Algebra and its Applications*, vol. 320, no. 1–3, pp. 51–61, 2000.
- [8] J. B. Muskat, “Generalized Fibonacci and Lucas sequences and rootfinding methods,” *Mathematics of Computation*, vol. 61, no. 203, pp. 365–372, 1993.
- [9] N. Y. Özgür, “On the sequences related to Fibonacci and Lucas numbers,” *Journal of the Korean Mathematical Society*, vol. 42, no. 1, pp. 135–151, 2005.
- [10] D. D. Wall, “Fibonacci series modulo m ,” *The American Mathematical Monthly*, vol. 67, pp. 525–532, 1960.
- [11] S. W. Knox, “Fibonacci sequences in finite groups,” *The Fibonacci Quarterly*, vol. 30, no. 2, pp. 116–120, 1992.
- [12] E. Karaduman and H. Aydın, “On the periods of 2-step general Fibonacci sequences in dihedral groups,” *Matematički Vesnik*, vol. 58, no. 1-2, pp. 47–56, 2006.
- [13] C. M. Campbell and P. P. Campbell, “The Fibonacci lengths of binary polyhedral groups and related groups,” *Congressus Numerantium*, vol. 194, pp. 95–102, 2009.
- [14] Ö. Deveci, E. Karaduman, C. M. Campbell, and J. Zhang, “On the k -nacci sequences in finite binary polyhedral groups,” *Algebra Colloquium*, vol. 18, spec. 1, pp. 945–954, 2011.
- [15] H. S. M. Coxeter and W. O. J. Moser, *Generators and Relations for Discrete Groups*, Springer, New York, NY, USA, 3rd edition, 1972.
- [16] C. M. Campbell, H. Doostie, and E. F. Robertson, “Fibonacci length of generating pairs in groups,” in *Applications of Fibonacci Numbers*, G. E. Bergum et al., Ed., vol. 3, pp. 27–35, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1990.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

