



One-Pot Synthesis and Antimicrobial Activity of Novel α -Aminophosphonates Using TMG

B. SIVA KUMAR*, Y. HARANADHA REDDY,
C. RADHA RANI and G. CHNADRA SEKHAR REDDY

*Department of Chemistry
Vel Tech Engineering College, Avadi, Chennai-600 062, India
Department of Chemistry
Sri Venakateswara University, Tirupati-517 502, India
bskumar_badur@yahoo.com

Received 2 March 2011; Accepted 28 April 2011

Abstract: α -Aminophosphonates (**4a-j**) were synthesized in one-pot simultaneous reaction of 4-bromo-3-methyl benzenamine (**1**), dimethylphosphite (**3**) and different aromatic aldehydes (**2a-j**) by Kabachnik-Fields reaction in the presence of tetramethylguanidine (TMG) (10 mole%) as catalyst in toluene at reflux temperature afforded **4a-j** in good yields. All these compounds were found to exhibit moderate to good antimicrobial activity.

Keywords: α -Aminophosphonate, 4-Bromo-3-methylbenzenamine, Tetramethyl-guanidine, Antimicrobial activity

Introduction



Synthesis of α -aminophosphonates exhibiting high bio-activity has recently attracted a lot of attention¹⁻³. Their diverse applications include inhibition of synthase⁴, HIV protease⁵, renin⁶, and PTPases^{7,8}. Some of these derivatives are potential antibiotics⁹ and herbicides¹⁰. α -Aminophosphonates are chief substrates in the synthesis of phosphonopeptides¹¹. Due to their structural analogy with α -amino acids, these types of organophosphorus compounds are widely used for the development of new inhibitors of enzymes, neuroactive compounds and plant growth regulators^{12,13}. Among the number of synthetic approaches to α -aminophosphonates, one of the most powerful methods is the Kabachnik-Fields reaction^{14,15}. Previous results demonstrated that tetramethylguanidine (TMG) catalyzes the Michael addition of nitromethane to α,β -unsaturated ketones^{16,17}. TMG has been used only sporadically and has not yet received full recognition as a strong base in organic synthesis. Its catalytic activity in the Kabachnik-Fields reaction is explored in the present investigation.

Experimental

IR spectra were recorded in KBr pellets on a Perkin-Elmer 683 spectrophotometer. ^1H NMR spectra were recorded at 300 MHz in CDCl_3 using TMS as internal standard reference. ^{31}P NMR (121.4 MHz) was taken in CDCl_3 using 85% H_3PO_4 as external standard with broadband ^1H decoupling. ^{13}C NMR spectra measurements were performed at 75.4 MHz using TMS as internal standard. ^1H , ^{13}C , ^{31}P NMR spectra were taken on Varian Gemini 300 MHz spectrometer and mass spectra were recorded on JEOL MSD- instrument.

General procedure for the synthesis of α -aminophosphonates (**4a-j**)

To a stirred solution of 4-bromo-3-methylbenzenamine (**1**) (0.930 g, 0.005 mole) the aldehyde (**2a-j**) (0.005 mole) in anhydrous toluene (20 mL) was added drop wise and then TMG (10 mole %) was added and stirring continued at RT for 1h. Then dimethylphosphite (**3**) (0.55 g, 0.005 mole) in anhydrous toluene (20 mL) was added drop wise. Stirring was continued at RT for another 0.5 h and then the mixture was heated at gentle reflux for 5-6 h. The progress of the reaction was monitored by TLC analysis. After completion of the reaction the solvent was removed under reduced pressure. The residue was purified by column chromatography on silica gel (80-120 mesh) using hexane-ethyl acetate (8:2) as eluent.



Scheme 1. Physical, analytical and spectral data for compounds **4a-j**

Dimethyl (4-bromo-3-methylphenylamino) (4-hydroxyphenyl) methylphosphonate (4a)

Yield (68%), mp 108-109 °C; IR (KBr) cm^{-1} 3255 (P-N-H), 1233 (P=O), 762 (P-CH); ^1H -NMR ($\text{DMSO}-d_6$): δ 6.12-7.09 (m, 7H, Ar-H), 4.98 (s, 1H, NH), 4.62-4.69 (m, 1H, PCH), 3.37-3.42 (d, $^3J_{\text{P-H}} = 10.2$ Hz, P-OCH₃), 3.52-3.64 (d, $^3J_{\text{P-H}} = 10.9$ Hz, P-OCH₃), 2.26 (s, 3H, Ar-CH₃); ^{13}C -NMR($\text{DMSO}-d_6$), δ 146.5 (C-1), 112.7 (C-2), 132.5 (C-3), 112.4 (C-5), 139.1

(C-5), 115.7 (C-6), 128.6 (C-1'), 136.5 (2C, C-2'&6'), 115.5 (2C, C-3'&5'), 156.9 (C-4'), 54.90(d, P-CH, $J = 7.6$ Hz), 53.50 (d, $J = 6.3$ Hz, P-OCH₃); ³¹P-NMR (DMSO-*d*₆): δ 26.28; GCMS *m/z* (%): 401(M+2), 399(M+), 385.0 (11), 277.2 (100), 231.1 (15), 185.0 (36), 110.1 (4); Anal. Calcd. for C₁₆H₁₉BrNO₄P: C, 48.02; H, 4.79; N, 3.50. Found: C, 47.91; H, 4.73; N, 3.47%

Dimethyl (4-bromo-3-methylphenylamino) (2-hydroxyphenyl) methylphosphonate (4b)

Yield (73%) mp 109-110 °C; IR (KBr) cm⁻¹; 3375 (P-N-H), 1216 (P=O), 756 (P-CH); ¹H-NMR (DMSO-*d*₆) δ 6.41-7.32 (m, 7H, Ar-H), 4.97 (s, 1H, NH), 4.68-4.80 (m, 1H, P-CH), 3.42-3.51 (d, ³*J*_{P-H} = 10.1 Hz, P-OCH₃), 3.63-3.72 (d, ³*J*_{P-H} = 10.7 Hz, P-OCH₃) 2.31 (s, 3H, Ar-CH₃); ¹³C-NMR (DMSO-*d*₆), δ 145.9 (C-1), 113.1 (C-2), 133.0 (C-3), 113.2 (C-4), 140.1 (C-5), 115.9 (C-6), 121.6 (C-1'), 154.5 (C-2'), 115.7 (C-3'), 128.9 (C-4') 121.7 (C-5'), 128.5 (C-6') 55.60 (d, $J = 7.5$ Hz, P-CH), 53.91 (d, $J = 6.5$ Hz, P-OCH₃); ³¹P-NMR (DMSO-*d*₆): δ 29.37; Anal. Calcd. for C₁₆H₁₉BrNO₄P: C, 48.02; H, 4.79; N, 3.50. Found: C, 47.89 H, 4.73 ; N, 3.47%

Dimethyl (4-bromo-3-methylphenylamino) (3,4-dihydroxyphenyl) methyl phosphonate (4c)

Yield (71%), mp 105-106 °C; IR (KBr) cm⁻¹; 3345 (P-NH), 1213 (P=O), 775 (P-CH); ¹H-NMR (DMSO-*d*₆) δ 6.56-7.51 (m, 6H, Ar-H), 4.87 (s, 1H, NH), 4.71-4.82 (m, 1H, P-CH), 3.38-3.47 (d, ³*J*_{P-H} = 10.1 Hz, P-OCH₃), 3.66-3.74 (d, ³*J*_{P-H} = 10.9 Hz, P-OCH₃), 2.35 (s, 3H, Ar-CH₃); ¹³C-NMR(DMSO-*d*₆), δ 146.9 (C-1), 114.9 (C-2), 132.7 (C-3), 113.2 (C-4), 139.1 (C-5), 115.3 (C-6), 130.8 (C-1'), 114.3 (C-2'), 147.7 (C-3'), 145.9 (C-4') 117.5 (C-5'), 122.1 (C-6'), 55.81 (d, P-CH, $J = 8.1$ Hz), 53.27 (d, $J = 6.5$ Hz, P-OCH₃); ³¹P NMR (DMSO-*d*₆); δ 25.58. Anal. Calcd. for C₁₆H₁₉BrNO₅P: C, 44.80; H, 4.26; N, 3.48. Found: C, 44.68; H, 4.21; N, 3.43%.

Dimethyl (4-bromo-3-methylphenylamino) (3-hydroxy-4-methoxyphenyl) methyl phosphonate (4d)

Yield (68%), mp 97-98 °C; IR (KBr) cm⁻¹; 3390 (P-NH), 1232 (P=O), 742 (P-CH); ¹H NMR (DMSO-*d*₆) δ 6.62-7.37 (m, 6H, Ar-H), 4.86 (s, 1H, NH), 4.66-4.78 (m, 1H, PCH), 3.81 (s, 3H, Ar-OCH₃), 3.23-3.28 (d, ³*J*_{P-H} = 10.0 Hz, P-OCH₃), 3.57-3.65 (d, ³*J*_{P-H} = 11.1 Hz, P-OCH₃) 2.28 (s, 3H, Ar-CH₃); ¹³C NMR (DMSO-*d*₆) δ 122.7 (C-2), 112.5 (C-3), 121.8 (C-4), 120.6 (C-5), 119.6 (C-6), 111.5 (C-7), 131.2 (C-8), 179 (C-9), 151.5 (C-3'), 113.5 (C-4'), 160.9 (C-5'), 55.83 (d, $J = 7.9$ Hz, P-CH), 54.0 (d, $J = 6.4$ Hz, P-OCH₃), 15.8 (Ar-CH₃); ³¹P NMR (DMSO-*d*₆); δ 31.29; Anal. Calcd. for C₁₇H₂₁BrNO₅P : C, 47.46; H, 4.92; N, 3.26. Found: C, 47.34; H, 4.85; N, 3.19%.

Dimethyl (4-bromo-3-methylphenylamino) (4-methoxyphenyl) methyl phosphonate (4e)

Yield (74%), mp 110-111 °C; IR (KBr) cm⁻¹; 3225 (P-NH), 1228 (P=O), 750 (P-CH); ¹H-NMR (DMSO-*d*₆) δ 6.72-7.54 (m, 7H, Ar-H), 5.08 (s, 1H, NH), 4.75-4.93 (m, 1H, PCH), 3.87 (s, 3H, Ar-OCH₃), 3.22-3.29 (d, ³*J*_{P-H} = 10.0 Hz, P-OCH₃), 3.61-3.68 (d, ³*J*_{P-H} = 11.2 Hz, P-OCH₃), 2.15(s, 3H, Ar-CH₃); ³¹P NMR (DMSO-*d*₆); δ 28.62. Anal. Calcd. for C₁₇H₂₁BrNO₄P: C, 49.29; H, 5.11; N, 3.38. Found: C, 49.17; H, 5.07; N, 3.32%.

Dimethyl (4-bromo-3-methylphenylamino) (3,4-dimethoxyphenyl) methyl phosphonate (4f)

Yield (69%), mp 123-124 °C; IR (KBr) cm⁻¹; 3385 (P-NH), 1235 (P=O), 762 (P-CH); ¹H-NMR (DMSO-*d*₆) δ 6.60-7.64 (m, 6H, Ar-H), 4.97 (s, 1H, NH), 4.85-4.89 (m, 1H, P-CH), 3.87-3.98 (m 6H, Ar-OCH₃), 3.29-3.37 (d, ³*J*_{P-H} = 10.0 Hz, P-OCH₃), 3.59-3.68 (d, ³*J*_{P-H} = 11.1 Hz, P-OCH₃); 2.37 (s, 3H, Ar-CH₃). ³¹P NMR (DMSO-*d*₆); δ 26.76. Anal. Calcd. for C₁₈H₂₃BrNO₅P: C, 48.66; H, 5.22; N, 3.15. Found: C, 48.55; H, 5.16; N, 3.08%.

Dimethyl(4-bromo-3-methylphenylamino)(4-N, N-dimethylphenyl) methyl phosphonate (4g)

Yield (60%), mp 145-146 °C; IR (KBr) cm^{-1} ; 3385 (P-NH), 1238 (P=O), 758 (P-CH); $^1\text{H-NMR}$ (DMSO- d_6) δ 6.72-7.58 (m, 7H, Ar-H), 5.14 (s, 1H, NH), 4.86-4.90 (m, 1H, P-CH); 3.31-3.39 (d, $^3J_{\text{P-H}} = 10.1$ Hz, P-OCH₃); 3.55-3.65 (d, $^3J_{\text{P-H}} = 11.3$ Hz, P-OCH₃), 2.84 (s, 6H, N(CH₃)₂), 2.28 (s, 3H, Ar-CH₃); ^{31}P NMR (DMSO- d_6); δ : 31.30. Anal. Calcd. for C₁₈H₂₄BrN₂O₃P: C, 50.60; H, 5.66; N, 6.56. Found: C, 50.48; H, 5.57; N, 6.51%.

Dimethyl (4-bromo-3-methylphenylamino) (phenyl) methylphosphonate (4h)

Yield (69%), mp 143-144°C; IR (KBr) cm^{-1} ; 3387 (P-NH), 1232 (P=O), 755 (P-CH); $^1\text{H-NMR}$ (DMSO- d_6) δ 6.52-7.66 (m, 8H, Ar-H), 5.04 (s, 1H, NH), 4.70-4.85(m, 1H,P-CH), 3.29-3.41 (d, $^3J_{\text{P-H}} = 10.0$ Hz, P-OCH₃); 3.54-3.63 (d, $^3J_{\text{P-H}} = 11.2$ Hz, P-OCH₃) 2.26 (s, 3H, Ar-CH₃); ^{31}P NMR (DMSO- d_6) δ 30.30. GCMS m/z (%): 384 (M+), 385.0 (8), 369.4 (16), 262.3 (100), 215.6 (18), 172.5 (9), 110.2 (5): Anal.Calcd. for C₁₆H₁₉BrNO₃P: C, 50.02; H, 4.98; N, 3.65. Found: C, 49.91; H, 4.91; N, 3.61%.

Dimethyl (4-bromo-3-methylphenylamino) (3-chlorophenyl) methylphosphonate (4i)

Yield (73%), mp 161-162°C; IR (KBr) cm^{-1} ; 3350 (P-NH), 1218 (P=O), 780 (P-CH); ^1H NMR(DMSO- d_6) δ : 6.53-7.76 (m, 7H, Ar-H), 5.03 (s, 1H, NH), 4.76-4.80 (m, 1H, P-CH), 3.36-3.44 (d, $^3J_{\text{P-H}} = 10.1$ Hz, P-OCH₃); 3.61-3.69 (d, $^3J_{\text{P-H}} = 11.1$ Hz, P-OCH₃), 2.21 (s, 3H, Ar-CH₃); ^{31}P NMR(DMSO- d_6) δ 20.18. Anal. Calcd. for C₁₆H₁₈BrClNO₃P: C, 45.90; H, 4.33; N, 3.35. Found: C, 45.81; H, 4.27; N, 3.29%.

Dimethyl (4-bromo-3-methylphenylamino) (4-chlorophenyl) methylphosphonate (4j)

Yield (74%), mp 157-158 °C; IR (KBr) cm^{-1} 3372 (P-NH), 1211 (P=O), 776 (P-CH). $^1\text{H-NMR}$ (DMSO- d_6) δ 6.75-7.54(m, 7H, Ar-H), 5.12 (s, 1H, NH), 4.79-4.84 (m, 1H, P-CH), 3.42-3.49 (d, $^3J_{\text{P-H}} = 9.8$ Hz, P-OCH₃), 3.71-3.78 (d, $^3J_{\text{P-H}} = 10.5$ Hz, P-OCH₃), 2.17 (m, 3H, Ar-CH₃); ^{31}P NMR (DMSO- d_6) δ 31.80. Anal. Calcd. for C₁₆H₁₈BrClNO₃P: C, 45.90; H, 4.33; N, 3.35. Found: C, 45.78; H, 4.25; N, 3.28%.

Results and Discussion

4-Bromo 3-methyl benzenamine (**1**) was treated with different aldehydes (**2a-j**) and dimethyl phosphite (**3**) in the presence of 10 mole % of tetramethylguanidine (TMG) in dry toluene at RT. The mixture was stirred at RT for 1 h and at 70-80 °C for another 5 h. The progress of the reaction was monitored by thin layer chromatography. The reaction proceeded smoothly, and completed in 5-6 h to afford the corresponding α -aminophosphonates in high yield (60-74%). This showed that TMG acts as an effective catalyst in this reaction. An important feature is that the TMG can be easily recovered from the reaction mixture after completion of the reaction and can be reused. The chemical structures of all the new compounds were confirmed by elemental analysis, IR, ^1H , ^{13}C and ^{31}P NMR spectra. Compounds **4a-j** exhibited characteristic IR stretching frequencies in the regions 3225-3390, 1211-1238, 742-780 cm^{-1} for N-H, P=O and P-C(aliphatic) respectively¹⁸. The aromatic protons of the benzene rings of the α -aminophosphonates (**4a-j**) showed a complex multiplet at δ 6.12-7.76. The P-C-H proton signal appeared as a multiplet¹⁹ at δ 4.62-4.93 due to its coupling with both phosphorus and the N-H proton. The exocyclic N-H proton signal appeared at δ 4.87-5.14 as a singlet. The methoxy group protons of the dimethylphosphite moiety resonated as two distinct doublets in the range of δ 3.22 – 3.49 (d, $^3J_{\text{P-H}} = 9.8$ -10.1 Hz) and δ 3.54-3.79 (d, $^3J_{\text{P-H}} = 10.5$ -11.30)^{19,20} indicating their non-equivalence. The carbon chemical shifts for P-CH-, P-O-CH₃

in the title compounds were observed at δ 54.90-55.83, 53.27-54.00 respectively^{18,21,22}. The ³¹P NMR signals²³ appeared in the region δ 20.18-31.80 for these compounds.

Antimicrobial activity

The antibacterial activity of (**4a-j**) was assayed²⁴ against the growth of *staphylococcus aureus*(gram +ve) and *Escherichia coli* (gram -ve) at three concentrations (100, 50, 25 ppm) (Table 1). The majority of the compounds exhibited moderate to good activity against both the bacteria. The same (**4a-j**) were screened for their antifungal activity²⁵ (Table 2) against *Aspergillus niger* and *Helminthosporium oryzae* species along with the standard fungicide Griseofulvin at three different concentrations (100, 50, 25 ppm). It is gratifying to observe that the majority of the compounds **4a-j** exhibited moderate to good antifungal activity when compared with the Griseofulvin reference.

Table 1. Antibacterial activity of α -aminophosphonates (**4a-j**)

Compound	Zone of inhibition, %					
	<i>Escherichia coli</i>			<i>Staphylococcus aureus</i>		
	100	50	25	100	50	25
4a	08	05	02	09	05	01
4b	07	05	03	09	04	02
4c	09	04	-	08	04	-
4d	05	03	01	07	04	-
4e	09	04	-	06	05	-
4f	08	05	03	10	06	02
4g	08	04	-	09	04	01
4h	08	05	-	08	05	02
4i	08	05	01	06	03	-
4j	05	02	-	07	03	-
Pencillin	12	07	-	11	08	-

Table 2. Antifungal activity of α -aminophosphonates (**4a-j**)

Compound	Zone of inhibition, %					
	<i>Aspergillus Niger</i>			<i>Helmenthosphorium oryzae</i>		
	100	50	25	100	50	25
4a	07	05	03	10	06	03
4b	07	05	03	08	05	02
4c	09	06	04	09	05	03
4d	05	03	01	07	04	-
4e	09	04	-	06	05	-
4f	08	05	03	10	06	02
4g	08	04	-	09	04	01
4h	09	04	-	08	04	-
4i	08	05	02	09	05	01
4j	07	03	-	08	04	01
Griseo – fulvin	11	08	-	13	08	-

Acknowledgment

The authors express their thanks to Prof C. Suresh Reddy, S.V. University, Tirupati for his suggestions and Dr. C. Chellaram Department of Biomedical Engineering, VelTech MultiTech Dr.R.R.& Dr.S.R Engineering college for his help in biological studies. We extend our thanks to Director IISc Bangalore for providing spectral data.

References

1. Kaboudin B, Nazari R, *Tetrahedron Lett.*, 2001, **42**, 8211.
2. Matveeva E D, Podrugina T A, Tishkovskaya E V, Tomilova L G and Zefirov N S, *Synlett.*, 2003, 2321-2324.
3. Pavlov V. Ya, Kabachinik M M, Zobnina E V, Ponomarev G V and Beletskaya I P, *Synlett.*, 2003, 2193.
4. Sikorski J A, Miller M J, Braccolino D S, Cleary D G, Corey S D, Ream J E, Schnur D, Shah A and Walker M C, *Phosphorus Sulfur Silicon.*, 1993, **76**, 375.
5. Stowasser B, Budt K H, Jain-Qi L, Peyman A and Ruppert D, *Tetrahedron Lett.*, 1992, **33**, 6625.
6. Patel D V, Reilly-Gauvin K and Ryono D E, *Tetrahedron Lett.*, 1990, **31**, 5587.
7. (a) Bruke T R, Jr Brachi J J, Jr George C, Wolf G, Shoelson S E and Yan X, *J Med Chem.*, 1995, **38**, 1386-1396; (b) Bruke T R, Jr Kole H K and Roller P P, *Biochem Biophys Res Commun.*, 1994, **204**, 129-134.
8. Peyman A, Budt K H, Paning J S, Stowasser B and Ruppert D, *Tetrahedron Lett.*, 1992, **33**, 4549-4552.
9. Atherton F R, Hassall C H and Lambert R W, *J Med Chem.*, 1986, **29**, 29-40.
10. Kafarski P, Lejack B and Mastalerz P, *Beitr Wirk Forsh.*, 1985, H25; *Chem Abstr.*, 1985, **103**, 174532.
11. (a) Maier L and Lea P J, *Phosphorus Sulfur Silicon* 1983, **17**, 1; (b) Giannousis P P, Bartlett P A, *J Med Chem.*, 1987, **30**, 1603-1609; (c) Gancarz R and Wieczorek J S, *Synthesis*, 1977, 625; (d) Baylis E K, Campbell C D and Dingwall J G, *J Chem Soc., PerkinTrans 1*, 1984, 2845-2853.
12. Fields S C, *Tetrahedron*, 1999, **55**, 12237.
13. Kleszczynska H and Sarapuk J, *Cell Mol Biol Lett.*, 2001, **6(1)**, 83-91.
14. (a) Kabachinik M J and Medved T, *Izv Akad Nauk SSSR*, 1953, 1126; (b) Kabachinik M J and Medved T, *Izv Akad Nauk SSSR*, 1954, 1024.
15. Fields E K, *J Am Chem Soc.*, 1952, **74**, 1528.
16. Pollini G P, Barco A and Degiuli G, *Synthesis*, 1924, 44.
17. Hewson A. T and Macpherson D T, *Tetrahedron Lett.*, 1983, **24**, 647.
18. Silverstein R M and Webster F X, *Spectrometric Identification of Organic Compounds*, 6th Edn., Wiley: New York, 1998.
19. Tongcharoensirikul P, Suarez A I, Voelker T and Thomson C M, *J Org Chem.*, 2004, **69**, 2322.
20. Heydari A, Zarei M, Alijanizadeh R and Tavakol H, *Tetrahedron Lett.*, 2001, **42**, 3629.
21. Qian C and Huang, T, *J Org Chem.*, 1998, **63**, 4125.
22. Semenzin D, Moghadam G E, Albiouy D, Diallo O and Koenig M, *J Org Chem.*, 1997, **62**, 2414.
23. Quin L D and Verkade J G, *Phosphorus-31 NMR Spectral Properties in Compound Characterization and Structural Analysis*; VCH: New York, 1994
24. Vincent J C and Vincent H W, *Proc Soc Expt Biol Med.*, 1944, **55**, 162.
25. Benson H J, *Microbiological Applications*, 5th Edn., Brown W C, Publications: Boston, 1990.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

