

Review Article

The Role of Endoplasmic Reticulum Stress in Autoimmune-Mediated Beta-Cell Destruction in Type 1 Diabetes

Jixin Zhong,^{1,2,3} Xiaoquan Rao,^{2,3} Jun-Fa Xu,⁴ Ping Yang,^{1,2} and Cong-Yi Wang^{1,2,4}

¹The Center for Biomedical Research, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, 1095 Jiefang Avenue, Wuhan 430030, China

²The Center for Biotechnology and Genomic Medicine, Medical College of Georgia, 1120 15th Street, CA4098, Augusta, GA 30912, USA

³Affiliated Hospital of Guangdong Medical College, 57 Ren-Ming Road, Zhanjiang 524001, China

⁴The Department of Clinical Immunology, Guangdong Medical College, 1 Xincheng Avenue, Dongguan 523808, China

Correspondence should be addressed to Cong-Yi Wang, cwang@georgiahealth.edu

Received 11 July 2011; Accepted 27 November 2011

Academic Editor: Muthuswamy Balasubramanyam

Copyright © 2012 Jixin Zhong et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Unlike type 2 diabetes which is caused by the loss of insulin sensitivity, type 1 diabetes (T1D) is manifested by the absolute deficiency of insulin secretion due to the loss of β mass by autoimmune response against β -cell self-antigens. Although significant advancement has been made in understanding the pathoetiology for type 1 diabetes, the exact mechanisms underlying autoimmune-mediated β -cell destruction, however, are yet to be fully addressed. Accumulated evidence demonstrates that endoplasmic reticulum (ER) stress plays an essential role in autoimmune-mediated β -cell destruction. There is also evidence supporting that ER stress regulates the functionality of immune cells relevant to autoimmune progression during T1D development. In this paper, we intend to address the role of ER stress in autoimmune-mediated β -cell destruction during the course of type 1 diabetes. The potential implication of ER stress in modulating autoimmune response will be also discussed. We will further dissect the possible pathways implicated in the induction of ER stress and summarize the potential mechanisms underlying ER stress for mediation of β -cell destruction. A better understanding of the role for ER stress in T1D pathoetiology would have great potential aimed at developing effective therapeutic approaches for the prevention/intervention of this devastating disorder.

1. Introduction

Recent epidemiologic studies revealed that the incidence of type 1 diabetes (T1D) in most regions worldwide has been increasing by 2% to 5% [1]. Particularly, in some developing countries such as China, the rapid economic development along with changes in lifestyle and presumably the living environment has rendered this country with an annual increase of 7.4% for T1D prevalence [2]. Given that T1D is typically developed in children and juveniles, its impact on the quality of life is far more significant than that of type 2 diabetes, in which it usually occurs in adults. Although exogenous insulin therapy partly compensates the function of β cells, it cannot regulate blood glucose as accurately as the action of endogenous insulin. As a result, long-term improperly control of blood glucose homeostasis predisposes T1D patients

to the development of diverse complications such as diabetic retinopathy [3–5], nephropathy [6, 7], neuropathy [8–10], foot ulcers [11–13], and cardiovascular diseases [14–16]. Although the underlying mechanisms leading to T1D have yet to be fully addressed, extensive studies have consistently demonstrated that endoplasmic reticulum (ER) stress plays a critical role in autoimmune-mediated β -cell destruction during the course of T1D development.

The pancreatic β -cells are equipped with highly developed endoplasmic reticulum (ER) to fulfill the requirement of secreting a large amount of insulin. This physiological feature renders β cells particularly vulnerable to ER stress [17]. Exhaustion of β cells is essential for the onset of T1D, which requires the residual β cells for compensated insulin secretion. While this compensated action is beneficial for control of blood glucose homeostasis, it also increases ER burden

associated with the induction of unfolded protein response (UPR) and ER stress, which further exacerbates β -cell death. Although the implication of ER stress in β -cell death has been extensively emphasized, the underlying mechanisms, however, are yet to be fully elucidated. As such, understanding the role of ER stress in the loss of β mass and dissecting the mechanisms underlying ER stress would be important for developing therapeutic approaches aimed at prevention and intervention of type 1 diabetes. In the present paper, we will first intend to address the overall role of ER stress in autoimmune-mediated β -cell destruction based on published genetic and experimental data. The impact of ER stress on modulation of autoimmune response during the course of T1D development will be next discussed. We will finally focus on the possible pathways implicated in the induction of ER stress and summarize the potential mechanisms underlying ER stress for mediation of β -cell destruction.

2. The Endoplasmic Reticulum (ER)

ER is a membranous network of tubules, vesicles, and cisternae that are interconnected by the cytoskeleton in the cytoplasm of eukaryotic cells. ER is responsible for many general cellular functions, including the facilitation of protein folding and assembly [18–20], manufacture of the membranes [21], biosynthesis of lipid and sterol, storage of intracellular Ca^{2+} , and transport of synthesized proteins in cisternae.

ER can be categorized into rough endoplasmic reticulum (RER) and smooth endoplasmic reticulum (SER). RER is responsible for protein synthesis, while SER is in charge of the synthesis of lipids and steroids, regulation of calcium concentration, attachment of receptors on cell membrane proteins, and detoxification of drugs. As featured by its name, RER bears ribosomes on the outer surfaces of the cisternae and looks bumpy and rough under a microscope. The newly synthesized proteins by RER are sequestered in cisternae and sent to Golgi complex or membrane via small vesicles. In contrast, SER does not have ribosomes on its cisternae and appears to have a smooth surface under the microscope. SER is found commonly in places such as in the liver and muscle. It is important for the liver to detoxify poisonous substances. Sarcoplasmic reticulum (SR) is a special type of SER, which is found in smooth and striated muscle. SR is responsible for the regulation of calcium levels. It sequesters a large store of calcium and releases them when the muscle cell is stimulated.

3. ER Stress

ER stress is the cellular responses to the disturbances of normal function of ER. The most focused and well-studied ER stress is that caused by protein misfolding. The accumulation of unfolded proteins leads to a protective pathway to restore ER function, termed as unfolded protein response (UPR). ER employs a type of special proteins called chaperones as a quality control mechanism. Chaperones attach to the newly synthesized proteins and assist them to fold into their native conformations. In addition, chaperones also help to break

down unfolded or incorrectly folded proteins in the ER via a process called ER-associated degradation (ERAD). Protein folding requires a serial of ER-resident protein folding machinery. Exhaustion of those protein folding machineries or insufficient energy supply increases the accumulation of unfolded or misfolded proteins in ER, leading to the activation of UPR. Various physiological and pathological insults such as increased general protein synthesis, failure of post-translational modifications, hypoxia, nutrient/glucose starvation, and alterations in calcium homeostasis can result in the accumulation of unfolded or misfolded proteins in ER which then causes ER stress [22]. For example, altered expression of antithrombin III [23, 24] or blood coagulation factor VIII [25, 26] results in the exhaustion of protein-folding machinery and thus induces UPR. Some physiological processes such as the differentiation of B lymphocytes into plasma cells along with the development of highly specialized secretory capacity can also cause accumulation of unfolded proteins and induce UPR [27–29]. In response to certain physiological and pathological insults, cells undergo UPR to get rid of the unfolded or misfolded proteins. Therefore, UPR is a protective mechanism by which it monitors and maintains the homeostasis of ER. For instance, UPR increases the folding capacity by upregulating ER chaperones and foldases, and attenuates the biosynthetic burden of secretory pathway through downregulating the expression of secreted proteins [30–32]. In addition, UPR also activates ERAD to eliminate unfolded proteins [33–35] (Figure 1). However, once the stress is beyond the compensatory capacity of UPR, the cells would undergo apoptosis. As such, UPR and ER stress are reported to be implicated in a variety of pathological processes, including diabetes, neurodegenerative diseases, pathogenic infections, atherosclerosis, and ischemia [22, 36].

As aforementioned, there is a monitoring mechanism to ensure the correct protein folding in ER. The unfolded proteins usually have a higher number of hydrophobic surface patches than that of proteins with native conformation [37]. Thus, unfolded proteins are prone to aggregate with each other in a crowded environment and directed to degradative pathway [38]. Molecular chaperones in ER are the major mechanisms to promote protein folding. They preferentially interact with hydrophobic surface patches on unfolded proteins and create a private folding environment by preventing unfolded proteins from interaction and aggregation with other unfolded proteins. In addition, the concentration of Ca^{2+} in ER also impairs protein folding by inhibiting the activity of ER-resident chaperones and foldases [39–42]. ER is the major site for Ca^{2+} storage in mammalian cells. The concentration of Ca^{2+} in ER is thousands times higher than that in the cytosol [43]. Most chaperones and foldases in ER are vigorous Ca^{2+} binding proteins. Their activity, therefore, is affected by the concentration of Ca^{2+} in ER. A variety of posttranslational modifications including N-linked glycosylation, disulfide bond formation, lipidation, hydroxylation, and oligomerization occur in ER. Disruption of those post-translational modifications can also result in the accumulation of incorrectly folded proteins and thereby induce UPR or ER stress. For example, glucose deprivation impairs




FIGURE 1: The regulatory role of unfolded protein response (UPR). Various physiological and pathological insults can result in the accumulation of unfolded proteins which then induces UPR and ER stress. In response to stressful insults, UPR regulates secretory pathway via following mechanisms: (1) enhancing (red arrow) the expression of ER chaperones and foldases to increase the folding capacity of ER; (2) attenuating (blue) the biosynthetic burden of secretory pathway through downregulating the expression of secreted proteins (purple arrow); (3) promoting the clearance of unfolded proteins by activating ERAD (orange arrow).

the process for N-linked protein glycosylation and thus leads to ER stress [44].

UPR is mediated by three major pathways, which are initiated by the three transmembrane signaling proteins located on the ER membrane. Those transmembrane proteins function as a bridge to link cytosol and ER with their C-terminal in the cytosol and N-terminal in the ER lumen. The N-terminal is usually engaged by an ER-resident chaperone BiP (Grp78) to avoid aggregation. When unfolded proteins accumulate in ER, chaperones are occupied by unfolded proteins and release the transmembrane signaling proteins: which include the following three axes of signals: the pancreatic endoplasmic reticulum kinase (PERK), the inositol-requiring enzyme 1 (IRE1), and the activating transcription factor 6 (ATF6). The release of these proteins triggers UPR and ER stress (Figure 2). PERK is a Ser/Thr protein kinase uniquely present in ER. Once released from BiP, PERK becomes oligomerized and autophosphorylated. PERK inactivates eukaryotic initiation factor 2 α (eIF2 α) by phosphorylation of Ser51 to reduce mRNA translation and protein load on ER. Deficiency of PERK results in an abnormally elevated protein synthesis in response to the accumulation of unfolded proteins in ER. IRE1 is another axis of signal involved in UPR. IRE1 increases the production of X box protein-1 (XBP-1), a bZIP-family transcriptional factor, by promoting its mRNA splicing [45]. XBP-1 heterodimerizes with NF-Y and enhances gene transcription by binding to the ER stress enhancer (ERSE) and unfolded protein response element

(UPRE) in the promoters of targeted genes. Unlike PERK and IRE1 which oligomerize upon UPR, when released from BiP, ATF6, the third axis of signal, translocates into the Golgi apparatus where its transmembrane domain is cleaved [46]. The cleaved ATF6 is then relocated into the nucleus to regulate the expression of targeted genes. For example, once released from the ER membrane, ATF6 enhances the transcription of XBP-1 mRNA which is further regulated by IRE1 [45].

4. ER Stress in Autoimmune-Mediated β -Cell Destruction

Accumulative evidence supports that ER stress is implicated in autoimmune-mediated β -cell destruction in type 1 diabetes [47, 48]. It was noted that loss of β cells is the direct causing factor for insufficient insulin secretion in T1D patients. As described earlier, pancreatic β cells have a very well-developed ER to fulfill their biological function for secreting insulin and other glycoproteins, and therefore, β cells are highly sensitive to ER stress and the subsequent unfolded protein response (UPR). Severe or long-term ER stress would direct β cells undergoing apoptosis [47]. For example, mice deficient in PERK, a molecule responsible for regulating UPR, are extremely susceptible to diabetes. The *null* mice display a progressive loss of β mass and hyperglycemia with aging [49]. Consistent with the observations in these mice, some infant-onset diabetes in humans have also been




FIGURE 2: Signaling pathways relevant to UPR. PERK, IRE1, and ATF6 act as ER stress sensors by binding to the ER chaperone BiP, and by which they remain inactive under normal condition. Upon the accumulation of unfolded proteins, BiP preferentially binds to the unfolded proteins, which results in the release of PERK, IRE1, and ATF6. Once released from BiP, PERK becomes activated and dimerized. Activated PERK phosphorylates eIF2 α to suppress the overall transcription of mRNAs while selectively enhance the transcription of genes implicated in UPR such as the ATF4 mRNA, and through which ATF4 initiates the transcription of UPR target genes. Similar to PERK, IRE1 is dimerized and activated after detached from BiP. IRE1 induces XBP-1 by promoting the splicing of its mRNA. XBP-1 activates the transcription of its target genes to enhance UPR. The release of ATF6 from BiP results in the translocation of ATF6 to the Golgi apparatus, where ATF6 is cleaved and then translocates into the nucleus, and by which ATF6 initiates the transcription of target genes.

confirmed to be associated with the mutations in PERK. For example, loss of *EIF2AK3* (the gene encodes PERK) develops Wolcott-Rallison syndrome, an autosomal recessive disorder characterized by early infancy insulin-dependent diabetes and multisystemic manifestations including growth retardation, hepatic/renal dysfunction, mental retardation, and cardiovascular abnormalities [50, 51]. Similarly, disruption of UPR by mutating eIF2 α , a protein that controls mRNA translation upon ER stress, enhances the sensitivity to ER stress-induced apoptosis and results in defective gluconeogenesis. Mice carrying a homozygous Ser51Ala mutation for eIF2 α show defective in pancreatic β cells manifested by the smaller core of insulin-secreting β cells and attenuated insulin secretion [52]. Altogether, defects in PERK/eIF2 α signaling render β cells highly vulnerable to ER stress in both humans and mice [53, 54].

In type 1 diabetes, ER stress in the pancreatic β cells is primarily induced by proinflammatory cytokines produced by infiltrated immune cells, which then contributes to β -cell destruction. During the course of autoimmunity, proinflammatory cytokines are secreted by the infiltrated autoreactive immune cells in the milieu of pancreatic islets. For example, nitric oxygen (NO) is an inflammatory mediator resulted from autoimmune response during the course of type 1 diabetes. Studies have shown that excessive NO production induces β -cell apoptosis in a CHOP-dependent manner [55].

Other than ER stress caused by autoimmunity, misfolding of insulin in β cells can also directly induce chronic ER stress as evidenced by the observations in Akita mice. The Akita mouse carries a mutation for the *Ins2* gene which disrupts a disulfide bond between the α and β chain of proinsulin, leading to the mis-folding of the mutated insulin, and by which the mutated insulin induces ER stress in β cells to cause diabetes [56].

It is likely that inflammatory cytokines produced by islet-infiltrated autoreactive immune cells are the major factors causing β -cell death in type 1 diabetes [57]. In the early stage of type 1 diabetes, the autoreactive immune cells such as macrophages and T lymphocytes infiltrate into the pancreatic islets along with the secretion of inflammatory cytokines such as IL-1 β , IFN- γ , and TNF- α , which then induce ER stress to mediate β -cell destruction. The damaged or dying β cells also release danger signals such as high-mobility group box 1 and heat shock proteins (HSPs), to alert the immune system for the presence of β -cell injury, which in turn further promotes autoimmune progression [57–60]. Studies have shown that stimulation of β cells with IL-1 β and IFN- γ induces the expression of death protein 5 (DP5), and through which these cytokines mediate β -cell apoptosis via ER stress [61]. Knockdown of DP5 provides protection for β cells against inflammatory cytokine-induced ER stress [61]. Insult of β cells with IL-1 β and IFN- γ has also been found to

decrease the expression of sarcoendoplasmic reticulum pump Ca^{2+} ATPase (SERCA) 2b, which controls the storage of ER Ca^{2+} [62]. It has been well demonstrated that altered ER Ca^{2+} concentration induces the accumulation of unfolded proteins in ER associated with the induction of UPR and ER stress in β cells [63].

Given that hyperglycemia only occurs when β cells fail to compensate the increased demand for insulin, β cells are usually “exhausted” in T1D patients [54]. Therefore, other than the ER stress induced by autoimmune response, β cells in T1D patients are also under ER stress caused by altered insulin synthesis. In later case, the increased insulin demand requires the remaining functional β cells to increase insulin synthesis to compensate the decrease of β mass. While this process in short term is beneficial for control of blood glucose homeostasis, it also induces ER stress, which in turn exacerbates β -cell dysfunction to promote disease progression and diabetes onset. Collectively, there is convincing evidence that ER stress plays an essential role in β -cell destruction during the course of T1D development.

5. The Impact of ER Stress on Modulation of Autoimmune Response

Unlike its well-defined effect on autoimmune-mediated β -cell destruction in type 1 diabetes, the impact of ER stress in modulating autoimmune response during the course of type 1 diabetes, however, remains poorly elucidated. There is evidence supporting that other than its critical roles played in β -cell destruction, ER stress also modulates the functionality of immune cells with implications in autoimmune response in type 1 diabetes.

It has been well accepted that the presence of β -cell-specific autoantibodies serves as a marker for the initiation and progression of autoimmunity in type 1 diabetes [64]. Studies have shown that IRE1, a key molecule in UPR, modulates the differentiation of antibody-producing B lymphocytes. Deficiency of IRE1 hampers pro-B cells differentiating into pre-B cells [65], and XBP-1, an IRE1 downstream molecule, is required for antibody production by mature B cells [66]. It was found that the engagement of B-cell receptor (BCR) induces ubiquitin-mediated degradation of BCL-6, a repressor for B-lymphocyte-induced maturation protein 1 (BLIMP1) [67], while BLIMP1 negatively regulates the expression of B-cell-lineage-specific activator protein (BSAP) [68], and BSAP is suggested to function as a repressor for XBP-1 [69]. In line with these results, B lymphocytes deficient in BLIMP1 failed to express XBP-1 in response to LPS stimulation [66].

Recent studies highlighted the importance of innate immunity in the pathogenesis of type 1 diabetes [59, 60], while elements of the UPR pathway are found to regulate innate immune response [70]. The expression of CREBH, an ER stress-associated transcription factor, can be induced by inflammatory cytokines such as IL-1 β and IL-6, which in turn regulates the transcription of serum amyloid P-component and C-reactive protein, the two critical factors implicated in innate immune responses [71]. Furthermore, the differentiation of dendritic cells (DCs), the most critical innate

immune cells, is regulated by UPR signaling element, XBP-1 [72]. High levels of mRNA splicing for XBP-1 are found in DCs, and mice deficient in XBP-1 show altered development of both conventional and plasmacytoid DCs. Loss of XBP-1 renders DCs vulnerable to ER stress-induced apoptosis [72]. Moreover, the capacity for DCs secretion of inflammatory cytokine IL-23 is regulated by CHOP, a UPR mediator. CHOP can directly bind to the *IL-23* gene and regulate its transcription. ER stress combined with Toll-like receptor (TLR) agonists was found to markedly increase the mRNA of IL-23 p19 subunit and the secretion of IL-23, while knock-down of CHOP suppressed the induction of IL-23 by ER stress and TLR signaling [73].

Richardson and coworkers reported that innate immune response induced by *P. aeruginosa* infection causes ER stress in *C. elegans*, and mutations with loss of function for XBP-1 lead to larval lethality [74]. In consistent with this result, the polymorphisms within the *XBP-1* gene were found to be associated with Crohn’s disease and ulcerative colitis in humans [75], and the two autoimmune diseases share similar properties as type 1 diabetes. Loss of XBP-1 in intestinal epithelial cells induces Paneth cell dysfunction and overactive epithelium, leading to impaired mucosal defense to *Listeria monocytogenes* and increased sensitivity to colitis [75].

Other than the IRE1/XBP-1 axis, the PERK/eIF2 α /ATF4 axis of UPR is also found to be associated with innate response. TLR signaling, the most important innate signaling pathway, is reported to induce selective suppression of the ATF-4/CHOP axis of UPR pathway [76]. TLR signaling decreases eIF2 α -induced ATF4 translation. For example, pretreatment of LPS, an agonist for TLR4, suppressed ATF4/CHOP signaling and prevented systemic ER stress-induced apoptosis in macrophages, renal tubule cells, and hepatocytes [76]. In contrast, loss of Toll-IL-1R-containing adaptor inducing IFN- β (TRIF), an important adapter for TLR signaling, abrogated the protective effect of LPS on systemic ER stress-induced renal dysfunction and hepatosteatosis, suggesting that TLR signaling suppresses ATF4/CHOP via a TRIF-dependent pathway [76].

6. Pathways for Cytokines Induction of ER Stress

Upon the insults of pathogens, mutated self-antigens, or tissue damage, the immune system initiates inflammatory response by releasing copious amount of cytokines. UPR and ER stress are interconnected with inflammatory cytokines through multiple mechanisms including reactive oxygen species (ROS), NF κ B, and JNK (Figure 3). ROS are highly reactive small molecules with unpaired electrons. They are important mediators of inflammatory response. The accumulation of ROS, referred to as oxidative stress, was confirmed to be associated with ER stress [77]. Oxidizing condition is required for the disulphide bond formation during the process of protein folding [78]. Increased protein folding load may lead to oxidative stress. The PERK axis of UPR signaling is reported to be able to activate antioxidant pathway by promoting ATF4 and nuclear factor-erythroid-derived




FIGURE 3: The possible implication of UPR in inflammatory response. UPR is associated with inflammation via a variety of mechanisms involving ROS, JNK, and NF κ B. PERK promotes ATF4 and NRF2, which then suppress ROS production by activating antioxidant pathway. Upon activation, IRE1/TRAF2 recruits IKK, leading to the phosphorylation of I κ B α and subsequent activation of NF κ B. IRE1/TRAF2 can also activate AP1, resulting in the activation of JNK. XBP-1 induced by IRE1 can further induce the expression of various genes implicated in inflammation. Furthermore, ATF6, the other axis of UPR signaling, can promote inflammation via activating NF κ B.

2-related factor 2 (NRF2) [79, 80]. Therefore, loss of PERK markedly increases ROS accumulation induced by toxic chemicals [79, 81]. The IRE1/TRAF2 axis of UPR can recruit I κ B kinase (IKK), leading to the activation of NF κ B, a key regulator in inflammation [82]. As a result, NF κ B activation and TNF- α production are reduced in cells lacking IRE1 [82]. Furthermore, the IRE1/TRAF2 axis can activate JNK, and by which it induces the expression of inflammatory genes by activating activator protein 1 (AP1) [83]. ATF6, the third axis of UPR signaling, can also activate NF κ B pathway, in which suppression of ATF6 reduces NF κ B activation caused by BiP degradation [84].

Other than the above described pathways, cytokines may also induce ER stress via inducible nitric oxide (NO) synthase (iNOS) and JNK pathway. JNK pathway is activated by IL-1 β . Suppression of JNK by its inhibitor SP600125 protected β cells from IL-1 β -induced apoptosis [85]. Cytokines have been evidenced to induce the expression of iNOS, leading to excessive NO production. Stimulation with IL-1 β and IFN- γ activates ER stress pathway and induces β -cell apoptosis via NO synthesis [62]. NO has been suggested to be an important mediator of β -cell death in type 1 diabetes. Inflammatory cytokines including IL-1 β , IFN- γ , and TNF- α can induce iNOS expression in β cells which then produces copious amount of NO [50]. Excessive NO induces DNA damage and thus results in β -cell apoptosis through p53 pathway or necrosis through poly(ADP-ribose) polymerase (PARP) pathway [54]. Moreover, NO depletes ER Ca²⁺ stores via activating Ca²⁺ channels or inhibiting Ca²⁺ pumps [86–88]. Depletion of Ca²⁺ then leads to ER stress and apoptosis in β cells via the induction of CHOP signaling [55, 89].

7. Mechanisms Underlying ER-Stress-Induced β -Cell Death

ER stress is a key mediator for β -cell death in type 1 diabetes. The primary purpose of ER stress or UPR is to compensate the damage caused by the disturbances of normal ER function. However, continuous ER dysfunction would eventually render cells undergoing apoptosis. The mechanisms by which ER stress induces cell death are not fully elucidated, due to the fact that multiple potential participants involved but little clarity on the dominant death effectors in a particular cellular context. In general, ER stress induction of cell death can be illustrated in three phases: adaptation, alarm, and apoptosis [44].

The phase for adaptation response is initiated to restore the homeostasis of ER and to protect cells from damage induced by the disturbances of ER function. As described earlier, the signaling for UPR involves three axes of responses: IRE1, PERK, and ATF6. These axes interact between each other and form a feedback regulatory mechanism to control the activity of UPR. The accumulation of unfolded proteins in ER results in the engagement of ER-resident chaperon BiP, and as a consequence, IRE1, PERK, and ATF6 are released from BiP. Therefore, overexpression of BiP can prevent cell death induced by oxidative stress, Ca²⁺ disturbances, and hypoxia [90]. PERK is oligomerized and phosphorylated when released from BiP. Activated PERK inactivates eIF2 α to reduce mRNA translation and protein load on ER. Therefore, PERK deficiency results in an abnormally elevated protein synthesis in response to the accumulation of unfolded proteins in ER, which renders cells highly sensitive to ER stress

and ER stress-induced apoptosis [91]. Similarly, as a downstream molecule of PERK, eIF2 α is required for cell survival upon the insult of ER stress, and a mutation at the phosphorylation site of eIF2 α (Ser51Ala) abolishes the translational suppression in response to ER stress [52]. Similar as PERK, IRE1 becomes dimerized and activated once released from BiP. IRE1 induces XBP-1 by promoting the splicing of its mRNA [45]. XBP-1 is a transcriptional factor belonging to the bZIP-family and is responsible for the transcription of many adaptation genes implicated in UPR. Unlike PERK and IRE1, ATF6 translocates into the Golgi apparatus upon the release from BiP. The transmembrane domain of ATF6 is cleaved in the Golgi apparatus and is then relocated into the nucleus, by which it regulates gene expression [46].

During the alarm phase, many signal pathways are activated, and the expression of responsive genes has been induced to alert the system. For example, the cytoplasmic part of IRE1 binds to TNF receptor-associated factor 2 (TRAF2), a key adaptor for TNF-mediated innate immune signaling. TRAF2 would then activate NF κ B pathway via activating IKK and activate the signaling for c-Jun N-terminal kinases (JNK) by apoptosis signal-regulating kinase 1 (Ask1). Studies have shown that dominant negative TRAF2 suppresses the activation of JNK by IRE1 in response to ER stress [92]. Importantly, TRAF2 is also a critical component for E3 ubiquitin-protein ligase complex [93], which binds to Ubc13 and promotes the noncanonical ubiquitination of substrates. The Ubc13-dependent ubiquitination of TRAF2 is suggested to be required for the activation of JNK [94]. In addition, IRE1 can further activate JNK signaling through interacting with c-Jun N-terminal inhibitory kinase (JIK) [95].

Although the purpose for the initiation of adaptation response is to restore the homeostasis of ER, apoptosis however could occur, once the accumulation of unfolded proteins exceeds the cellular regulatory capacity. The action for apoptosis is initiated by the activation of several proteases such as caspase-12, caspase-4, caspase-2, and caspase-9. Studies in rodents provided evidence supporting that caspase-12 is involved in ER stress-induced apoptosis. Caspase-12 is activated by IRE1 upon the insult of ER stress. Mice deficient in caspase-12 are resistant to ER stress-induced apoptosis, but remain susceptible to apoptosis induced by other stimuli [96]. There is evidence that caspase-12 can also be activated by interacting with TRAF2, a signaling molecule downstream of IRE1 [95]. In response to ER stress, caspase-7 is translocated from the cytosol to the ER surface, which then activates procaspase-12 as well [97]. The human caspase-4 is the closest paralog of rodent caspase-12, which is normally located on the ER membrane. However, caspase-4 can only be activated by ER stress-inducing reagents not by the other apoptotic reagents, and knockdown of caspase-4 by siRNA reduces ER stress-induced apoptosis in neuroblastoma cells [98]. Similarly, caspase-2 and caspase-9 are found to be activated in the early phase of ER stress and inhibition of their activation either by inhibitors or siRNA reduces ER stress-induced apoptosis [99]. Studies also suggest that some members of inhibitor of apoptosis protein family prevent ER stress-induced cell death via interacting with caspase-2 and caspase-9 [99].

Other than the implication of caspases, Ask1 kinase and CCAAT/enhancer binding protein (C/EBP) homologous protein (CHOP) are also critical mediators for ER stress-induced cell death. IRE1/TRAF2 complex recruits Ask1 and activates subsequent JNK signaling. Studies have shown that the activation of JNK inhibits antiapoptotic protein BCL-2 [100] and induces proapoptotic protein Bim [101, 102]. Loss of *Ask1* suppresses ER stress-induced JNK activation and provides protection for cells against ER stress-induced death [103]. CHOP is a transcription factor belonging to basic leucine zipper transcription factor (bZIP) family. Many inducers of UPR including ATF4, ATF6, and XBP-1 up-regulate CHOP expression, and phosphorylation of CHOP at ser78 and ser81 by p38 MAPK enhances its transcriptional activity [44, 104]. Upon its activation, CHOP suppresses anti-apoptotic protein BCL-2 which in turn induces cells undergoing apoptosis [105–107].

8. Conclusion and Future Directions

There is convincing evidence that ER stress plays an essential role in autoimmune-mediated β -cell destruction. Feasible evidence also supports that ER stress modulates autoimmune response during T1D development (Table 1). ER stress in β cells can be either triggered by autoimmune responses against β -cell self-antigens and/or by the increase of compensated insulin synthesis. During the course of type 1 diabetes, autoreactive immune cells secrete copious amount of inflammatory cytokines such as IL-1 β , TNF- α , and IFN- γ into the islet milieu, which stimulate excessive production of NO in β cells and mediate β -cell destruction by inducing ER stress. Recent studies further suggest that ER stress also modulates the functionality of immune cells with implications in autoimmune progression. The absolute insulin deficiency in T1D patients renders the residual β cells for compensated insulin secretion to meet the demands of insulin for maintaining blood glucose homeostasis. This increase in insulin biosynthesis could overwhelm the folding capacity of ER, leading to UPR and ER stress in β cells, which in turn exacerbates β -cell dysfunction and T1D onset.

It should be kept in mind that the mechanisms underlying autoimmune-mediated β -cell destruction in type 1 diabetes are complex, and ER stress is unlikely the exclusive mechanism implicated in disease process. Despite recent significant advancement in this field, there are still many questions yet to be addressed. Can ER stress be served as a biomarker for β -cell destruction and autoimmune progression in the clinic setting? Are there additional factors for induction of ER stress in β cells during T1D development? Does modulation of ER stress in immune cells attenuate autoimmune progression? Does blockade of ER stress protect β cells from autoimmune-mediated destruction? Future studies aimed at dissecting these questions would provide a broadened insight for T1D pathogenesis and would have great potential for developing novel therapeutic strategies against this devastating disorder.

TABLE 1: Publications relevant to ER stress in the regulation of immune response and β -cell destruction.

Author	Defective/mutant gene	Species	Major finding	Reference
Harding et al.	PERK ^{-/-}	Mouse	PERK-deficient mice are extremely susceptible to diabetes. They display a progressive β -cell loss and hyperglycemia with aging.	[49]
Delépine et al.	PERK ^{-/-}	Human	Deficiency of PERK in human results in Wolcott-Rallison syndrome, which is characterized by early infancy insulin-dependent diabetes and multisystemic dysfunction.	[50]
Scheuner et al.	eIF2 α mutant (Ser51Ala)	Mouse	Ser51Ala mutation of eIF2 α shows a deficiency in pancreatic β cells manifested by the smaller core of insulin-secreting β cells and attenuated insulin secretion, and the mice die from hypoglycemia at their early infancy.	[52]
Ron et al.	Ins2 mutation	Mouse	Ins2 mutation in Akita mice disrupts disulfide bond between the α and β chain of proinsulin, which leads to the mis-folding of the mutated insulin and further induces ER stress in β cells and diabetes.	[56]
Zhang et al.	IRE1 ^{-/-}	Mouse	Pro-B cells failed to differentiate into pre-B cells when deficient for IRE1.	[65]
Iwakoshi et al.	XBP-1 ^{-/-}	Mouse	Deficiency of XBP-1 results in the impacted development of both conventional and plasmacytoid DCs. Loss of XBP-1 renders DCs vulnerable to ER stress-induced apoptosis.	[72]
Goodall et al.	CHOP knockdown		Knockdown of CHOP suppressed the production of IL-23 induced by ER stress and TLR signaling.	[73]
Richardson et al.	XBP-1 mutation	<i>C. elegans</i>	Innate immune response induced by <i>P. aeruginosa</i> infection causes ER stress in <i>C. elegans</i> , and mutations with loss of function for XBP-1 lead to larval lethality.	[74]
Kaser et al.	XBP-1 polymorphisms	Human	Loss of XBP-1 in intestinal epithelial cells induces Paneth cell dysfunction and overactive epithelium, leading to impaired mucosal defense to <i>Listeria monocytogenes</i> and increased sensitivity to colitis, an inflammatory disease sharing similar properties with T1D. The polymorphisms within the XBP-1 gene are associated with Crohn's disease and ulcerative colitis in humans.	[75]
Nakagawa et al.	Caspase-12 ^{-/-}	Mouse	Caspase-12 is involved in ER stress-induced apoptosis. Mice deficient in caspase-12 are resistant to ER stress-induced apoptosis, but remain susceptible to apoptosis induced by other stimuli.	[96]
Hitomi et al.	Caspase-4 knockdown	Human	Human caspase-4, the closest paralog of rodent caspase-12, is involved in ER stress-induced apoptosis. Knockdown of caspase-4 by siRNA reduces ER stress-induced apoptosis.	[98]
Nishitoh et al.	Ask1 ^{-/-}	Mouse	Loss of Ask1 suppresses ER stress-induced JNK activation and protects cells from ER stress-induced death.	[103]

Abbreviations

AP1:	Activator protein 1	ER:	Endoplasmic reticulum
Ask1:	Apoptosis signal-regulating kinase 1	ER stress:	Endoplasmic reticulum stress
ATF6:	Activating transcription factor 6	ERAD:	ER-associated degradation
BCR:	B-cell receptor	ERSE:	ER stress enhancer
BLIMP1:	B-lymphocyte-induced maturation protein 1	HMGB1:	High-mobility group box 1
BSAP:	B-cell-lineage-specific activator protein	HSPs:	Heat shock proteins
bZIP:	Basic leucine zipper transcription factor	IKK:	I κ B kinase
C/EBP:	CCAAT/enhancer binding protein	iNOS:	Inducible nitric oxide synthase
CHOP:	C/EBP homologous protein	IRE1:	Inositol-requiring enzyme 1
CPA:	Cyclopiazonic acid	IRS-1:	Insulin receptor substrate-1
CREBH:	Cyclic-AMP-responsive-element-binding protein H	JIK:	c-Jun N-terminal inhibitory kinase
DC:	Dendritic cell	JNK:	c-Jun N-terminal kinases
DP5:	Death protein 5	NO:	Nitric oxygen
eIF2 α :	Eukaryotic initiation factor 2 α	NRF2:	Nuclear factor-erythroid-derived 2-related factor 2

PARP: Poly(ADP-ribose) polymerase
 PERK: Pancreatic endoplasmic reticulum kinase
 RER: Rough endoplasmic reticulum
 ROS: Reactive oxygen species
 SER: Smooth endoplasmic reticulum
 SERCA: Sarcoendoplasmic reticulum pump Ca^{2+} ATPase
 SR: Sarcoplasmic reticulum
 T1D: Type 1 diabetes
 TLR: Toll-like receptor
 TRAF2: TNF receptor-associated factor 2
 TRIF: Toll-IL-1R-containing adaptor inducing IFN- β
 UPR: Unfolded protein response
 UPRE: Unfolded protein response element
 XBP-1: X box protein-1.

Acknowledgments

Our research is supported by grants from the Juvenile Diabetes Research Foundation International (JDRFI), the EFSD/CDC/Lilly Program for Collaborative Diabetes Research between China and Europe, the Synergy Award from the Diabetes and Obesity Discovery Institute (DODI), and the National Natural Science Foundation of China (81130014/H0704 and 81101553/H1604). The authors declare that they have no financial conflict of interests.

References

- [1] D. M. Maahs, N. A. West, J. M. Lawrence, and E. J. Mayer-Davis, "Epidemiology of type 1 diabetes," *Endocrinology and Metabolism Clinics of North America*, vol. 39, no. 3, pp. 481–497, 2010.
- [2] P. Yang, H.-L. Li, and C.-Y. Wang, "FUT2 nonfunctional variant: a "missing link" between genes and environment in type 1 diabetes?" *Diabetes*, vol. 60, no. 11, pp. 2685–2687, 2011.
- [3] E. Bandurska-Stankiewicz and D. Wiatr, "Programme preventing vision loss due to diabetes," *Klinika Oczna*, vol. 109, no. 7–9, pp. 359–362, 2007.
- [4] N. Cheung and T. Y. Wong, "Diabetic retinopathy and systemic vascular complications," *Progress in Retinal and Eye Research*, vol. 27, no. 2, pp. 161–176, 2008.
- [5] S. Studholme, "Diabetic retinopathy," *Journal of Perioperative Practice*, vol. 18, no. 5, pp. 205–210, 2008.
- [6] V. Monhart, "Diabetes mellitus, hypertension and kidney," *Vnitřní Lekarství*, vol. 54, no. 5, pp. 499–507, 2008.
- [7] J. F. Navarro-González and C. Mora-Fernández, "The role of inflammatory cytokines in diabetic nephropathy," *Journal of the American Society of Nephrology*, vol. 19, no. 3, pp. 433–442, 2008.
- [8] A. J. Boulton, "Diabetic neuropathy: classification, measurement and treatment," *Current Opinion in Endocrinology, Diabetes and Obesity*, vol. 14, no. 2, pp. 141–145, 2007.
- [9] R. S. Cornell and I. Ducic, "Painful diabetic neuropathy," *Clinics in Podiatric Medicine and Surgery*, vol. 25, no. 3, pp. 347–360, 2008.
- [10] E. Otto-Buczowska, Z. Kazibutowska, J. Sołyk, and Ł. Machnica, "Neuropathy and type 1 diabetes mellitus," *Endokrynologia, Diabetologia i Choroby Przemiany Materii Wieku Rozwojowego*, vol. 14, no. 2, pp. 109–116, 2008.
- [11] S. E. Gardner and R. A. Frantz, "Wound bioburden and infection-related complications in diabetic foot ulcers," *Biological Research for Nursing*, vol. 10, no. 1, pp. 44–53, 2008.
- [12] D. Malgrange, "Physiopathology of the diabetic foot," *Revue de Medecine Interne*, vol. 29, supplement 2, pp. S231–S237, 2008.
- [13] O. Ochoa, F. M. Torres, and P. K. Shireman, "Chemokines and diabetic wound healing," *Vascular*, vol. 15, no. 6, pp. 350–355, 2007.
- [14] M. Anselmino, H. Gohlke, L. Mellbin, and L. Rydén, "Cardiovascular prevention in patients with diabetes and prediabetes," *Herz*, vol. 33, no. 3, pp. 170–177, 2008.
- [15] T. Inoguchi and R. Takayanagi, "Role of oxidative stress in diabetic vascular complications," *Fukuoka Igaku Zasshi*, vol. 99, no. 3, pp. 47–55, 2008.
- [16] T. H. Marwick, "Diabetic heart disease," *Postgraduate Medical Journal*, vol. 84, no. 990, pp. 188–192, 2008.
- [17] W. D'Hertog, M. Maris, G. B. Ferreira et al., "Novel insights into the global proteome responses of insulin-producing INS-1E cells to different degrees of endoplasmic reticulum stress," *Journal of Proteome Research*, vol. 9, no. 10, pp. 5142–5152, 2010.
- [18] S. C. Hubbard and R. J. Ivatt, "Synthesis and processing of asparagine-linked oligosaccharides," *Annual Review of Biochemistry*, vol. 50, pp. 555–583, 1981.
- [19] R. Kornfeld and S. Kornfeld, "Assembly of asparagine-linked oligosaccharides," *Annual Review of Biochemistry*, vol. 54, pp. 631–664, 1985.
- [20] S. W. Fewell, K. J. Travers, J. S. Weissman, and J. L. Brodsky, "The action of molecular chaperones in the early secretory pathway," *Annual Review of Genetics*, vol. 35, pp. 149–191, 2001.
- [21] F. Paltauf, S. D. Kohlwein, and S. A. Henry, "Regulation and compartmentalization of lipid synthesis in yeast," in *The Molecular and Cellular Biology of the Yeast Saccharomyces*, E. W. Jones and J. R. Broach, Eds., pp. 415–500, Cold Spring Harbor Laboratory Press, New York, NY, USA, 1992.
- [22] A. S. Lee, "The glucose-regulated proteins: stress induction and clinical applications," *Trends in Biochemical Sciences*, vol. 26, no. 8, pp. 504–510, 2001.
- [23] M. Schröder and P. Friedl, "Overexpression of recombinant human antithrombin III in Chinese hamster ovary cells results in malformation and decreased secretion of recombinant protein," *Biotechnology and Bioengineering*, vol. 53, no. 6, pp. 547–559, 1997.
- [24] M. Schröder, R. Schäfer, and P. Friedl, "Induction of protein aggregation in an early secretory compartment by elevation of expression level," *Biotechnology and Bioengineering*, vol. 78, no. 2, pp. 131–140, 2002.
- [25] A. J. Dorner, L. C. Wasley, and R. J. Kaufman, "Increased synthesis of secreted proteins induces expression of glucose-regulated proteins in butyrate-treated Chinese hamster ovary cells," *Journal of Biological Chemistry*, vol. 264, no. 34, pp. 20602–20607, 1989.
- [26] R. J. Kaufman, L. C. Wasley, and A. J. Dorner, "Synthesis, processing, and secretion of recombinant human factor VIII expressed in mammalian cells," *Journal of Biological Chemistry*, vol. 263, no. 13, pp. 6352–6362, 1988.
- [27] M. Schröder and R. J. Kaufman, "ER stress and the unfolded protein response," *Mutation Research*, vol. 569, no. 1–2, pp. 29–63, 2005.
- [28] A. M. Reimold, N. N. Iwakoshi, J. Manis et al., "Plasma cell differentiation requires the transcription factor XBP-1," *Nature*, vol. 412, no. 6844, pp. 300–307, 2001.

- [29] N. N. Iwakoshi, A. H. Lee, P. Vallabhajosyula, K. L. Otipoby, K. Rajewsky, and L. H. Glimcher, "Plasma cell differentiation and the unfolded protein response intersect at the transcription factor XBP-1," *Nature Immunology*, vol. 4, no. 4, pp. 321–329, 2003.
- [30] I. M. Martínez and M. J. Chrispeels, "Genomic analysis of the unfolded protein response in Arabidopsis shows its connection to important cellular processes," *Plant Cell*, vol. 15, no. 2, pp. 561–576, 2003.
- [31] T. M. Pakula, M. Laxell, A. Huuskonen, J. Uusitalo, M. Saloheimo, and M. Penttilä, "The effects of drugs inhibiting protein secretion in the filamentous fungus *Trichoderma reesei*. Evidence for down-regulation of genes that encode secreted proteins in the stressed cells," *Journal of Biological Chemistry*, vol. 278, no. 45, pp. 45011–45020, 2003.
- [32] H. P. Harding, Y. Zhang, and D. Ron, "Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase," *Nature*, vol. 397, no. 6716, pp. 271–274, 1999.
- [33] R. Casagrande, P. Stern, M. Diehn et al., "Degradation of proteins from the ER of *S. cerevisiae* requires an intact unfolded protein response pathway," *Molecular Cell*, vol. 5, no. 4, pp. 729–735, 2000.
- [34] R. Friedlander, E. Jarosch, J. Urban, C. Volkwein, and T. Sommer, "A regulatory link between ER-associated protein degradation and the unfolded-protein response," *Nature Cell Biology*, vol. 2, no. 7, pp. 379–384, 2000.
- [35] K. J. Travers, C. K. Patil, L. Wodicka, D. J. Lockhart, J. S. Weissman, and P. Walter, "Functional and genomic analyses reveal an essential coordination between the unfolded protein response and ER-associated degradation," *Cell*, vol. 101, no. 3, pp. 249–258, 2000.
- [36] R. J. Kaufman, "Orchestrating the unfolded protein response in health and disease," *Journal of Clinical Investigation*, vol. 110, no. 10, pp. 1389–1398, 2002.
- [37] F. J. Stevens and Y. Argon, "Protein folding in the ER," *Seminars in Cell and Developmental Biology*, vol. 10, no. 5, pp. 443–454, 1999.
- [38] M. Schröder, "Endoplasmic reticulum stress responses," *Cellular and Molecular Life Sciences*, vol. 65, no. 6, pp. 862–894, 2008.
- [39] C. K. Suzuki, J. S. Bonifacino, A. Y. Lin, M. M. Davis, and R. D. Klausner, "Regulating the retention of T-cell receptor α chain variants within the endoplasmic reticulum: Ca^{2+} -dependent association with BiP," *Journal of Cell Biology*, vol. 114, no. 2, pp. 189–205, 1991.
- [40] L. J. Li, X. Li, A. Ferrario et al., "Establishment of a Chinese hamster ovary cell line that expresses grp78 antisense transcripts and suppresses A23187 induction of both GRP78 and GRP94," *Journal of Cellular Physiology*, vol. 153, no. 3, pp. 575–582, 1992.
- [41] E. F. Corbett, K. Oikawa, P. Francois et al., " Ca^{2+} regulation of interactions between endoplasmic reticulum chaperones," *Journal of Biological Chemistry*, vol. 274, no. 10, pp. 6203–6211, 1999.
- [42] J. X. Zhang, I. Braakman, K. E. S. Matlack, and A. Helenius, "Quality control in the secretory pathway: the role of calreticulin, calnexin and BiP in the retention of glycoproteins with C-terminal truncations," *Molecular Biology of the Cell*, vol. 8, no. 10, pp. 1943–1954, 1997.
- [43] S. Orrenius, B. Zhivotovsky, and P. Nicotera, "Regulation of cell death: the calcium-apoptosis link," *Nature Reviews Molecular Cell Biology*, vol. 4, no. 7, pp. 552–565, 2003.
- [44] C. Xu, B. Bailly-Maitre, and J. C. Reed, "Endoplasmic reticulum stress: cell life and death decisions," *Journal of Clinical Investigation*, vol. 115, no. 10, pp. 2656–2664, 2005.
- [45] K. Lee, W. Tirasophon, X. Shen et al., "IRE1-mediated unconventional mRNA splicing and S2P-mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein response," *Genes and Development*, vol. 16, no. 4, pp. 452–466, 2002.
- [46] J. Ye, R. B. Rawson, R. Komuro et al., "ER stress induces cleavage of membrane-bound ATF6 by the same proteases that process SREBPs," *Molecular Cell*, vol. 6, no. 6, pp. 1355–1364, 2000.
- [47] D. L. Eizirik, A. K. Cardozo, and M. Cnop, "The role for endoplasmic reticulum stress in diabetes mellitus," *Endocrine Reviews*, vol. 29, no. 1, pp. 42–61, 2008.
- [48] D. R. Laybutt, A. M. Preston, M. C. Åkerfeldt et al., "Endoplasmic reticulum stress contributes to beta cell apoptosis in type 2 diabetes," *Diabetologia*, vol. 50, no. 4, pp. 752–763, 2007.
- [49] H. P. Harding, H. Zeng, Y. Zhang et al., "Diabetes mellitus and exocrine pancreatic dysfunction in *Perk*^{-/-} mice reveals a role for translational control in secretory cell survival," *Molecular Cell*, vol. 7, no. 6, pp. 1153–1163, 2001.
- [50] M. Delépine, M. Nicolino, T. Barrett, M. Golamaully, G. Mark Lathrop, and C. Julier, "EIF2AK3, encoding translation initiation factor 2- α kinase 3, is mutated in patients with Wolcott-Rallison syndrome," *Nature Genetics*, vol. 25, no. 4, pp. 406–409, 2000.
- [51] E. Araki, S. Oyadomari, and M. Mori, "Endoplasmic reticulum stress and diabetes mellitus," *Internal Medicine*, vol. 42, no. 1, pp. 7–14, 2003.
- [52] D. Scheuner, B. Song, E. McEwen et al., "Translational control is required for the unfolded protein response and in vivo glucose homeostasis," *Molecular Cell*, vol. 7, no. 6, pp. 1165–1176, 2001.
- [53] D. Mathis, L. Vence, and C. Benoist, " β -cell death during progression to diabetes," *Nature*, vol. 414, no. 6865, pp. 792–798, 2001.
- [54] S. Oyadomari and M. Mori, "Roles of CHOP/GADD153 in endoplasmic reticulum stress," *Cell Death and Differentiation*, vol. 11, no. 4, pp. 381–389, 2004.
- [55] S. Oyadomari, K. Takeda, M. Takiguchi et al., "Nitric oxide-induced apoptosis in pancreatic β cells is mediated by the endoplasmic reticulum stress pathway," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 98, no. 19, pp. 10845–10850, 2001.
- [56] D. Ron, "Proteotoxicity in the endoplasmic reticulum: lessons from the Akita diabetic mouse," *Journal of Clinical Investigation*, vol. 109, no. 4, pp. 443–445, 2002.
- [57] D. L. Eizirik, M. L. Colli, and F. Ortis, "The role of inflammation in insulinitis and β -cell loss in type 1 diabetes," *Nature Reviews Endocrinology*, vol. 5, no. 4, pp. 219–226, 2009.
- [58] J. Zhong, P. Yang, and C. Wang, "Environmental triggers and endogenous alarmins linking innate immunity to the pathogenesis of type 1 diabetes," in *Type 1 Diabetes Mellitus: Etiology, Diagnosis and Treatment*, pp. 177–206, Nova Science, 2011.
- [59] J. Han, J. Zhong, W. Wei et al., "Extracellular high-mobility group box 1 acts as an innate immune mediator to enhance autoimmune progression and diabetes onset in NOD mice," *Diabetes*, vol. 57, no. 8, pp. 2118–2127, 2008.
- [60] S. Zhang, J. Zhong, P. Yang, F. Gong, and C. Y. Wang, "HMGB1, an innate alarmin, in the pathogenesis of type 1 diabetes," *International Journal of Clinical and Experimental Pathology*, vol. 3, no. 1, pp. 24–38, 2010.
- [61] E. N. Gurzov, F. Ortis, D. A. Cunha et al., "Signaling by IL-1 β +IFN- γ and ER stress converge on DP5/Hrk activation:

- a novel mechanism for pancreatic β -cell apoptosis," *Cell Death and Differentiation*, vol. 16, no. 11, pp. 1539–1550, 2009.
- [62] A. K. Cardozo, F. Ortis, J. Storling et al., "Cytokines down-regulate the sarcoendoplasmic reticulum pump Ca^{2+} ATPase 2b and deplete endoplasmic reticulum Ca^{2+} , leading to induction of endoplasmic reticulum stress in pancreatic β -cells," *Diabetes*, vol. 54, no. 2, pp. 452–461, 2005.
- [63] D. T. Rutkowski and R. J. Kaufman, "A trip to the ER: coping with stress," *Trends in Cell Biology*, vol. 14, no. 1, pp. 20–28, 2004.
- [64] S. Baekkeskov, J. N. Nielsen, and B. Marnier, "Autoantibodies in newly diagnosed diabetic children immunoprecipitate human pancreatic islet cell proteins," *Nature*, vol. 298, no. 5870, pp. 167–169, 1982.
- [65] K. Zhang, H. N. Wong, B. Song, C. N. Miller, D. Scheuner, and R. J. Kaufman, "The unfolded protein response sensor IRE1 α is required at 2 distinct steps in B cell lymphopoiesis," *Journal of Clinical Investigation*, vol. 115, no. 2, pp. 268–281, 2005.
- [66] A. L. Shaffer, M. Shapiro-Shelef, N. N. Iwakoshi et al., "XBP1, downstream of Blimp-1, expands the secretory apparatus and other organelles, and increases protein synthesis in plasma cell differentiation," *Immunity*, vol. 21, no. 1, pp. 81–93, 2004.
- [67] H. Niu, B. H. Ye, and R. Dalla-Favera, "Antigen receptor signaling induces MAP kinase-mediated phosphorylation and degradation of the BCL-6 transcription factor," *Genes and Development*, vol. 12, no. 13, pp. 1953–1961, 1998.
- [68] K. I. Lin, C. Angelin-Duclos, T. C. Kuo, and K. Calame, "Blimp-1-dependent repression of Pax-5 is required for differentiation of B cells to immunoglobulin M-secreting plasma cells," *Molecular and Cellular Biology*, vol. 22, no. 13, pp. 4771–4780, 2002.
- [69] A. M. Reimold, P. D. Ponath, Y. S. Li et al., "Transcription factor B cell lineage-specific activator protein regulates the gene for human X-box binding protein 1," *Journal of Experimental Medicine*, vol. 183, no. 2, pp. 393–401, 1996.
- [70] L. Zhao and S. L. Ackerman, "Endoplasmic reticulum stress in health and disease," *Current Opinion in Cell Biology*, vol. 18, no. 4, pp. 444–452, 2006.
- [71] K. Zhang, X. Shen, J. Wu et al., "Endoplasmic reticulum stress activates cleavage of CREBH to induce a systemic inflammatory response," *Cell*, vol. 124, no. 3, pp. 587–599, 2006.
- [72] N. N. Iwakoshi, M. Pypaert, and L. H. Glimcher, "The transcription factor XBP-1 is essential for the development and survival of dendritic cells," *Journal of Experimental Medicine*, vol. 204, no. 10, pp. 2267–2275, 2007.
- [73] J. C. Goodall, C. Wu, Y. Zhang et al., "Endoplasmic reticulum stress-induced transcription factor, CHOP, is crucial for dendritic cell IL-23 expression," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 41, pp. 17698–17703, 2010.
- [74] C. E. Richardson, T. Kooistra, and D. H. Kim, "An essential role for XBP-1 in host protection against immune activation in *C. elegans*," *Nature*, vol. 463, no. 7284, pp. 1092–1095, 2010.
- [75] A. Kaser, A. H. Lee, A. Franke et al., "XBP1 links ER stress to intestinal inflammation and confers genetic risk for human inflammatory bowel disease," *Cell*, vol. 134, no. 5, pp. 743–756, 2008.
- [76] C. W. Woo, D. Cui, J. Arellano et al., "Adaptive suppression of the ATF4-CHOP branch of the unfolded protein response by toll-like receptor signalling," *Nature cell biology*, vol. 11, no. 12, pp. 1473–1480, 2009.
- [77] J. D. Malhotra and R. J. Kaufman, "Endoplasmic reticulum stress and oxidative stress: a vicious cycle or a double-edged sword?" *Antioxidants and Redox Signaling*, vol. 9, no. 12, pp. 2277–2293, 2007.
- [78] B. P. Tu and J. S. Weissman, "Oxidative protein folding in eukaryotes: mechanisms and consequences," *Journal of Cell Biology*, vol. 164, no. 3, pp. 341–346, 2004.
- [79] H. P. Harding, Y. Zhang, H. Zeng et al., "An integrated stress response regulates amino acid metabolism and resistance to oxidative stress," *Molecular Cell*, vol. 11, no. 3, pp. 619–633, 2003.
- [80] S. B. Cullinan, D. Zhang, M. Hannink, E. Arvisais, R. J. Kaufman, and J. A. Diehl, "Nrf2 is a direct PERK substrate and effector of PERK-dependent cell survival," *Molecular and Cellular Biology*, vol. 23, no. 20, pp. 7198–7209, 2003.
- [81] S. B. Cullinan and J. A. Diehl, "PERK-dependent activation of Nrf2 contributes to redox homeostasis and cell survival following endoplasmic reticulum stress," *Journal of Biological Chemistry*, vol. 279, no. 19, pp. 20108–20117, 2004.
- [82] P. Hu, Z. Han, A. D. Couvillon, R. J. Kaufman, and J. H. Exton, "Autocrine tumor necrosis factor alpha links endoplasmic reticulum stress to the membrane death receptor pathway through IRE1 α -mediated NF- κ B activation and down-regulation of TRAF2 expression," *Molecular and Cellular Biology*, vol. 26, no. 8, pp. 3071–3084, 2006.
- [83] R. J. Davis, "Signal transduction by the JNK group of MAP kinases," *Cell*, vol. 103, no. 2, pp. 239–252, 2000.
- [84] H. Yamazaki, N. Hiramatsu, K. Hayakawa et al., "Activation of the Akt-NF- κ B pathway by subtilase cytotoxin through the ATF6 branch of the unfolded protein response," *Journal of Immunology*, vol. 183, no. 2, pp. 1480–1487, 2009.
- [85] Q. Wang, H. Zhang, B. Zhao, and H. Fei, "IL-1 β caused pancreatic β -cells apoptosis is mediated in part by endoplasmic reticulum stress via the induction of endoplasmic reticulum Ca^{2+} release through the c-Jun N-terminal kinase pathway," *Molecular and Cellular Biochemistry*, vol. 324, no. 1–2, pp. 183–190, 2009.
- [86] U. K. Messmer and B. Brüne, "Nitric oxide-induced apoptosis: p53-dependent and p53-independent signalling pathways," *Biochemical Journal*, vol. 319, no. 1, pp. 299–305, 1996.
- [87] R. I. Viner, D. A. Ferrington, T. D. Williams, D. J. Bigelow, and C. Schöneich, "Protein modification during biological aging: selective tyrosine nitration of the SERCA2a isoform of the sarcoplasmic reticulum Ca^{2+} -ATPase in skeletal muscle," *Biochemical Journal*, vol. 340, no. 3, pp. 657–669, 1999.
- [88] K. Y. Xu, D. L. Huso, T. M. Dawson, D. S. Bredt, and L. C. Becker, "Nitric oxide synthase in cardiac sarcoplasmic reticulum," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 96, no. 2, pp. 657–662, 1999.
- [89] L. Xu, J. P. Eu, G. Meissner, and J. S. Stamler, "Activation of the cardiac calcium release channel (Ryanodine receptor) by poly-S-nitrosylation," *Science*, vol. 279, no. 5348, pp. 234–237, 1998.
- [90] H. Liu, R. C. Bowes III, B. Van De Water, C. Sillence, J. F. Nagelkerke, and J. L. Stevens, "Endoplasmic reticulum chaperones GRP78 and calreticulin prevent oxidative stress, Ca^{2+} disturbances, and cell death in renal epithelial cells," *Journal of Biological Chemistry*, vol. 272, no. 35, pp. 21751–21759, 1997.
- [91] H. P. Harding, Y. Zhang, A. Bertolotti, H. Zeng, and D. Ron, "Perk is essential for translational regulation and cell survival during the unfolded protein response," *Molecular Cell*, vol. 5, no. 5, pp. 897–904, 2000.

- [92] F. Urano, X. Wang, A. Bertolotti et al., "Coupling of stress in the ER to activation of JNK protein kinases by transmembrane protein kinase IRE1," *Science*, vol. 287, no. 5453, pp. 664–666, 2000.
- [93] C. Zheng, V. Kabaleeswaran, Y. Wang, G. Cheng, and H. Wu, "Crystal structures of the TRAF2: cIAP2 and the TRAF1: TRAF2: cIAP2 complexes: affinity, specificity, and regulation," *Molecular Cell*, vol. 38, no. 1, pp. 101–113, 2010.
- [94] H. Habelhah, S. Takahashi, S. G. Cho, T. Kadoya, T. Watanabe, and Z. Ronai, "Ubiquitination and translocation of TRAF2 is required for activation of JNK but not of p38 or NF- κ B," *EMBO Journal*, vol. 23, no. 2, pp. 322–332, 2004.
- [95] T. Yoneda, K. Imaizumi, K. Oono et al., "Activation of caspase-12, an endoplasmic reticulum (ER) resident caspase, through tumor necrosis factor receptor-associated factor 2-dependent mechanism in response to the ER stress," *Journal of Biological Chemistry*, vol. 276, no. 17, pp. 13935–13940, 2001.
- [96] T. Nakagawa, H. Zhu, N. Morishima et al., "Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid- β ," *Nature*, vol. 403, no. 6765, pp. 98–103, 2000.
- [97] R. V. Rao, E. Hermel, S. Castro-Obregon et al., "Coupling endoplasmic reticulum stress to the cell death program. Mechanism of caspase activation," *Journal of Biological Chemistry*, vol. 276, no. 36, pp. 33869–33874, 2001.
- [98] J. Hitomi, T. Katayama, Y. Eguchi et al., "Involvement of caspase-4 in endoplasmic reticulum stress-induced apoptosis and A β -induced cell death," *Journal of Cell Biology*, vol. 165, no. 3, pp. 347–356, 2004.
- [99] H. H. Cheung, N. Lynn Kelly, P. Liston, and R. G. Korneluk, "Involvement of caspase-2 and caspase-9 in endoplasmic reticulum stress-induced apoptosis: a role for the IAPs," *Experimental Cell Research*, vol. 312, no. 12, pp. 2347–2357, 2006.
- [100] K. Yamamoto, H. Ichijo, and S. J. Korsmeyer, "BCL-2 is phosphorylated and inactivated by an ASK1/Jun N-terminal protein kinase pathway normally activated at G₂/M," *Molecular and Cellular Biology*, vol. 19, no. 12, pp. 8469–8478, 1999.
- [101] G. V. Putcha, S. Le, S. Frank et al., "JNK-mediated BIM phosphorylation potentiates BAX-dependent apoptosis," *Neuron*, vol. 38, no. 6, pp. 899–914, 2003.
- [102] K. Lei and R. J. Davis, "JNK phosphorylation of Bim-related members of the Bcl2 family induces Bax-dependent apoptosis," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 5, pp. 2432–2437, 2003.
- [103] H. Nishitoh, A. Matsuzawa, K. Tobiume et al., "ASK1 is essential for endoplasmic reticulum stress-induced neuronal cell death triggered by expanded polyglutamine repeats," *Genes and Development*, vol. 16, no. 11, pp. 1345–1355, 2002.
- [104] X. Wang and D. Ron, "Stress-induced phosphorylation and activation of the transcription factor CHOP (GADD153) by p38 MAP kinase," *Science*, vol. 272, no. 5266, pp. 1347–1349, 1996.
- [105] K. D. McCullough, J. L. Martindale, L. O. Klotz, T. Y. Aw, and N. J. Holbrook, "Gadd153 sensitizes cells to endoplasmic reticulum stress by down-regulating Bcl2 and perturbing the cellular redox state," *Molecular and Cellular Biology*, vol. 21, no. 4, pp. 1249–1259, 2001.
- [106] M. Matsumoto, M. Minami, K. Takeda, Y. Sakao, and S. Akira, "Ectopic expression of CHOP (GADD153) induces apoptosis in M1 myeloblastic leukemia cells," *FEBS Letters*, vol. 395, no. 2-3, pp. 143–147, 1996.
- [107] H. Zinszner, M. Kuroda, X. Wang et al., "CHOP is implicated in programmed cell death in response to impaired function of the endoplasmic reticulum," *Genes and Development*, vol. 12, no. 7, pp. 982–995, 1998.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

