

Research Article

Repurposing Lesogaberan to Promote Human Islet Cell Survival and β -Cell Replication

Jide Tian, Hoa Dang, Angela Hu, Willem Xu, and Daniel L. Kaufman

Department of Molecular and Medical Pharmacology, University of California, Los Angeles, CA, USA

Correspondence should be addressed to Daniel L. Kaufman; dkaufman@mednet.ucla.edu

Received 12 May 2017; Accepted 26 July 2017; Published 5 September 2017

Academic Editor: Peter Thule

Copyright © 2017 Jide Tian et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The activation of β -cell's A- and B-type gamma-aminobutyric acid receptors (GABA_A-Rs and GABA_B-Rs) can promote their survival and replication, and the activation of α -cell GABA_A-Rs promotes their conversion into β -cells. However, GABA and the most clinically applicable GABA-R ligands may be suboptimal for the long-term treatment of diabetes due to their pharmacological properties or potential side-effects on the central nervous system (CNS). Lesogaberan (AZD3355) is a peripherally restricted high-affinity GABA_B-R-specific agonist, originally developed for the treatment of gastroesophageal reflux disease (GERD) that appears to be safe for human use. This study tested the hypothesis that lesogaberan could be repurposed to promote human islet cell survival and β -cell replication. Treatment with lesogaberan significantly enhanced replication of human islet cells *in vitro*, which was abrogated by a GABA_B-R antagonist. Immunohistochemical analysis of human islets that were grafted into immune-deficient mice revealed that oral treatment with lesogaberan promoted human β -cell replication and islet cell survival *in vivo* as effectively as GABA (which activates both GABA_A-Rs and GABA_B-Rs), perhaps because of its more favorable pharmacokinetics. Lesogaberan may be a promising drug candidate for clinical studies of diabetes intervention and islet transplantation.

1. Introduction

A major goal in diabetes research is to develop agents that can safely promote human β -cell survival and replication. Most mitogens and growth factors that have been shown to promote rodent β -cell replication fail to promote human β -cell replication (reviewed in [1, 2]). β -Cells have been long known to express the GABA synthetic enzyme glutamic acid decarboxylase (GAD), as well as GABA_A-Rs and GABA_B-Rs [3–7]. Although GABA_A-Rs and GABA_B-Rs share GABA as an agonist, these receptors are encoded by distinct gene families and their activation induces different pathways; GABA_A-Rs are fast-acting chloride channels and GABA_B-Rs are slow-acting G-protein-coupled receptors [8, 9]. Recently, GABA administration has been shown to protect rodent and human β -cells from apoptosis and to promote their replication both *in vitro* and *in vivo* [10–14]. This response is mediated by both GABA_A-R and GABA_B-Rs [10–12, 14, 15].

GABA-mediated enhancement of β -cell replication did not attenuate after five weeks of GABA treatment and led to an eventual increase in β -cell mass in a nonautoimmune context [14]. Notably, GABA treatment enhanced β -cell replication and survival in newly diabetic NOD mice [11, 16, 17], indicating that GABA-R activation can be beneficial in an autoimmune context even when little β -cell mass remains.

β -Cells express GABA_A-Rs and GABA_B-Rs [3, 4, 7, 18, 19]. α -Cells express GABA_A-Rs but may not express functional GABA_B-Rs, while PCR detects GABA_B-R transcripts in isolated α -cells (but not the α -cell line α -TC9), a GABA_B-R agonist failed to modulate any of the tested α -cell functions [3, 18–21]. We are unaware of any evidence of functional GABA_B-Rs on δ or PP islet cells. Very recently, long-term treatment with antimalarial drugs that target gephyrin (a protein that participates in GABA_A-R transport to the membrane), or treatment with GABA, was shown to promote islet α -cell transdifferentiation into β -cells [22, 23].

This conversion appears to be mediated by GABA_A-Rs [22, 23] and not the GABA_B-Rs that are the focus of this study.

GABA appears safe for human consumption [24] and its inability to pass through the blood-brain barrier (BBB) avoids potential CNS effects. GABA, however, has a relatively low affinity for its receptors, a fast off-rate (presumably so that neurons can quickly respond to the next stimulus), and a short half-life (about 20 minutes in blood [25]) and therefore may be pharmacologically suboptimal. In the clinic, there are many BBB-permeable drugs that can modulate neuronal GABA-Rs and ameliorate CNS disorders such as seizures, anxiety, and insomnia. Their CNS effects, however, raise concerns about their long-term use for the treatment of diabetes. It would be ideal if peripherally restricted GABA-R agonists that were safe for human use could be repurposed for treating diabetes.

Lesogaberan (AZD3355) is a peripherally restricted GABA_B-R-specific agonist that was developed for the treatment of GERD [26–28]. It has an EC₅₀ of 9 nM (compared to GABA's EC₅₀ of 160 nM) and a K_i of 5 nM (versus 110 nM for GABA) for GABA_B-Rs [26] and a half-life of about 11 hours in peripheral blood [29, 30]. While treatment of GERD patients with lesogaberan (oral 60–240 mg twice daily) for up to 28 days in several phase IIb clinical trials did not result in sufficient beneficial effects, there were no treatment-related serious adverse events, suggesting that lesogaberan may be safe for human use [27, 29, 31]. Here, we tested the potential of repurposing lesogaberan to promote human islet cell survival and β -cell replication. Our findings suggest that targeting GABA_B-Rs can promote human β -cell replication and islet cell survival to a similar extent as GABA.

2. Materials and Methods

2.1. Chemicals. Lesogaberan was supplied by AstraZeneca (London, UK). The development and structure of lesogaberan have been previously described [28]. Saclofen, streptozotocin (STZ), and 5-bromo-2-deoxyuridine (BrdU) were purchased from Sigma Aldrich.

2.2. Islet Cell Proliferation Assay. Fresh human islets were obtained from the Integrated Islet Distribution Program (IIDP). The islets (50–75 IEQ/well) were treated in triplicate with, or without, the indicated dosages of lesogaberan in CMRL medium (0.1% glucose, Gibco) containing 10% human AB-type sera (MP Biomedicals, Santa Ana, USA) and 1.5 μ Ci/ml of ³H-thymidine in the presence or absence of the competitive GABA_B-R antagonist saclofen (10⁻⁴ M) for 4 days. Islets cultured in medium alone served as controls. The ³H-thymidine uptake in individual wells was measured by β -counter. Data were analyzed by the proliferation index formula: CPM of experimental wells / CPM of controls.

2.3. Analysis of Human β -Cell Replication In Vivo. All animal experiments were approved by UCLA's Animal Research Committee. NOD/scid mice were injected intraperitoneally with STZ to induce diabetes and implanted with about 2000 human islets under their kidney capsule. The mice were

randomized and treated for 12 days with plain water containing 0.8 mg/ml of BrdU as the control or water containing the same dose of BrdU and lesogaberan (0.08 mg/ml) or positive control GABA (6 mg/ml). At the end of treatment, the percentages of BrdU⁺insulin⁺ and Ki67⁺insulin⁺ β -cells in at least 2000 islet cells of 10 fields (magnification \times 400) of each islet graft were determined by immunofluorescence in a blinded manner, as our previous report [12].

2.4. Analysis of Human β -Cell Apoptosis In Vivo. STZ-rendered diabetic NOD/scid mice were implanted with about 2000–3000 human islets under their kidney capsule. The mice were randomized and given plain water or water containing lesogaberan (0.08 mg/ml) or positive control GABA (6 mg/ml). Forty-eight hours later, the percentages of insulin⁺ β -cells or TUNEL⁺ apoptotic islet cells in total islet cells within the grafts of individual recipients were determined by immunofluorescence in a blinded manner, as in our previous report [12].

2.5. Statistical Analysis. Data are expressed as the mean \pm SEM of individual groups ($n = 4–9$ mice per group) from two separate experiments. The difference among groups was analyzed by ANOVA and post hoc Fisher's least significant difference and the difference between groups was determined Student *t*-test. A *p* value of <0.05 was considered statistically significant.

3. Results

3.1. Lesogaberan Enhances Human Islet Cell Proliferation In Vitro. Human islets were treated with lesogaberan over a dose range in the presence, or absence, of the GABA_B-R competitive antagonist saclofen. The effect of drug treatment on islet cell proliferation was determined by ³H-thymidine incorporation. We observed that lesogaberan at 3 nM had a small but nonsignificant promitotic effect, while treatment at higher dosages (10 and 30 nM) led to a 2–3-fold increase in proliferation relative to that of islets cultured in medium alone (Figure 1). Lesogaberan effects were not dose-dependent at the concentrations tested, and this may reflect its bimodal effect that was noted in some early preclinical GERD studies [26], but was not observed in subsequent clinical studies [27, 29, 31–33]. The promitotic effect of lesogaberan was blocked by saclofen (Figure 1). Hence, activation of GABA_B-Rs enhanced human islet cell proliferation *in vitro*.

Since β -cells may be the only islet cell type that expresses functional GABA_B-Rs and α -cell transdifferentiation would require long-term GABA_A-R activation and would not involve ³H-thymidine incorporation into new DNA, the vast majority of the proliferating islet cells are likely to be β -cells. That contention, however, requires verification, which we pursued using quantitative immunohistochemical analysis of β -cells in human xenografts below.

3.2. Oral Lesogaberan Promotes Human β -Cell Replication In Vivo. Next, we quantitatively assessed lesogaberan's effects on human β -cell replication *in vivo*. We implanted \sim 2000 human islets under the kidney capsule of STZ-rendered diabetic NOD/scid mice. Islet recipients were provided with




FIGURE 1: Lesogaberan enhances human islet cell proliferation *in vitro*. Human islets were treated in triplicate with, or without, the indicated dosages of lesogaberan in the presence (black bars) or absence (open bars) of saclofen (10^{-4} M) for 4 days. Data shown are the average rate of proliferation \pm SEM relative to that of cultures with medium alone (designated as 1) using islets from two donors, each of which were studied in separate experiments. Treatment with saclofen alone did not affect the proliferation of human islet in our experimental system (data not shown). ** $p < 0.01$ versus lesogaberan.

water containing BrdU with, or without, lesogaberan (0.08 mg/ml) or GABA (6 mg/ml) for 12 days. The lesogaberan dose was chosen based on the results of previous preclinical studies in GERD models [26, 34] and our pilot dosing studies. All islet recipients, including controls given plain water, quickly became normoglycemic after receiving the islet graft and maintained normoglycemia throughout the experimental period. There was no significant difference in their food and water consumption (~ 4 ml/day), as well as body weights among these groups of mice (data not shown).

Immunofluorescent staining indicated that treatment with GABA significantly enhanced human β -cell replication (Figure 2), similar to previous observations [14, 16]. Lesogaberan treatment also significantly increased the frequency of BrdU⁺insulin⁺ and Ki67⁺insulin⁺ islet β -cells relative to that in human islets from the control mice (Figure 2). There was no significant difference in the percentages of replicative β -cells between the lesogaberan-treated and GABA-treated groups of mice. Hence, activation of peripheral GABA_B-Rs effectively promotes human islet β -cell replication *in vivo*.

3.3. Oral Lesogaberan Limits Human Islet Cell Apoptosis following Islet Transplantation. Following human islet transplantation, the islets are subjected to hypoxic, metabolic, and inflammatory stressors which cause the apoptosis of a large proportion of islet cells within the first few days after implantation [35, 36]. We examined the ability of lesogaberan administration to limit islet cell apoptosis in a human islet xenograft

model. We implanted STZ-rendered hyperglycemic NOD/scid mice with human islets under their kidney capsule and placed them on plain water or water containing lesogaberan or positive control GABA. Two days later, their kidneys were removed and tissue sections were stained by TUNEL and anti-insulin antibodies (Figure 3). In comparison with mice given plain water, treatment with GABA significantly protected human islet cells from apoptosis, consistent with previous observations [14, 16]. Furthermore, oral feeding with lesogaberan significantly reduced the percentages of apoptotic islet cells (Figure 3(b)) and increased the frequency of insulin⁺ β -cells in human islet grafts (Figure 3(c)). Thus, lesogaberan-mediated activation of GABA_B-Rs limits islet cell apoptosis in transplanted human islets, consistent with previous islet cell survival studies with the prototypic GABA_B-R-specific agonist baclofen, a drug unsuited for long-term human use [10, 12, 14].

4. Discussion

The modulation of GABA-Rs on β -cells is emerging as a new strategy to help promote human β -cell survival and replication in the context of T1D. Although GABA is safe for human consumption, its pharmacokinetics may be suboptimal. There are a number of available drugs that have been in wide clinical use to modulate GABA-Rs on CNS neurons; however, their CNS effects raise concerns for long-term diabetes treatment. We chose to test lesogaberan's potential for diabetes treatment based on (1) its high affinity for GABA_B-Rs, (2) its peripheral restriction circumvents CNS effects, and (3) its apparent safety in early clinical studies.

Our initial studies showed that lesogaberan at low dosages (30 and 10 nM) increased human islet cell proliferation *in vitro* by about 2-3-fold (resp.). This promitotic effect was abrogated by a GABA_B-R antagonist, confirming that the effect was mediated through GABA_B-Rs. The proliferating islet cells are likely to be primarily β -cells because β -cells may be the only islet cell type that express functional GABA_B-Rs and α -cell transdifferentiation requires long-term GABA_A-R activation [22].

After implanting human islets into scid mice, lesogaberan significantly increased β -cell replication in the islet grafts. The level of lesogaberan-induced β -cell replication *in vivo* was similar to that induced by GABA at the dosages used in our model and similar to the maximum level of β -cell replication that takes place shortly after birth in humans [37]. The activation of β -cell GABA_B-Rs causes the opening of potassium channels, release of Ca²⁺ from intracellular storage, PKA activation, and Ca²⁺-dependent activation of PI3K-Akt and CREB (reviewed in [14]). It is notable that lesogaberan promoted β -cell replication as effectively as GABA (which activates both GABA_B-Rs and GABA_A-Rs) which may be due to its superior pharmacokinetics. Our short-term treatments with lesogaberan and GABA should not have induced α -cell transdifferentiation (which became detectable after 2 months of GABA_A-R activation [22, 23]), and the small increase they induced in β -cell replication should not alter islet size. Finally, we observed that




FIGURE 2: Oral lesogaberan promotes human islet β -cell replication in mice. Mildly hyperglycemic NOD/scid mice were transplanted with human islets under their kidney capsule. The mice were randomized and provided with water containing with BrdU, with or without GABA (6 mg/ml) or lesogaberan (0.08 mg/ml) for 12 days. The percentages of replicated β -cells were determined by immunofluorescent assays using anti-insulin and anti-BrdU or anti-Ki67, followed by counterstaining with DAPI. (a) Representative image of islet cells (magnification $\times 400$) costained with anti-insulin (green) and anti-BrdU (red) (arrows). (b) Representative image of islet cells costained with anti-insulin (green) and anti-Ki67 (red) (arrows). (c) Graphical representation of the percentages of BrdU⁺insulin⁺ β -cells and (d) Ki67⁺insulin⁺ islet cells in total insulin⁺ β -cells. Data are mean \pm SEM from two independent experiments, each using islets from a human donor that were implanted into 4–9 NOD/scid mice. The percentages of BrdU⁺insulin⁺ and Ki67⁺insulin⁺ β -cells in at least 2000 islet cells of 10 fields of each islet graft were determined as described in Materials and Methods. * $p < 0.05$, ** $p < 0.01$ versus the control.

lesogaberan treatment significantly preserved human islet cells from apoptosis in human islet grafts.

It is thought that the amount of residual β -cell mass following T1D onset is a major factor determining the success of interventional therapy such that even a short-term treatment with lesogaberan may help preserve residual β -cell mass and thereby improve the outcome of interventional therapies. Indeed, GABA treatment enhanced β -cell replication and survival in newly diabetic NOD mice [11, 16], demonstrating that GABA-R activation can be beneficial in an autoimmune context even when there is little residual β -cell mass. It is worth noting that using FDA guidelines for scaling mouse doses to the equivalent human dosage (<http://www.fda.gov/downloads/Drugs/.../Guidances/UCM078932.pdf>), the lesogaberan dose we used was about 2- to 7-fold lower than the doses used in phase IIb GERD clinical trials [27, 32, 33]. This suggests that it may be possible to use lower dosage lesogaberan for diabetes treatment.

Conceivably, long-term lesogaberan treatment may lead to increased β -cell mass in T1D patients if autoimmune responses can be sufficiently controlled. In regard to

controlling β -cell autoimmunity, recent studies indicate that activation of GABA_B-Rs on immune cells has anti-inflammatory effects and can ameliorate collagen-induced arthritis and contact dermatitis in mouse models [38, 39]. Therefore, lesogaberan may also have anti-inflammatory effects that help control the pathogenic autoreactive T-cell responses that mediate β -cell destruction in T1D. It will be of interest to further investigate how activation of GABA_B-Rs modulates the functions of different types of immunocompetent cells, and this may be a fertile area for new anti-inflammatory drug development. Additionally, lesogaberan may be combined with other immune modulators to potentially increase therapeutic effects, as was shown by the synergistic effect of combining GABA and antigen-specific immunotherapy to reverse hyperglycemia in newly diabetic NOD mice [16].

5. Conclusions

We found that the GABA_B-R agonist lesogaberan promoted human islet cell proliferation *in vitro*, as well as β -cell




FIGURE 3: Oral lesogaberan protects human islet β -cells from apoptosis in islet grafts in mice. Diabetic NOD/scid mice were implanted with human islets and treated with plain water or water containing lesogaberan (0.08 mg/ml) or GABA (6 mg/ml) for 48 h. The percentages of apoptotic islet cells and islet β -cells in total human islet cells were determined by TUNEL assay and costaining with anti-insulin as well as DAPI. At least 2000 human islet cells in 10 fields (magnification $\times 400$) from individual grafts were counted. Data are representative image or expressed as the mean $\% \pm$ SEM for each group of mice ($n = 5 - 7$) from three separate experiments. (a) A representative image with white arrows indicating TUNEL⁺ cells (green for TUNEL⁺, red for anti-insulin⁺, and light blue for DAPI staining). (b) Percent apoptotic islet cells and (c) percent insulin⁺ islet cells. ** $p < 0.01$ versus the control.

replication and islet cell survival *in vivo*, as effectively as GABA. Accordingly, GABA_B-R agonists may provide a new drug class to help maintain residual β -cell mass and function after diabetes onset. Our findings also suggest that including lesogaberan in the drug regimen following clinical human islet transplantation for even a brief period may reduce β -cell loss due to stressors and may thereby reduce the number of islets required to achieve insulin independence. Lesogaberan's apparent safety and pharmacokinetic profile make it an excellent candidate for testing in clinical trials.

Conflicts of Interest

The authors declare that there is no duality of interest associated with this manuscript.

Acknowledgments

The authors thank the IIDP for providing isolated human islets. This work was supported by a grant from the JDRF (17-2013-403) and the National Institutes of Health (DK092480) to Daniel L. Kaufman.

References

- [1] S. Bonner-Weir, W. C. Li, L. Ouziel-Yahalom, L. Guo, G. C. Weir, and A. Sharma, " β -Cell growth and regeneration: replication is only part of the story," *Diabetes*, vol. 59, no. 10, pp. 2340–2348, 2010.
- [2] R. N. Kulkarni, E. B. Mizrahi, A. G. Ocana, and A. F. Stewart, "Human β -cell proliferation and intracellular signaling: driving in the dark without a road map," *Diabetes*, vol. 61, no. 9, pp. 2205–2213, 2012.
- [3] P. Rorsman, P. O. Berggren, K. Bokvist et al., "Glucose-inhibition of glucagon secretion involves activation of GABAA-receptor chloride channels," *Nature*, vol. 341, no. 6239, pp. 233–236, 1989.
- [4] M. Braun, R. Ramracheya, M. Bengtsson et al., " γ -Aminobutyric acid (GABA) is an autocrine excitatory transmitter in human pancreatic β -cells," *Diabetes*, vol. 59, no. 7, pp. 1694–1701, 2010.
- [5] A. Reetz, M. Solimena, M. Matteoli, F. Folli, K. Takei, and P. De Camilli, "GABA and pancreatic beta-cells: colocalization of glutamic acid decarboxylase (GAD) and GABA with synaptic-like microvesicles suggests their role in GABA storage and secretion," *The EMBO Journal*, vol. 10, no. 5, pp. 1275–1284, 1991.

- [6] M. Braun, A. Wendt, B. Birnir et al., "Regulated exocytosis of GABA-containing synaptic-like microvesicles in pancreatic β -cells," *The Journal of General Physiology*, vol. 123, no. 3, pp. 191–204, 2004.
- [7] J. Taneera, Z. Jin, Y. Jin et al., " γ -Aminobutyric acid (GABA) signalling in human pancreatic islets is altered in type 2 diabetes," *Diabetologia*, vol. 55, no. 7, pp. 1985–1994, 2012.
- [8] R. W. Olsen and A. J. Tobin, "Molecular biology of GABAA receptors," *The FASEB Journal*, vol. 4, no. 5, pp. 1469–1480, 1990.
- [9] B. Bettler, K. Kaupmann, J. Mosbacher, and M. Gassmann, "Molecular structure and physiological functions of GABA(B) receptors," *Physiological Reviews*, vol. 84, no. 3, pp. 835–867, 2004.
- [10] B. Ligon, J. Yang, S. B. Morin, M. F. Ruberti, and M. L. Steer, "Regulation of pancreatic islet cell survival and replication by γ -aminobutyric acid," *Diabetologia*, vol. 50, no. 4, pp. 764–773, 2007.
- [11] N. Soltani, H. Qiu, M. Aleksic et al., "GABA exerts protective and regenerative effects on islet beta cells and reverses diabetes," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 108, no. 28, pp. 11692–11697, 2011.
- [12] J. Tian, H. Dang, Z. Chen et al., " γ -Aminobutyric acid regulates both the survival and replication of human β -cells," *Diabetes*, vol. 62, no. 11, pp. 3760–3765, 2013.
- [13] G. J. Prud'homme, Y. Glinka, C. Hasilo, S. Paraskevas, X. Li, and Q. Wang, "GABA protects human islet cells against the deleterious effects of immunosuppressive drugs and exerts immunoinhibitory effects alone," *Transplantation*, vol. 96, no. 7, pp. 616–623, 2013.
- [14] I. Purwana, J. Zheng, X. Li et al., "GABA promotes human β -cell proliferation and modulates glucose homeostasis," *Diabetes*, vol. 63, no. 12, pp. 4197–4205, 2014.
- [15] J. Tian, H. Dang, B. Middleton, and D. L. Kaufman, "Clinically applicable GABA receptor positive allosteric modulators promote β -cell replication," *Scientific Reports*, vol. 7, no. 1, p. 374, 2017.
- [16] J. Tian, H. Dang, A. V. Nguyen, Z. Chen, and D. L. Kaufman, "Combined therapy with GABA and proinsulin/alum acts synergistically to restore long-term normoglycemia by modulating T-cell autoimmunity and promoting β -cell replication in newly diabetic NOD mice," *Diabetes*, vol. 63, no. 9, pp. 3128–3134, 2014.
- [17] J. Tian, H. Dang, and D. L. Kaufman, "Combining antigen-based therapy with GABA treatment synergistically prolongs survival of transplanted β -cells in diabetic NOD mice," *PLoS One*, vol. 6, no. 9, article e25337, 2011.
- [18] M. Braun, A. Wendt, K. Buschard et al., "GABAB receptor activation inhibits exocytosis in rat pancreatic β -cells by G-protein-dependent activation of calcineurin," *The Journal of Physiology*, vol. 559, Part 2, pp. 397–409, 2004.
- [19] A. Wendt, B. Birnir, K. Buschard et al., "Glucose inhibition of glucagon secretion from rat alpha-cells is mediated by GABA released from neighboring β -cells," *Diabetes*, vol. 53, no. 4, pp. 1038–1045, 2004.
- [20] N. L. Brice, A. Varadi, S. J. Ashcroft, and E. Molnar, "Metabotropic glutamate and GABA(B) receptors contribute to the modulation of glucose-stimulated insulin secretion in pancreatic beta cells," *Diabetologia*, vol. 45, no. 2, pp. 242–252, 2002.
- [21] S. J. Bailey, M. A. Ravier, and G. A. Rutter, "Glucose-dependent regulation of γ -aminobutyric acid (GABA_A) receptor expression in mouse pancreatic islet α -cells," *Diabetes*, vol. 56, no. 2, pp. 320–327, 2007.
- [22] N. Ben-Othman, A. Vieira, M. Courtney et al., "Long-term GABA administration induces alpha cell-mediated beta-like cell neogenesis," *Cell*, pp. 31523–31529, 2016.
- [23] J. Li, T. Casteels, T. Frogne et al., "Artemisinins target GABAA receptor signaling and impair alpha cell identity," *Cell*, vol. 168, no. 1-2, article e115, pp. 86–100, 2017.
- [24] J. Li, Z. Zhang, X. Liu et al., "Study of GABA in healthy volunteers: pharmacokinetics and pharmacodynamics," *Frontiers in Pharmacology*, vol. 6, p. 260, 2015.
- [25] D. B. Tower and E. Roberts, Eds., *Inhibition in the nervous system and GABA*, pp. 562–578, Pergamon Press, New York, 1960.
- [26] A. Lehmann, M. Antonsson, A. A. Holmberg et al., "(R)-(3-Amino-2-fluoropropyl) phosphinic acid (AZD3355), a novel GABAB receptor agonist, inhibits transient lower esophageal sphincter relaxation through a peripheral mode of action," *The Journal of Pharmacology and Experimental Therapeutics*, vol. 331, no. 2, pp. 504–512, 2009.
- [27] N. J. Shaheen, H. Denison, K. Bjorck, M. Karlsson, and D. G. Silberg, "Efficacy and safety of lesogaberan in gastro-oesophageal reflux disease: a randomised controlled trial," *Gut*, vol. 62, no. 9, pp. 1248–1255, 2013.
- [28] C. Alstermark, K. Amin, S. R. Dinn et al., "Synthesis and pharmacological evaluation of novel gamma-aminobutyric acid type B (GABAB) receptor agonists as gastroesophageal reflux inhibitors," *Journal of Medicinal Chemistry*, vol. 51, no. 14, pp. 4315–4320, 2008.
- [29] M. Niazi, S. Skrtic, M. Ruth, and A. A. Holmberg, "Pharmacokinetic profile of lesogaberan (AZD3355) in healthy subjects: a novel GABA(B)-receptor agonist reflux inhibitor," *Drugs in R&D*, vol. 11, no. 1, pp. 77–83, 2011.
- [30] B. Fransson, D. G. Silberg, M. Niazi, F. Miller, M. Ruth, and A. A. Holmberg, "Effect of food on the bioavailability of lesogaberan given as an oral solution or as modified-release capsules in healthy male volunteers," *International Journal of Clinical Pharmacology and Therapeutics*, vol. 50, no. 4, pp. 307–314, 2012.
- [31] G. E. Boeckxstaens, H. Beaumont, V. Mertens et al., "Effects of lesogaberan on reflux and lower esophageal sphincter function in patients with gastroesophageal reflux disease," *Gastroenterology*, vol. 139, no. 2, pp. 409–417, 2010.
- [32] G. E. Boeckxstaens, H. Beaumont, J. G. Hatlebakk et al., "A novel reflux inhibitor lesogaberan (AZD3355) as add-on treatment in patients with GORD with persistent reflux symptoms despite proton pump inhibitor therapy: a randomised placebo-controlled trial," *Gut*, vol. 60, no. 9, pp. 1182–1188, 2011.
- [33] M. Niazi, D. G. Silberg, F. Miller, M. Ruth, and A. A. Holmberg, "Evaluation of the pharmacokinetic interaction between lesogaberan (AZD3355) and esomeprazole in healthy subjects," *Drugs in R&D*, vol. 10, no. 4, pp. 243–251, 2010.
- [34] L. Branden, A. Fredriksson, E. Harring, J. Jensen, and A. Lehmann, "The novel, peripherally restricted GABAB receptor agonist lesogaberan (AZD3355) inhibits acid reflux and reduces esophageal acid exposure as measured with 24-h pHmetry in dogs," *European Journal of Pharmacology*, vol. 634, no. 1–3, pp. 138–141, 2010.
- [35] J. A. Emamullee and A. M. Shapiro, "Factors influencing the loss of beta-cell mass in islet transplantation," *Cell Transplantation*, vol. 16, no. 1, pp. 1–8, 2007.

- [36] A. M. Davalli, Y. Ogawa, C. Ricordi, D. W. Scharp, S. Bonner-Weir, and G. C. Weir, "A selective decrease in the beta cell mass of human islets transplanted into diabetic nude mice," *Transplantation*, vol. 59, no. 6, pp. 817–820, 1995.
- [37] B. E. Gregg, P. C. Moore, D. Demozay et al., "Formation of a human β -cell population within pancreatic islets is set early in life," *The Journal of Clinical Endocrinology and Metabolism*, vol. 97, no. 9, pp. 3197–3206, 2012.
- [38] S. Huang, J. Mao, B. Wei, and G. Pei, "The anti-spasticity drug baclofen alleviates collagen-induced arthritis and regulates dendritic cells," *Journal of Cellular Physiology*, vol. 230, no. 7, pp. 1438–1447, 2015.
- [39] B. Duthey, A. Hubner, S. Diehl, S. Boehncke, J. Pfeffer, and W. H. Boehncke, "Anti-inflammatory effects of the GABA(B) receptor agonist baclofen in allergic contact dermatitis," *Experimental Dermatology*, vol. 19, no. 7, pp. 661–666, 2010.



Hindawi
Submit your manuscripts at
<https://www.hindawi.com>

