

Research Article

A Pilot Study of Plantar Mechanics Distributions and Fatigue Profiles after Running on a Treadmill: Using a Support Vector Machine Algorithm

Qian Liu,¹ Hairong Chen,¹ Anand Thirupathi,¹ Meimei Yang ,^{2,3} Julien S. Baker ,⁴ and Yaodong Gu ^{1,5}

¹Faculty of Sports Science, Ningbo University, Ningbo 315211, China

²Department of International Office, Ningbo University, Ningbo 315211, China

³CEEC Economic and Trade Cooperation Institute, Ningbo University, Ningbo 315211, China

⁴Centre for Health and Exercise Science Research, Department of Sport, Physical Education and Health, Hong Kong Baptist University, Hong Kong 999077, China

⁵Faculty of Engineering, University of Szeged, Szeged 6724, Hungary

Correspondence should be addressed to Meimei Yang; yangmeimei@nbu.edu.cn and Yaodong Gu; guyaodong@nbu.edu.cn

Received 26 July 2022; Revised 6 October 2022; Accepted 12 October 2022; Published 21 February 2023

Academic Editor: Xian-Bo Wang

Copyright © 2023 Qian Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The treadmill is widely used in running fatigue experiments, and the variation of plantar mechanical parameters caused by fatigue and gender, as well as the prediction of fatigue curves by a machine learning algorithm, play an important role in providing different training programs. This experiment aimed to compare changes in peak pressure (PP), peak force (PF), plantar impulse (PI), and gender differences of novice runners after they were fatigued by running. A support vector machine (SVM) was used to predict the fatigue curve according to the changes in PP, PF, and PI before and after fatigue. 15 healthy males and 15 healthy females completed two runs at a speed of $3.3 \text{ m/s} \pm 5\%$ on a footscan pressure plate before and after fatigue. After fatigue, PP, PF, and PI decreased at hallux ($T1$) and second-fifth toes ($T2-5$), while heel medial (HM) and heel lateral (HL) increased. In addition, PP and PI also increased at the first metatarsal ($M1$). PP, PF, and PI at $T1$ and $T2-5$ were significantly higher in females than in males, and metatarsals 3–5 ($M3-5$) were significantly lower in females than in males. The SVM classification algorithm results showed the accuracy was above average level using the $T1$ PP/HL PF (train accuracy: 65%; test accuracy: 75%), $T1$ PF/HL PF (train accuracy: 67.5%; test accuracy: 65%), and HL PF/ $T1$ PI (train accuracy: 67.5%; test accuracy: 70%). These values could provide information about running and gender-related injuries, such as metatarsal stress fractures and hallux valgus. Application of the SVM to the identification of plantar mechanical features before and after fatigue. The features of the plantar zones after fatigue can be identified and the learned algorithm of plantar zone combinations with above-average accuracy ($T1$ PP/HL PF, $T1$ PF/HL PF, and HL PF/ $T1$ PI) can be used to predict running fatigue and supervise training. It provided an important idea for the detection of fatigue after running.

1. Introduction

The most serious threat to health in modern times has been identified as sedentary behavior with insufficient physical activity [1]. Running has long been a popular leisure activity. Athletes have much lower resting heart rate, body weight, body mass index (BMI), and triglyceride levels compared to the general population [2], indicating that regular physical exercise can minimize the

risks of cardiovascular disease. At the same time, running carries a considerable risk of injury. In follow-up cases in the population, the incidence of running-related injury was reported to be 2.5 to 33.0 cases per 1000 h [3]. However, the causes of injuries are varied. Most running-related lower limb injuries, for example, are the result of avoidable training errors [4, 5]. In addition, accumulating long and strong training may lead to an increase in shin pain [6].

Muscle tiredness is a complicated physiological state induced not only by changes in muscle capacity but also by the central nervous system's inability to appropriately drive motor neurons [7]. Long-term running has been proven to cause central fatigue, which diminishes the strength of the maximal autonomic plantar flexor muscle. Plantar flexor fatigue can limit the power of these muscles during the propulsion phase of running, and lower limb strength can be lowered by 30 to 40% after running [8, 9]. The biomechanical features of the lower limbs change as a result of exhaustion, which is crucial in preventing sports injuries. Changes in knee angle and moment because of fatigue, for example, can be used to predict anterior cruciate ligament injuries [10].

Several measurement approaches have been utilized in many studies to quantify the association between foot dynamics and lower extremity overuse injuries. Plantar mechanical measurement has been frequently utilized to evaluate overall running performance as a result of this [11, 12]. The second and third metatarsals exhibit a 10% increase in peak pressure immediately after fatigue, and an 11% increase after 30 mins, with a significant 12% drop in load at the first toe [13]. It is worth noting that increased load under the metatarsal bone can produce biomechanical imbalance, which could lead to metatarsalgia [14]. Furthermore, the increasing plantar load will promote stretching stresses on the plantar aponeurosis, which leads to microtraumas and degradation of connective tissues, promoting the development of plantar fasciitis [15, 16]. In conclusion, there is an urgent need to reflect on and evaluate fatigue and fatigue injuries through changes in plantar mechanical parameters. Insole technologies for activity classification couple plantar pressure with accelerometer data, increasing technology cost, and complexity [17, 18]. The advantage of the platform is that it is easy to use because it is stationary and flat and can be well applied to the laboratory environment [19]. Therefore, we used the footscan force platform to detect the mechanical characteristics of the plantar. Treadmills have been widely used in laboratory studies to easily control speed gradients. Previous studies have also shown that treadmill running is different from running on the ground. Whether treadmill running can simulate running on the ground is still a controversial issue [20]. This experiment only examined the change form of plantar mechanical parameters after fatigue running on a treadmill.

Males and females have different bone structures and muscle strength, and studies have shown that females are more likely than males to sustain lower limb injuries while running [21, 22]. Females are more prone than males to have ligamentous laxity of the ankle joint, and females are approximately twice as likely as males to have ankle sprains [23]. Plantar mechanical parameter distributions are affected by several factors, including weight, gender, foot structure, and even how a person stands or walks [24]. Experts in forensic science use variations in foot bones to determine gender [25]. There are, however, no consistent results on the gender differences in plantar pressure characteristics. According to research [26], there are no significant variations in the midfoot contact area and plantar pressure

between males and females. The pressure under the toes was higher in female adolescents than in male adolescents, while the pressure was higher in male adolescents only at the hindfoot, and the pressure at the metatarsophalangeal toe increased more significantly in females [27]. The difference in plantar mechanical parameters caused by gender can reflect a lot of practical problems. Therefore, it is necessary to explore the effect of gender differences on plantar mechanical parameters.

In biomechanical research, traditional statistical methodologies have limited the ability to classify groups based on many variables [28]. In recent years, a support vector machine (SVM) has emerged as a new tool for solving biological classification problems [29]. By creating discriminatory parameters to separate groups from one another, the SVM attempts to discover a hyperplane that maximizes the distance between groups [30]. The SVM has the advantage of producing classification results based on limited data sets while minimizing structural and empirical risk [31]. Injuries are common in individual sports and will cause serious physical outcomes. Reduced exercise capacity because of fatigue increases the incidence of musculoskeletal injuries [32]. As a result, forecasting the occurrence of sports injuries is critical to maintaining good health [33]. Previous research [34] used the SVM to predict diabetic foot ulceration based on plantar mechanical parameters. Aguirre et al. [35] proposed a computational model for predicting tiredness during exercise from a sitting to a standing posture, which could be useful for rehabilitation. Si et al. [36] employed the SVM and fractal analysis for gait recognition and test the identification performance, and the testing outcomes indicate an overall accuracy of 93.57% via radial basis function kernel. Jeong et al. [37] used the SVM to classify activity patterns based on plantar pressure characteristics, and the recognition rate reached 95.2%. Stetter et al. [38] used the SVM and identified the kinematic difference between fatigue and nonfatigue based on principal component analysis, the strides of fatigue and nonfatigue were separated, and the classification accuracy was 99.4%. Wang et al. [39] used inertial measurement unit (IMU) and SVM to distinguish fatigue and nonfatigue running states, and predict the degree of fatigue. The classification accuracy of tibia and thigh IMUs was 91.10%. The characteristics of plantar pressure were evaluated using leave-one-out cross-validation with machine learning algorithms: SVM, decision tree, discriminant analysis, and k-nearest neighbors in the study of Merry et al. [17]. The results showed that the SVM and decision tree have higher classification accuracy. In addition, other studies have shown that the SVM has the best performance in distinguishing gait characteristics [40]. Therefore, the SVM was used to predict fatigue in this study. In addition, many researchers have applied SVM to the recognition of gait characteristics before and after fatigue, but few studies have paid attention to the plantar mechanical characteristics before and after fatigue.

As a consequence, this research aimed to explore the differences in peak pressure (PP), peak force (PF), and plantar impulse (PI) before and after long-distance running fatigue in novice runners, as well as gender differences. We

also employed the SVM algorithm to predict fatigue based on plantar mechanical parameters. Based on previous studies, we assumed that the change in plantar mechanical parameters before and after fatigue mainly occurred in the toes. It was also assumed that gender differences in plantar mechanical parameters were mainly concentrated in the toes and metatarsal regions. In addition, it was assumed that the SVM can predict fatigue at a high level.

2. Materials and Methods

2.1. Participants. The experimental subjects for this investigation were 15 healthy males and 15 healthy females [13, 25] who were novice runners (Table 1) with dominant right legs. Participants were recruited from sports clubs at Ningbo University and via social media. There were no health issues, neuromuscular abnormalities, or recognized gait difficulties in any of the participants, and no lower limb injuries in six months before the experiment. High arches and flat feet were not allowed to participate in the recruitment process. All subjects were given and signed written consent granted by the Institutional Review Board before the experiment (RAGH20210922205.6).

2.2. Experimental Procedures. Figure 1 depicts the experimental procedure. All of the participants did fatigue-inducing running workouts. The 15-point Borg scale and heart rate monitor (Polar RS100, Polar Electro Oy, Woodbury, NY, USA) were used to record perceived exertion, and heart rate changes per minute during the fatigue intervention. The individuals began the experiment by running at a speed of 1.67 m/s on a treadmill (h/p/cosmos para graphics^R, Germany). During the experiment, the slope was maintained at 1% [41–43]. After which the speed was increased by 0.28 m/s every 2 minutes until the subjects reached a Borg intensity of 13. The subjects then continued at this speed until they reached Borg scale 17 or 90% of maximal heart rate (HR_{max} calculated at 220-age), at which point they slowly reduced the speed to a speed of their choice [44, 45]. Space constraints, repeatability, and better control of climate, speed, and slope were the reasons why treadmill running was selected by our research team [46].

In this experiment, a footscan pressure plate (Footscan[®] software 7.0 Gait 2nd Generation, RsScan International) was used to monitor dynamic plantar pressure. The footscan pressure plate was 2 m in length and the acquisition frequency was 126 Hz. Subjects were asked to perform a pressure measurement on the footscan pressure plate before and immediately after fatigue. To avoid injury during the test, the subjects familiarized themselves with the footscan pressure plate before the trial. After familiarity, the subjects were asked to run on the footscan pressure plate at a speed of 3.3 m/s ± 5% [44]. To manage running speed, Brower timing lights (Brower Timing System, Draper, UT, USA) were used. The subjects who completed a full gait cycle on the footscan pressure plate at the specified speed were regarded as successful. 5 groups of valid data were collected

from each subject before and after the fatigue intervention. In addition, during the fatigue intervention, we uniformly provided clothes and shoes to the subjects to avoid experimental differences and maintain consistency.

2.3. Data Analysis. We analyzed plantar mechanical parameters in the running stance phase. For each trial, ten anatomical zones were automatically identified by the software (Footscan[®] software 7.0 Gait 2nd Generation, RsScan International) and if necessary, manually corrected by adjusting the pixels per zone (Figure 2): hallux (*T1*), second-fifth toes (*T2–5*), metatarsal 1–5 (*M1, M2, M3, M4, M5*), midfoot (*MF*), heel medial (*HM*), and heel lateral (*HL*). During the adjustment, we performed strict controls to ensure that the adjustment conditions and adjustment levels were rigorous and careful. The average values of PP, PF, and PI for all ten regions were calculated.

2.4. Statistical Analysis and SVM Classification Algorithm. The calculated data were exported to a statistical software package SPSS 26.0 (SPSS, Chicago, IL, USA), and the peak pressure, peak force, and plantar impulse of each plantar zone before and after running were statistically processed. The data were initially assessed for normality using a Kolmogorov–Smirnov test. The data were normally distributed. To investigate the effects of fatigue, gender, and their interaction on the plantar mechanical parameters, a two-way analysis of covariance (ANCOVA) was conducted. The significance level was set as $P < 0.05$.

When the data sets were not easily separable, the SVM classifier is a supervised machine learning technique that translates the input data space to a higher dimensional space to obtain a more accurate classification [35]. In our study, we used the LIBSVM toolbox based on MATLAB 2016b (Mathworks, MA, USA). The linear kernel was used for the SVM in the study. The cross-validation technique we employed was the hold-out method. 66.7% of the sample size was randomly selected as the training set, and 33.3% of the sample size was used as the test set [41].

The SVM is suitable for small and medium data samples, and nonlinear, high-dimensional classification problems. It maps the feature vector of the instance to some points in space. The purpose of the SVM is to find a line that best distinguishes two types of points, and when new points are added later, this line can also make a good classification. The SVM will find the partitioning hyperplane that distinguishes the two classes and maximizes the separation. For any hyperplane, the data points on both sides have a minimum distance (vertical distance) from it, and the sum of these two minimum distances is the interval.

For this partitioned hyperplane, we can give the following equation:

$$\omega^T X + b = 0, \quad (1)$$

where ω is the weight of each feature and the column vector. b is the displacement value.

The distance from the point x_i to the surface is as follows:

TABLE 1: Demographic data.

	Age (years)	Height (m)	Body mass (kg)	BMI (kg/m ²)	Shoe size (cm)
Male	23.61 (0.92)	1.83 (0.12)	76.14 (8.12)	24.87 (2.32)	26.35 (0.49)
Female	23.07 (1.04)	1.69 (0.13)	54.73 (4.14)	20.13 (1.08)	24.22 (0.34)

*Values: mean (SD).




FIGURE 1: The experimental procedure.

$$\frac{\omega^T x_i + b}{\|\omega\|}. \quad (2)$$

Then,

$$\frac{\omega^T x_i + b}{\|\omega\|} \times y_i \geq d, \quad (3)$$

y_i is the predicted value of sample i (-1 or 1, doing sign transformation). d is the distance of the support vector to the hyperplane. We assume that d is $(2/\|\omega\|)$.

Making all the points meet:

$$y_i(\omega^T x_i + b) \geq 1. \quad (4)$$

The hyperplane we need is the one that needs to maximize the minimum interval, i.e.,

$$\operatorname{argmax}_{\omega, b} \left\{ \frac{1}{\|\omega\|} \min_i [y_i(\omega^T x_i + b)] \right\}. \quad (5)$$

Then, we need to calculate

$$\operatorname{argmax}_{\omega, b} \frac{1}{\|\omega\|}. \quad (6)$$

Equivalent to calculate

$$\min_{\omega, b} \frac{1}{2} \|\omega\|^2, \quad (7)$$

and

$$y_i(\omega^T x_i + b) \geq 1. \quad (8)$$

Using the Lagrange multiplier method:

$$= \frac{1}{2} \|\omega\|^2 - \sum_{i=1}^n a_i (y_i(\omega^T x_i + b) - 1). \quad (9)$$

The original problem is the minimax problem.

$$\min_{\omega, b} \max_{\alpha} L(\omega, b, \alpha). \quad (10)$$

The dual problem of the original problem is a maximin problem:

$$\max_{\omega, b} \min_{\alpha} L(\omega, b, \alpha). \quad (11)$$

Taking its partial derivative with respect to ω and b and making it equal to 0,




FIGURE 2: The location of ten anatomical zones on the peak mechanical footprint.

$$\omega = \sum_{i=1}^n \alpha_i y_i x_n, \quad (12)$$

$$\sum_{i=1}^n \alpha_i y_i = 0.$$

Then,

$$L(\omega, b, \alpha) = \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i=1}^n \alpha_i \alpha_j y_i y_j x_i x_j. \quad (13)$$

Combining the abovementioned condition:

$$\begin{aligned} \sum_{i=1}^n \alpha_i y_i &= 0 \\ \alpha_i &\geq 0, i = 1, 2 \dots n \end{aligned} \quad (14)$$

Then, we find the maximum value of α .

$$\min \frac{1}{2} \sum_{i=1}^n \alpha_i \alpha_j y_i y_j x_i x_j - \sum_{i=1}^n \alpha_i. \quad (15)$$

We continue to use Lagrange multipliers:

$$\begin{aligned} \omega^* &= \sum_{i=1}^N a_i^* y_i x_i, \\ b^* &= y_i - \sum_{i=1}^N a_i^* y_i x_i x_j. \end{aligned} \quad (16)$$

We find the final hyperplane.

3. Results

3.1. The Peak Pressure. According to Figure 3 and Table 2, fatigue mainly affected the PP at T1, T2–5, HM, and HL, and gender factors were mainly reflected at T1, T2–5, and M3–5. Specifically, PP decreased significantly in T1 and T2–5 regions after fatigue ($P < 0.05$) and increased significantly in HM and HL ($P < 0.05$). PP at T1 and T2–5 was significantly higher in females than in males ($P < 0.05$), and PP at M3–5 was significantly higher in males than in females ($P < 0.05$).

3.2. The Peak Force. According to Figure 3 and Table 2, the fatigue effect was mainly reflected in the T1, T2–5, M1, HM, and HL, while the gender effect was mainly reflected in T1, T2–5, and M3–5. PF was significantly decreased at T1 and T2–5 due to fatigue and significantly increased at M1, HM, and HL ($P < 0.05$). PF in females was significantly larger at T1 and T2–5 than that in males, and significantly smaller at M3–5 than that in males ($P < 0.05$).

3.3. The Impulse. According to Figure 3 and Table 2, the fatigue effect was mainly reflected in the toes, M1, and heel, while the gender effect was mainly reflected in T1, T2–5, and M3–5. PI decreased significantly at T1 and T2–5 after fatigue ($P < 0.05$), and increased significantly at M1, HM, and HL ($P < 0.05$). In addition, the PI at T1 and T2–5 showed that females were significantly larger than males, and at M3–5, females were significantly smaller than males ($P < 0.05$).

3.4. SVM Classification Algorithm. We selected combinations of plantar zone parameters with significant differences ($P < 0.001$). Figure 4 shows the best fit separating hyperplane lines of fatigue or not fatigue in different plantar zone parameter combinations. The accuracy of the different plantar zone parameter combinations in predicting fatigue is presented in Table 3. The results showed that the average accuracy was a moderate level (train accuracy: 62.5%; test accuracy: 62.5%). The accuracies of following combinations were above average and showed a high level: T1 PP/HL PF (train accuracy: 65%; test accuracy: 75%), T1 PF/HL PF (train accuracy: 67.5%; test accuracy: 65%), and HL PF/T1 PI (train accuracy: 67.5%; test accuracy: 70%).

4. Discussion

This research aimed to analyze how PP, PF, and PI changed before and after running fatigue in novice runners, as well as gender differences. Based on previous studies, we assumed that the changes in plantar mechanical parameters before and after fatigue mainly occurred in T1 and T2–5. It was also assumed that gender differences in plantar parameters were mainly concentrated in T1 and T2–5 and M1–5. In addition, it was assumed that SVM can predict fatigue at a high level. Our results are largely consistent with our previous assumptions.

The changes in plantar mechanical parameters caused by fatigue were mainly under the T1, T2–5, M1, HM, and HL. The plantar mechanical parameters in the toes region were also reduced in the research of Bisiaux and Moretto [13],




FIGURE 3: Description of PP, PF, and PI by gender before and after running fatigue.

TABLE 2: Changes in PP, PF, and PI before and after fatigue.

	Male/pre	Male/post	Female/pre	Female/post	<i>F</i>	<i>P</i> value <i>G</i>	<i>F</i> × <i>G</i>
PP (N/kg.m²)							
T1	0.06 (0.02)	0.04 (0.02)	0.12 (0.03)	0.10 (0.01)	0.001	<0.001	0.693
T2-5	0.01 (0.005)	0.01 (0.005)	0.04 (0.02)	0.03 (0.01)	0.002	<0.001	0.115
M1	0.23 (0.05)	0.20 (0.06)	0.24 (0.06)	0.22 (0.07)	0.583	0.396	0.056
M2	0.30 (0.09)	0.28 (0.07)	0.29 (0.07)	0.28 (0.07)	0.256	0.99	0.476
M3	0.24 (0.09)	0.24 (0.07)	0.21 (0.05)	0.18 (0.05)	0.423	0.031	0.156
M4	0.19 (0.08)	0.18 (0.04)	0.14 (0.05)	0.12 (0.03)	0.238	<0.001	0.842
M5	0.12 (0.05)	0.12 (0.04)	0.07 (0.03)	0.07 (0.02)	0.832	<0.001	0.983
MF	0.05 (0.01)	0.05 (0.003)	0.04 (0.02)	0.04 (0.01)	0.283	0.067	0.086
HM	0.19 (0.05)	0.19 (0.04)	0.14 (0.06)	0.16 (0.04)	0.034	0.16	0.453
HL	0.16 (0.04)	0.18 (0.05)	0.13 (0.04)	0.16 (0.05)	0.007	0.29	0.449
PF (N/kg)							
T1	0.96 (0.29)	0.81 (0.40)	2.20 (0.67)	1.69 (0.58)	<0.001	0.001	0.063
T2-5	0.25 (0.14)	0.13 (0.11)	0.83 (0.50)	0.54 (0.30)	0.003	0.004	0.098
M1	3.92 (1.46)	4.54 (1.07)	5.00 (2.03)	5.34 (2.09)	0.017	0.213	0.309
M2	2.87 (0.75)	3.00 (0.92)	2.60 (0.49)	2.83 (0.66)	0.163	0.235	0.975
M3	1.76 (0.67)	1.91 (0.92)	0.99 (0.30)	1.27 (0.33)	0.579	0.014	0.084
M4	1.45 (0.38)	1.55 (0.66)	0.88 (0.30)	0.92 (0.26)	0.578	0.001	0.334
M5	1.18 (0.31)	1.20 (0.50)	0.58 (0.27)	0.59 (0.29)	0.968	<0.001	0.936
MF	2.50 (0.90)	2.27 (0.86)	1.59 (0.87)	1.55 (0.80)	0.079	0.055	0.097
HM	3.14 (0.71)	3.59 (1.12)	3.34 (0.82)	3.85 (0.57)	0.005	0.483	0.838
HL	2.70 (0.73)	3.18 (0.81)	2.73 (0.64)	3.36 (0.89)	<0.001	0.702	0.285
PI (N.s/kg)							
T1	0.13 (0.03)	0.10 (0.06)	0.30 (0.09)	0.22 (0.09)	<0.001	<0.001	0.072
T2-5	0.04 (0.04)	0.03 (0.05)	0.10 (0.06)	0.06 (0.03)	0.017	0.001	0.078
M1	0.47 (0.21)	0.58 (0.21)	0.62 (0.27)	0.66 (0.23)	0.022	0.405	0.054
M2	0.38 (0.16)	0.43 (0.20)	0.36 (0.06)	0.37 (0.10)	0.296	0.162	0.234
M3	0.23 (0.10)	0.27 (0.16)	0.14 (0.04)	0.16 (0.05)	0.901	0.006	0.109
M4	0.18 (0.05)	0.22 (0.12)	0.10 (0.03)	0.11 (0.03)	0.369	<0.001	0.499
M5	0.13 (0.04)	0.14 (0.07)	0.06 (0.03)	0.06 (0.02)	0.597	<0.001	0.711
MF	0.22 (0.08)	0.20 (0.07)	0.13 (0.08)	0.13 (0.07)	0.243	0.066	0.185
HM	0.29 (0.15)	0.31 (0.12)	0.27 (0.07)	0.39 (0.07)	0.021	0.478	0.639
HL	0.23 (0.12)	0.24 (0.10)	0.16 (0.09)	0.23 (0.14)	0.011	0.386	0.413

Pre = before fatigue, post = after fatigue, *F* = fatigue, and *G* = gender. The values in bold in the table show significant differences; $P < 0.05$; values: mean (SD). The bold data in the table indicate statistical significance.

Karagounis et al. [47], and Willems et al. [48]. This may be due to the increased dorsiflexion of the metatarsophalangeal joint after fatigue, which leads to fewer toes contributing to running, and thus less load under the toes [49]. In a study of PP and center of pressure (COP), it was found that the PP under the toes decreased, with a retraction of the COP. According to

Stolwijk et al. [50], to avoid overuse of the forefoot and the risk of incurring forefoot pain, subjects adjusted their gait pattern. This could explain why plantar mechanical parameters under the toes were decreased. Nagel et al. [49] also noted a decline in toes load. However, in the study of Bisiaux and Moretto [13] and Weist et al. [51], the phenomenon of decreased load under




FIGURE 4: SVM classification algorithm results.

TABLE 3: The accuracy of the SVM classification algorithm results.

Plantar zone parameters	Train accuracy (%)	Test accuracy (%)
T1 PP/T1 PF	55	55
T1 PP/HL PF	65	75
T1 PP/T1 PI	65	55
T1 PF/HL PF	67.5	65
T1 PF/T1 PI	55	55
HL PF/T1 PI	67.5	70
Mean accuracy	62.5	62.5

The data in bold indicates that train accuracy and test accuracy are above mean accuracy concurrently.

the toes was not observed. Moreover, in the study of Willems et al. and Wu et al. [48, 52], it was also found that PF and PI at *M1* increased significantly after fatigue, which was consistent with our findings. Perhaps because of the reduction in mechanical parameters under the toes, the load was transferred to the metatarsal. However, increased submetatarsal load may contribute to the occurrence of metatarsal stress fractures [49]. An investigation also revealed that after running fatigue, the contact area of the HM grew while the HL reduced. Then led to greater pronation in the rearfoot [53]. If fatigued, the quadriceps need to play a greater role to avoid knee instability, resulting in less knee flexion, which leads to increased heel pressure. This explanation has also been confirmed by Stolwijk et al. [50].

Several gender-induced differences in plantar mechanical parameters were found. PP, PF, and PI were significantly

higher in females than in males at *T1* and *T2-5* and significantly higher in males than in females at *M3-5*. This was also reflected in studies of Ferrari et al. [54] and Demirbukan et al. [27]. The larger plantar mechanical parameters of females' *T1* and *T2-5* may be related to the fact that females wear high heels, which also raised the risk of chronic paraspinal muscle fatigue, which was linked to postural changes and pain [55]. Although this study did not include cases of hallux valgus (HV), females had a higher load of the hallux than males. Studies reported a meta-analysis that estimated that female HV prevalence (30%) was 2.3 times greater than that in males (13%) [21]. Although many studies cannot reach a unified conclusion, there was no denying that gender differences in plantar mechanical parameters may be one of the reasons for the increase in hallux valgus in females. Males had much more load in the forefoot area than females, which could be due to males' higher body weight, physical structural differences, and females' better flexibility [56, 57]. Further to this, males tend to have a higher vertical center of mass displacement during walking than females. This may also contribute to a higher load in *M3-5* [56]. Pressure is equal to force divided by area. In all regions of the foot, males had a considerably higher contact area than females, both statistically and clinically [25]. At the same time, because of the female hormone secretion, the foot ligament relaxation reduces the degree of stiffness and spreads the forces to a larger extent [58, 59]. This may be the reason that the PP, PF, and PI at *M3-5* are higher in males than in females to varying degrees. In this study, gender differences in PP, PF,

and PI were mainly found in the $T1$, $T2-5$, and $M3-5$. Although the literature's findings are not always consistent, factors including gender and foot anatomy are thought to be linked to metatarsal stress fractures and lower limb injuries [50].

During running, the feet are the only part of the body that makes direct touch with the ground, and they are critical to progress. Running actions may be hampered by muscle exhaustion and physical discomfort. As a result, it is theoretically possible to predict fatigue through plantar mechanical parameters. Previous research has shown that fatigue will affect plantar mechanical parameter distribution and fatigue is correlated with plantar mechanical parameters [48]. Interval maximization is the SVM classification algorithm, which may be characterized as a problem of solving convex quadratic programming and is equivalent to the regularized hinges loss work minimization issue. The SVM classification algorithm is an optimization algorithm for solving convex quadratic programming, as evidenced by our SVM classification algorithm results. The SVM classification algorithm results revealed that the mean accuracy was an above-moderate level. The accuracy was of an above-average level by using the $T1$ PP/HL PF, $T1$ PF/HL PF, and HL PF/ $T1$ PI. These indicated that fatigue can be predicted to a certain extent by monitoring plantar mechanical parameters before and after running fatigue. Running fatigue can be predicted using the learned SVM classification algorithm, which can also be used as a useful tool for fatigue supervision. The learned SVM classification algorithm can help coaches to better identify the physical state of athletes from start to the finish of a run by monitoring plantar mechanical parameters. The classification may also be useful in identifying injuries over the running season.

There are some limits of this study. In the experiment, a treadmill was used for the fatigue intervention. We only studied the plantar mechanical parameters under treadmill conditions but did not study the condition of running on the ground. Future studies should include subjects performing at different exercise levels, such as professional athletes. In addition, the sample size should be expanded to improve the accuracy of the SVM classification algorithm.

5. Conclusions

We found that the change of plantar mechanical parameters caused by fatigue was mainly concentrated in $T1$, $T2-5$, $M1$, HM , and HL . While the effect of gender was mainly found in the $T1$, $T2-5$, and $M3-5$. These may indicate injuries related to fatigue and gender factors, such as metatarsal stress fractures and HV. Plantar mechanical parameters can be monitored before and after long-distance running to predict fatigue to some extent. The learned algorithm of plantar zone combinations with above-average accuracy ($T1$ PP/HL PF, $T1$ PF/HL PF, and HL PF/ $T1$ PI) can predict long-distance running fatigue and provide supervised training strategies.

Data Availability

The data used to support the findings of this study are available from the corresponding authors upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

Q. L. and H. C. conceived the presented idea, developed the framework, and wrote the manuscript. A. T., J. S. B., M. Y., and Y. G. provided critical feedback and contributed to the final version. All authors were involved in the final direction of the paper and contributed to the final version of the manuscript. All authors have read and agreed to the published version of the manuscript.

Acknowledgments

This research was funded by the Zhejiang Provincial Natural Science Foundation of China for Distinguished Young Scholars (Grant no. LR22A020002); the Zhejiang Provincial Key Research and Development Program of China (Grant no. 2021C03130); the Philosophy and Social Sciences Project of Zhejiang Province, China (Grant nos. 22QNYC10ZD and 22NDQN223YB); the Public Welfare Science and Technology Project of Ningbo, China (Grant No. 2021S134); and the K. C. Wong Magna Fund in Ningbo University.

References

- [1] D. Cooper, R. Kavanagh, J. Bolton, and L. Keaver, "A pilot 6-week lifestyle intervention in women aged 50+ in Ireland," *Physical Activity and Health*, vol. 6, no. 1, pp. 180–188, 2022.
- [2] R. S. Schwartz, S. M. Kraus, J. G. Schwartz et al., "Increased coronary artery plaque volume among male marathon runners," *Missouri Medicine*, vol. 111, no. 2, pp. 89–94, 2014.
- [3] A. Hulme, R. O. Nielsen, T. Timpka, E. Verhagen, and C. Finch, "Risk and protective factors for middle-and-long-distancerunning-related injury," *Sports Medicine*, vol. 47, no. 5, pp. 869–886, 2017.
- [4] S. A. Paluska, "An overview of hip injuries in running," *Sports Medicine*, vol. 35, no. 11, pp. 991–1014, 2005.
- [5] B. T. Saragiotto, T. P. Yamato, L. C. Hespanhol Junior, M. J. Rainbow, I. S. Davis, and A. D. Lopes, "What are the main risk factors for running-related injuries?" *Sports Medicine*, vol. 44, no. 8, pp. 1153–1163, 2014.
- [6] S. S. Yeung, E. W. Yeung, and L. D. Gillespie, "Interventions for preventing lower limb soft-tissue running injuries," *Cochrane Database of Systematic Reviews*, vol. 6, no. 7, pp. CD001256–142, 2011.
- [7] Y. Chandra, R. Tewari, and A. Jain, "Experimental studies on acrylic dielectric elastomers as actuator for artificial skeletal muscle application," *International Journal of Biomedical Engineering and Technology*, vol. 37, no. 1, pp. 65–82, 2021.
- [8] A. Saldanha, M. M. Nordlund Ekblom, and A. Thorstensson, "Central fatigue affects plantar flexor strength after prolonged running," *Scandinavian Journal of Medicine & Science in Sports*, vol. 18, no. 3, pp. 383–388, 2008.
- [9] M. Giandolini, G. Vernillo, P. Samozino et al., "Fatigue associated with prolonged graded running," *European Journal of Applied Physiology*, vol. 116, no. 10, pp. 1859–1873, 2016.
- [10] K. Balachandar, N. Muralidharan, N. Jawahar, and K. Chockalingam, "Development of comfort fit lower limb prosthesis by reverse engineering and rapid prototyping

- methods and validated with gait analysis,” *International Journal of Biomedical Engineering and Technology*, vol. 35, no. 4, pp. 362–381, 2021.
- [11] G. J. Dowling, G. S. Murley, S. E. Munteanu et al., “Dynamic foot function as a risk factor for lower limb overuse injury: a systematic review,” *Journal of Foot and Ankle Research*, vol. 7, pp. 53–13, 2014.
 - [12] H. K. Kim, S. A. Mirjalili, and J. Fernandez, “Gait kinetics, kinematics, spatiotemporal and foot plantar pressure alteration in response to long-distance running: systematic review,” *Human Movement Science*, vol. 57, pp. 342–356, 2018.
 - [13] M. Bisiaux and P. Moretto, “The effects of fatigue on plantar pressure distribution in walking,” *Gait & Posture*, vol. 28, no. 4, pp. 693–698, 2008.
 - [14] J.-H. Kang, M.-D. Chen, S.-C. Chen, and W.-L. Hsi, “Correlations between subjective treatment responses and plantar pressure parameters of metatarsal pad treatment in metatarsalgia patients: a prospective study,” *BMC Musculoskeletal Disorders*, vol. 7, pp. 95–98, 2006.
 - [15] A. P. Ribeiro, F. Trombini-Souza, V. D. Tessutti, F. R. Lima, S. M. João, and I. C. Sacco, “The effects of plantar fasciitis and pain on plantar pressure distribution of recreational runners,” *Clinical Biomechanics*, vol. 26, no. 2, pp. 194–199, 2011.
 - [16] H. Chen, Y. Song, R. Xuan, Q. Hu, J. S. Baker, and Y. Gu, “Kinematic comparison on lower limb kicking action of fetuses in different gestational weeks: a pilot study,” *Healthcare*, vol. 9, p. 1057, 2021.
 - [17] K. J. Merry, E. Macdonald, M. MacPherson et al., “Classifying sitting, standing, and walking using plantar force data,” *Medical, & Biological Engineering & Computing*, vol. 59, no. 1, pp. 257–270, 2021.
 - [18] E. S. Sazonov, G. Fulk, J. Hill, Y. Schutz, and R. Browning, “Monitoring of posture allocations and activities by a shoe-based wearable sensor,” *IEEE Transactions on Biomedical Engineering*, vol. 58, no. 4, pp. 983–990, 2011.
 - [19] A. H. Abdul Razak, A. Zayegh, R. K. Begg, and Y. Wahab, “Foot plantar pressure measurement system: a review,” *Sensors*, vol. 12, no. 7, pp. 9884–9912, 2012.
 - [20] Y. Hong, L. Wang, J. X. Li, and J. H. Zhou, “Comparison of plantar loads during treadmill and overground running,” *Journal of Science and Medicine in Sport*, vol. 15, no. 6, pp. 554–560, 2012.
 - [21] T. Yamamoto, Y. Hoshino, N. Kanzaki et al., “Plantar pressure sensors indicate women to have a significantly higher peak pressure on the hallux, toes, forefoot, and medial of the foot compared to men,” *Journal of Foot and Ankle Research*, vol. 13, pp. 40–47, 2020.
 - [22] J. Dempster, F. Dutheil, and U. C. Ugbohue, “The prevalence of lower extremity injuries in running and associated risk factors: a systematic review,” *Physical Activity and Health*, vol. 5, no. 1, pp. 133–145, 2021.
 - [23] J. M. Wolf, L. Cannada, A. E. Van Heest, M. I. O’Connor, and A. L. Ladd, “Male and female differences in musculoskeletal disease,” *Journal of the American Academy of Orthopaedic Surgeons*, vol. 23, no. 6, pp. 339–347, 2015.
 - [24] C. K. Ang, M. I. Solihin, W. J. Chan, and Y. Y. Ong, “Study of plantar pressure distribution,” *MATEC Web of Conferences*, vol. 237, Article ID 01016, 2018.
 - [25] A. Putti, G. Arnold, and R. Abboud, “Foot pressure differences in men and women,” *Foot and Ankle Surgery*, vol. 16, no. 1, pp. 21–24, 2010.
 - [26] D. F. Murphy, B. D. Beynon, J. D. Michelson, and P. M. Vacek, “Efficacy of plantar loading parameters during gait in terms of reliability, variability, effect of gender and relationship between contact area and plantar pressure,” *Foot & Ankle International*, vol. 26, no. 2, pp. 171–179, 2005.
 - [27] İ. Demirbüken, B. Özgül, E. Timurtaş, S. U. Yurdalan, M. D. Çekin, and M. G. Polat, “Gender and age impact on plantar pressure distribution in early adolescence,” *Acta Orthopaedica et Traumatologica Turcica*, vol. 53, no. 3, pp. 215–220, 2019.
 - [28] R. K. Fukuchi, B. M. Eskofier, M. Duarte, and R. Ferber, “Support vector machines for detecting age-related changes in running kinematics,” *Journal of Biomechanics*, vol. 44, no. 3, pp. 540–542, 2011.
 - [29] Y.-Y. Chan, D. T.-P. Fong, M. M.-L. Chung et al., “Identification of ankle sprain motion from common sporting activities by dorsal foot kinematics data,” *Journal of Biomechanics*, vol. 43, no. 10, pp. 1965–1969, 2010.
 - [30] S. R. Sain, *The Nature of Statistical Learning Theory*, Springer, Berlin, Germany, 1996.
 - [31] J. Zhang, T. E. Lockhart, and R. Soangra, “Classifying lower extremity muscle fatigue during walking using machine learning and inertial sensors,” *Annals of Biomedical Engineering*, vol. 42, no. 3, pp. 600–612, 2014.
 - [32] J. Verschuere, B. Tassignon, K. De Pauw et al., “Does acute fatigue negatively affect intrinsic risk factors of the lower extremity injury risk profile? A systematic and critical review,” *Sports Medicine*, vol. 50, no. 4, pp. 767–784, 2020.
 - [33] H. Van Eetvelde, L. D. Mendonça, C. Ley, R. Seil, and T. Tischer, “Machine learning methods in sport injury prediction and prevention: a systematic review,” *Journal of experimental orthopaedics*, vol. 8, pp. 27–15, 2021.
 - [34] F. S. Botros, M. F. Taher, N. M. ElSayed, and A. S. Fahmy, “Prediction of diabetic foot ulceration using spatial and temporal dynamic plantar pressure,” in *Proceedings of the 2016 8th Cairo international biomedical engineering conference (CIBEC)*, pp. 43–47, Cairo, Egypt, December 2016.
 - [35] A. Aguirre, M. J. Pinto, C. A. Cifuentes, O. Perdomo, C. A. R. Díaz, and M. Múnera, “Machine learning approach for fatigue estimation in sit-to-stand exercise,” *Sensors*, vol. 21, no. 15, p. 5006, 2021.
 - [36] W. Si, G. Yang, X. Chen, and J. Jia, “Gait identification using fractal analysis and support vector machine,” *Soft Computing*, vol. 23, no. 19, pp. 9287–9297, 2019.
 - [37] G.-M. Jeong, P. H. Truong, and S.-I. Choi, “Classification of three types of walking activities regarding stairs using plantar pressure sensors,” *IEEE Sensors Journal*, vol. 17, no. 9, pp. 2638–2639, 2017.
 - [38] B. J. Stetter, F. Möhler, F. C. Krafft, S. Sell, and T. Stein, “Identification of fatigue-related kinematic changes in elite runners using a support vector machine approach,” *ISBS Proceedings Archive*, vol. 38, p. 264, 2020.
 - [39] G. Wang, X. Mao, Q. Zhang, and A. Lu, “Fatigue detection in running with inertial measurement unit and machine learning,” in *Proceedings of the 2022 10th International Conference on Bioinformatics and Computational Biology*, pp. 85–90, ICBCB), Hangzhou, China, May 2022.
 - [40] K. Trentzsch, P. Schumann, G. Śliwiński et al., “Using machine learning algorithms for identifying gait parameters suitable to evaluate subtle changes in gait in people with multiple sclerosis,” *Brain Sciences*, vol. 11, no. 8, p. 1049, 2021.
 - [41] F. Aghakeshizadeh, A. Letafatkar, P. A. Ataabadi, and M. Hosseinzadeh, “The effect of taping on maximum plantar pressure and ground reaction force in people with flat foot after applying a fatigue protocol,” vol. 7, pp. 203–226, 2021.

- [42] F. García-Pinillos, A. Cartón-Llorente, D. Jaén-Carrillo et al., “Does fatigue alter step characteristics and stiffness during running?” *Gait & Posture*, vol. 76, pp. 259–263, 2020.
- [43] A. C. Clansey, M. Hanlon, E. S. Wallace, and M. J. Lake, “Effects of fatigue on running mechanics associated with tibial stress fracture risk,” *Medicine & Science in Sports & Exercise*, vol. 44, no. 10, pp. 1917–1923, 2012.
- [44] M. Anbarian and H. Esmaeili, “Effects of running-induced fatigue on plantar pressure distribution in novice runners with different foot types,” *Gait & Posture*, vol. 48, pp. 52–56, 2016.
- [45] I. F. Koblbauer, K. S. van Schooten, E. A. Verhagen, and J. H. van Dieën, “Kinematic changes during running-induced fatigue and relations with core endurance in novice runners,” *Journal of Science and Medicine in Sport*, vol. 17, no. 4, pp. 419–424, 2014.
- [46] J. A. García-Pérez, P. Pérez-Soriano, S. Llana, A. Martínez-Nova, and D. Sánchez-Zuriaga, “Effect of overground vs. treadmill running on plantar pressure: influence of fatigue,” *Gait & Posture*, vol. 38, no. 4, pp. 929–933, 2013.
- [47] P. Karagounis, G. Prionas, E. Armenis, G. Tsiganos, and P. Baltopoulos, “The impact of the Spartathlon ultramarathon race on athletes’ plantar pressure patterns,” *Foot & Ankle Specialist*, vol. 2, no. 4, pp. 173–178, 2009.
- [48] T. M. Willems, R. De Ridder, and P. Roosen, “The effect of a long-distance run on plantar pressure distribution during running,” *Gait & Posture*, vol. 35, no. 3, pp. 405–409, 2012.
- [49] A. Nagel, F. Fernholz, C. Kibele, and D. Rosenbaum, “Long distance running increases plantar pressures beneath the metatarsal heads: a barefoot walking investigation of 200 marathon runners,” *Gait & Posture*, vol. 27, no. 1, pp. 152–155, 2008.
- [50] N. M. Stolwijk, J. Duysens, J. W. K. Louwerens, and N. L. W. Keijsers, “Plantar pressure changes after long-distance walking,” *Medicine & Science in Sports & Exercise*, vol. 42, no. 12, pp. 2264–2272, 2010.
- [51] R. Weist, E. Eils, and D. Rosenbaum, “The influence of muscle fatigue on electromyogram and plantar pressure patterns as an explanation for the incidence of metatarsal stress fractures,” *The American Journal of Sports Medicine*, vol. 32, no. 8, pp. 1893–1898, 2004.
- [52] W.-L. Wu, J.-J. Chang, J.-H. Wu, L.-Y. Guo, and H.-T. Lin, “EMG and plantar pressure patterns after prolonged running,” *Biomedical Engineering: Applications, Basis and Communications*, vol. 19, no. 06, pp. 383–388, 2007.
- [53] E. Escamilla-Martínez, A. Martínez-Nova, B. Gómez-Martín, R. Sánchez-Rodríguez, and L. M. Fernández-Seguín, “The effect of moderate running on foot posture index and plantar pressure distribution in male recreational runners,” *Journal of the American Podiatric Medical Association*, vol. 103, no. 2, pp. 121–125, 2013.
- [54] J. Ferrari and D. Watkinson, “Foot pressure measurement differences between boys and girls with reference to hallux valgus deformity and hypermobility,” *Foot & Ankle International*, vol. 26, no. 9, pp. 739–747, 2005.
- [55] M. Gimunová, M. Zvonař, and O. Mikeska, “The effect of aging and gender on plantar pressure distribution during the gait in elderly,” *Acta of Bioengineering and Biomechanics*, vol. 20, no. 4, pp. 139–144, 2018.
- [56] M.-J. Chung and M.-J. Wang, “Gender and walking speed effects on plantar pressure distribution for adults aged 20–60 years,” *Ergonomics*, vol. 55, no. 2, pp. 194–200, 2012.
- [57] B. Zhang and Q. Lu, “A current review of foot disorder and plantar pressure alternation in the elderly,” *Physical Activity and Health*, vol. 4, no. 1, pp. 95–106, 2020.
- [58] D. Xu, W. Quan, H. Zhou, D. Sun, J. S. Baker, and Y. Gu, “Explaining the differences of gait patterns between high and low-mileage runners with machine learning,” *Scientific Reports*, vol. 12, no. 1, p. 2981, 2022.
- [59] L. Xiang, Q. Mei, A. Wang, V. Shim, J. Fernandez, and Y. Gu, “Evaluating function in the hallux valgus foot following a 12-week minimalist footwear intervention: a pilot computational analysis,” *Journal of Biomechanics*, vol. 132, Article ID 110941, 2022.