

Review Article

Immunosuppressive Role of Myeloid-Derived Suppressor Cells and Therapeutic Targeting in Lung Cancer

Jie Ma,^{1,2} Huaxi Xu,² and Shengjun Wang^{1,2} 

¹Department of Laboratory Medicine, The Affiliated People's Hospital, Jiangsu University, Zhenjiang, China

²Institute of Laboratory Medicine, Jiangsu Key Laboratory of Laboratory Medicine, Jiangsu University School of Medicine, Zhenjiang, China

Correspondence should be addressed to Shengjun Wang; sjwjs@ujs.edu.cn

Received 27 July 2017; Revised 10 January 2018; Accepted 29 January 2018; Published 25 March 2018

Academic Editor: Nejat K. Egilmez

Copyright © 2018 Jie Ma et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Lung cancer is the leading cause of cancer death worldwide due to its late diagnosis and poor outcome. Immunotherapy is becoming more and more encouraging and promising in lung cancer therapy. Myeloid-derived suppressor cells (MDSCs) are the main tumor suppressor factors, and the treatment strategy of targeting MDSCs is gradually emerging. In this review, we summarize what is currently known about the role of MDSCs in lung cancer. In view of the emerging importance of MDSCs in lung cancer, the treatment of targeting MDSCs will be useful to the control of the development and progression of lung cancer. However, the occurrence, metastasis, and survival of cancer is the result of multiple factors and multiple mechanisms, so combined treatments using different strategies will become the major therapy method for lung cancer in the future.

1. Introduction

Lung cancer is a challenging health problem and the leading cause of cancer-related mortality in developed countries, where more than 1.0 million people die of the disease each year [1]. Despite advances in the treatment of lung cancer with chemotherapy and the integration of targeted therapy, the overall outcomes remain poor. A better understanding of the immunologic properties of lung cancer has led to novel treatment strategies, including immune checkpoint modulation and vaccine therapy [2]. Recent clinical trials in lung cancer demonstrate the potential of immunotherapeutics to increase the overall survival in patients with lung cancer compared to the current standard of care [3].

Myeloid-derived suppressor cells (MDSCs) are a heterogeneous population of cells that consists of myeloid progenitor cells and immature granulocytes, immature macrophages, and immature dendritic cells (DCs) [4]. MDSCs play a critical role in tumor-associated immunosuppressive function, which plays an important role in the effective immunotherapies for cancer. In mice, MDSCs are identified by the expression of CD11b and Gr-1 on the cell surface,

and the Gr-1 molecule includes Ly6G and Ly6C. CD11b⁺Ly6G⁻Ly6C^{high} cells showing monocytic-like morphology are called monocytic MDSCs (M-MDSCs), and CD11b⁺Ly6G⁺Ly6C^{low} cells showing granulocyte-like morphology are called granulocytic MDSCs (G-MDSCs) [5]. MDSCs also express histamine and histamine receptor 1 (HR1), which enhances the survival and expansion of MDSCs [6]. In humans, MDSCs are defined by the expression of CD33 on the cell surface but lack the expression of markers of mature myeloid and lymphoid cells [4]. The equivalents to PMN-MDSCs are defined as CD11b⁺CD14⁻CD15⁺ or CD11b⁺CD14⁻CD66b⁺, and equivalents to M-MDSCs, as CD11b⁺CD14⁺HLA-DR^{-/low}CD15⁻ in human peripheral blood mononuclear cells (PBMC) [7]. In addition, there is a third population of MDSCs in humans. The early-stage MDSCs are termed Lin⁻HLA-DR⁻CD33⁺ [7, 8]. In cancer patients, MDSCs could strongly inhibit the antitumor immune responses of CD4⁺ T cells, CD8⁺ T cells, and NK cells and promote the progression of tumors. Currently, strategies to target MDSCs in cancer immunotherapy mainly involve promoting the differentiation of MDSCs, inhibiting their suppressive effect, or eliminating the cells.

2. Mechanisms of MDSC-Mediated Immune Suppression

MDSCs comprise a heterogeneous population of immature myeloid cells that exert the protumor immune response function via a variety of mechanisms. It is believed that MDSCs are major contributors to mediating tumor escapes. MDSCs are able to induce tolerance to a variety of immune responses mediated by effector T cells and NK cells. Both M-MDSCs and G-MDSCs could inhibit effector T cells by different manners [4]. M-MDSCs predominantly play the role of immune suppressor by the production of Arg-1 and generation of NO, whereas G-MDSCs mainly produce ROS and Arg-1 [8].

2.1. Arg-1 and NO. MDSCs are able to express high levels of Arg-1 and NO, while these two molecules have the effect of inhibiting the function of T cells [9, 10]. The suppressive activity of Arg-1 is based on its role in the hepatic urea cycle, metabolizing L-arginine to L-ornithine. A study showed that Arg-1 was closely related to the proliferation of T cells [11]. A PEGylated form of the catabolic enzyme arginase-1 (peg-Arg-1) can enhance the growth of tumors in mice in a manner that correlated with higher MDSC numbers [12]. The enhancement of the activity of Arg-1 in MDSCs causes the decomposition of arginine, which leads to the decrease of L-arginine, and inhibits the proliferation of T cells by various mechanisms, including the downregulation of CD3 expression and the inhibition of cyclin D3 and cyclin-dependent kinase 4 expression [13]. NO can inhibit the function of JAK3 and STAT5 by inducing the apoptosis of T cells [14] or inhibit the proliferation of T cells by inhibiting the expression of MHC-II [15].

2.2. ROS. Another important factor associated with the immunosuppressive ability of MDSCs is reactive oxygen species (ROS). The upregulation of the expression of ROS in tumor-bearing mice and tumor patients is a major feature of MDSCs [16–19]. The expression of ROS in tumor-bearing mice and tumor patients could significantly enhance the immunosuppression of MDSCs [16]. Interestingly, the binding of integrin on the surface of MDSCs after the action between MDSCs and T cells increased the expression of ROS [20]. In addition, other factors such as GM-CSF, IL-10, TGF- β , IL-6, PDGF, and IL-3 can induce the production of ROS by MDSCs [21].

2.3. Peroxynitrite. The product of the superoxide anion and NO chemical reaction is another factor that inhibits effector T cells [20]. The expression level of peroxynitrite was significantly increased in the accumulation of MDSCs and inflammatory cells. In many different tumors, a high content of peroxynitrite is closely related to the process of tumor growth. In addition, this is related to the failure of T cells to respond [22–26]. Peroxynitrite can damage the expression of MHC-II and mediate T-cell apoptosis [4, 27]. Moreover, peroxynitrite leads to the nitration of tyrosines in the TCR-CD8 complex, which can damage the conformational flexibility of the complex, affecting its interaction with

peptide-loaded MHC-I and leading to the unresponsiveness of CD8⁺ T cells to antigen-specific stimulation [4, 27, 28]. In addition to inhibiting the activation of T cells, MDSCs were able to influence the immune response by interfering with the innate immune response, mainly through the influence of NK cells, macrophages, and NKT cells. The effect of MDSCs on NK cells is complex. Some subsets can inhibit the killing of NK cells by blocking the production of IFN- γ . Other subsets can activate NK cells and enhance the killing of them by expressing RAE-1, which interacts with NKG2D on the surface of NK cells [29, 30]. A recent report showed that IL-13 mediated the effect through the IL-4R-STAT6 pathway and induced TGF- β -producing CD11b⁺Gr-1⁺ MDSCs. The production of TGF- β , IL-13, and IL-4 impaired the function of NK cells [31].

2.4. Tregs. The population of regulatory T cells (Tregs) plays a crucial role in tumor immune escape [32, 33]. It has been reported that MDSCs could promote the development of Tregs [32, 33]. MDSCs have been shown to not be involved in the induction of Tregs; however, they may be involved in the differentiation of Tregs by releasing cytokines or cell-cell contact [34].

2.5. Exosomes. Exosomes are present in high abundance in the tumor microenvironment, where they transfer information between different cells. Deng et al. found that MDSC-derived exosomes polarize macrophages toward a tumor-promoting phenotype, demonstrating that some of the tumor-promoting functions of MDSC are mediated by MDSC-shed exosomes [35].

2.6. Metabolic Regulation. It has been noticed that MDSCs from tumors have a stronger immunosuppressive function than MDSCs in the peripheral lymphoid organs. Some newer studies suggest that MDSC maturation and function is under the control of metabolism in the tumor microenvironment [36, 37]. Compared to spleen-MDSCs, tumor-infiltrating MDSCs (T-MDSCs) increased fatty acid uptake and activated fatty acid oxidation (FAO) [38]. Husain et al. provide evidence of an immunosuppressive role of tumor-derived lactate in inhibiting innate immune responses against developing tumors via the regulation of MDSC activity [39]. In addition to the effects of lipid metabolism and lactate, the glycolysis pathway can also affect the maturation and function of MDSCs. Liu et al. showed that the SIRT1-mTOR/HIF-1 α glycolytic pathway was determined by the differentiation of MDSCs [40]. mTORC1 intrinsically controls CD11b⁺Ly6C^{high} M-MDSC maturation and function by mediating cellular glycolysis activity [36].

3. Potential Importance of MDSCs in Lung Cancer

MDSCs may provide predictive and prognostic information in lung cancer patients. The function of MDSCs as biomarkers of lung cancer involves measurements of different cell subsets in the peripheral blood of patients. Tian et al. demonstrated that the number and frequency of peripheral

CD14⁺HLA-DR^{-low} MDSCs were significantly increased in SCLC patients compared with those in controls and the frequency of MDSCs correlated with tumor stage [41]. Two years before that, Huang et al. reported similar results in that both the frequency and absolute number of CD14⁺HLA-DR^{-low} cells were significantly increased in the peripheral blood of NSCLC patients and indicated an association with metastasis, response to chemotherapy, and progression-free survival [42]. Expecting that the frequency and number of MDSCs could distinguish between lung cancer patients and healthy controls, immunoglobulin-like transcript 3 (ILT3), which is expressed by MDSCs [43], and arginase-1 (Arg-1) mRNA [44], which is expressed by MDSCs, could also be used as surrogate markers for the frequency of MDSCs in PBMC and as attractive targets for immune intervention. Patients with NSCLC had a significantly higher ratio of CD11b⁺CD14⁺ cells than healthy subjects, which was correlated with poor performance status and poor response to chemotherapy [45]. In a study by Zhang et al., the clinical data analysis indicated that a higher frequency of B7⁻H3⁺ MDSCs was associated with reduced recurrence-free survival in patients with NSCLC [46]. The data provide evidence that increased percentages of new M-MDSC subpopulations in advanced NSCLC patients are associated with an unfavorable clinical outcome [47].

Based on clinical experience, the treatment appears to influence levels of MDSCs in lung cancer patients. Wang et al. showed that in 20 patients with advanced NSCLC who received systemic chemotherapy, 9 partial remission cases had MDSC percentages that significantly decreased, 3 stable disease cases remained invariable, and 8 progressive disease cases had MDSC percentages that significantly increased [48]. Recently, results showed that three cycles of bevacizumab-containing regimens significantly reduced the percentage of granulocytic MDSCs compared with nonbevacizumab-based regimens [49]. Elevated serum levels of TNF- α , CCL-2, and CCL-4 associated with an increased NO production in circulating MDSCs might be an early indicator of incomplete radiofrequency ablation and, subsequently, a potential tumor relapse in NSCLC [50]. Taken together, the reduced levels of MDSCs may be interrelated with the efficacy of chemotherapy.

Murine models are regularly used to study the relationship between MDSCs and lung cancer. In murine models of lung cancer, B7⁻H3⁺ MDSCs were found only in the tumor microenvironment, and their frequencies increased during tumor progression [46]. Parallel increases in the level of galectin-3 with the number of MDSCs in vivo were detected after cisplatin treatment [51]. Furthermore, acute exposure to single-walled carbon nanotubes (SWCNT) induced the recruitment and accumulation of lung-associated MDSCs and the MDSC-derived production of TGF- β , resulting in an upregulated tumor burden in the lung [52]. As we know, smoking is the leading cause of lung cancer. The effect of smoking on MDSCs function is less reported. When Ortiz et al. exposed mice to cigarette smoke (CS) alone, it resulted in a significant accumulation in various organs of cells with typical MDSC phenotype, but these cells lacked immunosuppressive activity. When CS was combined with a single dose

of urethane, it led to Gr-1⁺CD11b⁺ cells accumulating in the spleen and lung, and they had potent immunosuppressive activity [53].

MDSCs have shown an increasing trend in lung cancer patients and murine models, correlating with tumor progression, increased severity of the disease, and poor prognosis and survival.

4. MDSCs Are a Potential Target for Therapeutic Development in Lung Cancer

Along with the development of MDSCs, many factors have been found to regulate MDSCs in recent years. Multiple signaling pathways and cytokines were found to participate in the regulation of MDSCs. Most of the factors regulate the differentiation and maturation of myeloid cells by the JAK-STAT and NF- κ B signaling pathways and then affect the production and activation of MDSCs. The interaction of all these factors constitutes a complex network control system that regulates the generation and function of MDSCs.

To succeed in the implementation of tumor immunotherapy, the tumor suppressor factors must be removed. MDSCs are the main tumor suppressor factors, so the treatment strategy of targeting MDSCs is gradually emerging (Figure 1). The tumor immunotherapy can be effectively enhanced by targeting the numbers and function of MDSCs. In general, the mechanisms that can be implemented to reverse the number and function of MDSCs focus on four main categories (Table 1).

4.1. Promotion of Myeloid Cell Differentiation. One of the most popular methods in treatment by targeting MDSCs is to promote the differentiation of immature MDSCs into myeloid cells. Retinoic acid, a product of the metabolism of vitamin A, can stimulate the differentiation of myeloid progenitor cells to dendritic cells or macrophages [54]. In a clinical trial of patients with extensive stage small cell lung cancer (SCLC), Iclozan et al. showed that vaccine alone did not affect the proportion of MDSCs, whereas in patients treated with vaccination in combination with MDSCs targeted by therapy with all-*trans*-retinoic acid (ATRA), the MDSCs decreased more than twofold [61]. In our previous study, we found that whole β -glucan particles (WGP) could promote the differentiation and maturation of MDSCs via the dectin-1 pathway in vitro and decrease the suppressive function of cells, thus leading to enhanced antitumor immune responses [55].

4.2. Inhibition of MDSC Expansion. The amplification of MDSCs is regulated by many factors, such as stem cell factor (SCF) and vascular endothelial growth factor (VEGF). c-KIT, the receptor of SCF, can inhibit the signaling pathway which is mediated by SCF, thus inhibiting the amplification of MDSCs and tumor angiogenesis [58]. VEGF is another factor that can promote the expansion of MDSCs, so it can be used as another effective target of MDSCs. However, the mechanism of VEGF in lung cancer has not been reported yet.

The STAT family, especially STAT3, plays an essential role in the regulation of the production, amplification, and




FIGURE 1: Targeting MDSCs in the treatment of lung cancer. Retinoic acid can stimulate the differentiation of myeloid progenitor cells to dendritic cells or macrophages, thereby inhibiting the differentiation of MDSCs. c-KIT can inhibit the signaling pathway which is mediated by SCF, inhibiting the amplification of MDSCs. The inhibition of COX2 expression in MDSCs can decrease the release of arginine-1, Nrf2 contributes to the clearance of ROS in MDSCs, and both COX2 and Nrf2 can inhibit the function of MDSCs. The above measures will inhibit MDSCs' immunosuppressive function on effector T cells and enhance the antitumor immunity. SCF: stem cell factor; VEGF: vascular endothelial growth factor; COX2: cyclooxygenase 2; Arg-1: arginine-1; Nrf2: nuclear factor erythroid 2 p45-related factor 2; ROS: reactive oxygen species; CTL: cytotoxic lymphocyte.

TABLE 1: Regulation of MDSCs in lung cancer.

Treatment	Mouse model versus patients	Effect on MDSCs	References
Gemcitabine	Mouse models	Inhibition of MDSC expansion	[29]
Retinoic acid	Mouse models	Promotion of the differentiation of immature MDSCs	[54]
WGP	Mouse models	Promotion of the differentiation and maturation of MDSCs	[55]
Cyclooxygenase 2 inhibitor	Mouse models	Inhibition of the suppressive effects of MDSCs	[11, 56]
CSF-1 receptor antagonist(GW2580)	Mouse models	Inhibition of the recruitment of MDSCs	[57]
Anti-c-KIT	Mouse models	Inhibition of the amplification of MDSCs	[58]
Anti-Gr-1 or anti-Ly6G antibodies	Mouse models	Depletion of MDSCs	[59]
Indomethacin	Mouse models	Inhibition of the suppressive effects of MDSCs	[60]
ATRA	Patients	Inhibition of MDSCs expansion	[61]
Triterpenoids	Mouse models and patients	Inhibition of the suppressive effects of MDSCs	[62]

function of MDSCs. STAT3 is, controversially, the main transcription factor which regulates the expansion of MDSCs. MDSCs from tumor-bearing mice have greatly increased levels of phosphorylated STAT3 compared with IMCs from naive mice, probably by upregulating the expression of STAT3 target genes, including B-cell lymphoma XL (BCL-XL), Myc, cyclin D1, and survivin. Blocking STAT3 expression in conditional knockout mice or STAT3 inhibitors could markedly reduce the expansion of MDSCs and

increase T-cell responses in tumor-bearing mice [63, 64]. STAT3 can also regulate MDSCs' expansion by inducing the expression of S100A8 and S100A9 proteins, which belong to the family of S100 calcium-binding proteins that have been reported to have an important role in inflammation [65]. Wu et al. verified that Stat3C promotes MDSCs' expansion and immune suppression during lung tumorigenesis [66]. A recent report suggested that the activation of STAT3 in MDSCs and macrophages promoted tumorigenesis

through pulmonary recruitment and increased resistance of suppressive cells to CD8⁺ T cells in lung cancer development [67]. The upregulation of CD45 tyrosine phosphatase activity in MDSCs exposed to hypoxia in a tumor site was responsible for the downregulation of STAT3, and STAT3 has a unique function in the tumor environment in controlling the differentiation of MDSCs into TAM [68]. In conclusion, the STAT regulatory pathway could be a potential target for lung cancer therapy.

4.3. Elimination of MDSCs. MDSCs can be directly eliminated in pathological settings by using some chemotherapeutic drugs and antibodies. The administration of gemcitabine to tumor-bearing mice resulted in a dramatic reduction in the number of MDSCs in the spleen and resulted in a great improvement in the antitumor response induced by immunotherapy [29]. Not only that, a combination treatment with gemcitabine and a superoxide dismutase mimetic that targets MDSCs in the tumor microenvironment can enhance the quantity and quality of both effector and memory CD8⁺ T-cell responses [69]. In addition, the generalized depletion of MDSCs, as obtained with anti-Gr-1 or anti-Ly6G antibodies, not only improves APCs, NK, and T-cell immune activities but also promotes angiostasis, leading to more efficient control of tumor growth [59].

4.4. Attenuation of MDSC Function. In addition, targeting MDSCs' function will be useful for controlling cancer growth and may be more efficient in combination with other immunomodulatory strategies [70]. The infiltration of MDSCs into the spleen of tumor-bearing mice was significantly decreased after being treated with gemcitabine, and the antitumor immune response was significantly enhanced [29, 71]. A recent publication reported that the treatment of mice bearing the LP07 lung adenocarcinoma with indomethacin (IND) inhibited the suppressive activity of splenic MDSCs, which restrained tumor growth through mechanisms involving CD8⁺ T cells [60]. Hoeppner et al. demonstrated that D₂R agonists may reduce lung tumor growth through the inhibition of immunosuppressive MDSCs as well as the abrogation of tumor angiogenesis [72]. Moreover, in our previous study, we found that the inhibition of miR-9 promoted the differentiation of MDSCs with significantly reduced immunosuppressive function, whereas the overexpression of miR-9 markedly enhanced the function of MDSCs. Notably, the knockdown of miR-9 significantly impaired the activity of MDSCs and inhibited the tumor growth of Lewis lung carcinoma in mice [55].

Cyclooxygenase 2 (COX2) is considered to be a potential target molecule. The inhibition of COX2 expression in MDSCs can decrease the release of arginine, thus promoting the antitumor immune response and enhancing the effect of immune therapy. The COX2 overexpression in lung cancer and the process of epithelial mesenchymal transition (EMT) are supposed to play an important role in the inhibition of antitumor immune response by MDSCs [73, 74]. The inhibitor of ROS can also effectively reduce the immunosuppression mediated by MDSCs. In a recent study, the results indicate that the antioxidant systems directed by Nrf2 and

selenoenzymes contribute to the clearance of ROS in MDSCs, efficiently preventing cancer cell metastasis [42, 75]. Zheng et al. found that cimetidine reduced MDSCs accumulating in the spleen, blood, and tumor tissue of tumor-bearing mice. Further investigation demonstrated that the NO production and Arg-1 expression of MDSCs were reduced, and MDSCs were prone to apoptosis by cimetidine treatment [76].

4.5. Blockade of Immune Checkpoint. In addition to the above, immune checkpoint inhibition is a new treatment approach that is undergoing extensive investigation in lung cancer. There is emerging evidence that signaling through programmed death-ligand 1 (PD-L1) plays an essential role in the immune escape of cancers linked to MDSCs [77]. Ajona et al. put forward that the combined blockade of PD-1/PD-L1 and C5a can restore antitumor immune responses, inhibit tumor cell growth, and improve outcomes of patients with lung cancer. This effect is accompanied by a negative association between the frequency of CD8 T cells and MDSCs within tumors [78]. After that, Ballbach et al. demonstrated that PD-L1 is expressed on granulocytic MDSCs upon coculture with T cells. Targeting PD-L1 also partially impaired MDSC-mediated T-cell suppression [79].

5. Concluding Remarks

In conclusion, MDSCs play an important role in the development and progression of lung cancer and can be used as a potential target for lung cancer treatment. We believe MDSC-targeted immunotherapy has good potential in the future. Of course, just relying on MDSCs is not enough, and the combination of different strategies should be considered, such as CAR-T immunotherapy [80, 81]. Currently, relatively new areas of research are mainly focused on the regulation of noncoding RNA in MDSCs [82] and the impact of changes in the metabolic status of MDSCs on its aggregation, differentiation, and function, which means the implications for metabolic reprogramming exist as a cancer therapeutic approach. An in-depth study of MDSCs immunotherapy will progress the treatment of lung cancer into a new era.

Abbreviations

MDSC:	Myeloid-derived suppressor cell
M-MDSC:	Monocytic MDSC
G-MDSC:	Granulocytic MDSC
DC:	Dendritic cell
HR1:	Histamine receptor 1
BCL-XL:	B-cell lymphoma XL
TLR:	Toll-like receptor
ADAM:	A disintegrin and metalloproteinase
miRNA:	MicroRNAs
ROS:	Reactive oxygen species
MEF2C:	Myeloid enhancement factor 2C
Arg-1:	Arginase-1
Treg:	Regulatory T cell
SCLC:	Small cell lung cancer
NSCLC:	Nonsmall cell lung cancer

SWCNT: Single-walled carbon nanotubes
 ATRA: All-*trans*-retinoic acid
 SCF: Stem cell factor
 VEGF: Vascular endothelial growth factor
 COX2: Cyclooxygenase 2
 EMT: Epithelial mesenchymal transition.

Ethical Approval

This article does not contain any studies with human participants performed by any of the authors.

Conflicts of Interest

The authors declare no conflicts of interest.

Acknowledgments

This work was supported by the Summit of the Six Top Talents Program of Jiangsu Province (Grant no. 2015-WSN-116), Jiangsu Province's Key Medical Talents Program (Grant no. ZDRCB2016018), Jiangsu Province "333" Project (Grant no. BRA2017128), Specialized Project for Clinical Medicine of Jiangsu Province (Grant no. BL2014065), Project Funded by China Postdoctoral Science Foundation (Grant no. 2016M600382), Jiangsu Postdoctoral Science Foundation Funded Project (Grant no. 1601082B), and Jiangsu University Science Foundation (Grant nos. 11JDG093 and FCJJ2015022).

References

- [1] R. L. Siegel, K. D. Miller, and A. Jemal, "Cancer statistics, 2015," *CA: A Cancer Journal for Clinicians*, vol. 65, no. 1, pp. 5–29, 2015.
- [2] E. Byron and M. Pinder-Schenck, "Systemic and targeted therapies for early-stage lung cancer," *Cancer Control*, vol. 21, no. 1, pp. 21–31, 2014.
- [3] R. D. Hall, J. E. Gray, and A. A. Chiappori, "Beyond the standard of care: a review of novel immunotherapy trials for the treatment of lung cancer," *Cancer Control*, vol. 20, no. 1, pp. 22–31, 2013.
- [4] D. I. Gabrilovich and S. Nagaraj, "Myeloid-derived suppressor cells as regulators of the immune system," *Nature Reviews Immunology*, vol. 9, no. 3, pp. 162–174, 2009.
- [5] J. I. Youn, S. Nagaraj, M. Collazo, and D. I. Gabrilovich, "Subsets of myeloid-derived suppressor cells in tumor-bearing mice," *Journal of Immunology*, vol. 181, no. 8, pp. 5791–5802, 2008.
- [6] R. K. Martin, S. J. Saleem, L. Folgosa et al., "Mast cell histamine promotes the immunoregulatory activity of myeloid-derived suppressor cells," *Journal of Leukocyte Biology*, vol. 96, no. 1, pp. 151–159, 2014.
- [7] V. Bronte, S. Brandau, S. H. Chen et al., "Recommendations for myeloid-derived suppressor cell nomenclature and characterization standards," *Nature Communications*, vol. 7, article 12150, 2016.
- [8] Y. Wang, J. Tian, and S. Wang, "The potential therapeutic role of myeloid-derived suppressor cells in autoimmune arthritis," *Seminars in Arthritis and Rheumatism*, vol. 45, no. 4, pp. 490–495, 2016.
- [9] V. Bronte and P. Zanovello, "Regulation of immune responses by L-arginine metabolism," *Nature Reviews Immunology*, vol. 5, no. 8, pp. 641–654, 2005.
- [10] P. C. Rodriguez and A. C. Ochoa, "Arginine regulation by myeloid derived suppressor cells and tolerance in cancer: mechanisms and therapeutic perspectives," *Immunological Reviews*, vol. 222, no. 1, pp. 180–191, 2008.
- [11] P. C. Rodriguez, C. P. Hernandez, D. Quiceno et al., "Arginase I in myeloid suppressor cells is induced by COX-2 in lung carcinoma," *The Journal of Experimental Medicine*, vol. 202, no. 7, pp. 931–939, 2005.
- [12] M. Fletcher, M. E. Ramirez, R. A. Sierra et al., "L-arginine depletion blunts antitumor T-cell responses by inducing myeloid-derived suppressor cells," *Cancer Research*, vol. 75, no. 2, pp. 275–283, 2015.
- [13] P. C. Rodriguez, D. G. Quiceno, and A. C. Ochoa, "L-arginine availability regulates T-lymphocyte cell-cycle progression," *Blood*, vol. 109, no. 4, pp. 1568–1573, 2007.
- [14] R. M. Bingisser, P. A. Tilbrook, P. G. Holt, and U. R. Kees, "Macrophage-derived nitric oxide regulates T cell activation via reversible disruption of the Jak3/STAT5 signaling pathway," *Journal of Immunology*, vol. 160, pp. 5729–5734, 1998.
- [15] L. Rivoltini, M. Carrabba, V. Huber et al., "Immunity to cancer: attack and escape in T lymphocyte-tumor cell interaction," *Immunological Reviews*, vol. 188, no. 1, pp. 97–113, 2002.
- [16] A. Szuster-Ciesielska, E. Hryciuk-Umer, A. Stepulak, K. Kupisz, and M. Kandefers-Szerszen, "Reactive oxygen species production by blood neutrophils of patients with laryngeal carcinoma and antioxidative enzyme activity in their blood," *Acta Oncologica*, vol. 43, no. 3, pp. 252–258, 2004.
- [17] G. Waris and H. Ahsan, "Reactive oxygen species: role in the development of cancer and various chronic conditions," *Journal of Carcinogenesis*, vol. 5, no. 1, p. 14, 2006.
- [18] G. Mantovani, A. Maccio, C. Madeddu et al., "Antioxidant agents are effective in inducing lymphocyte progression through cell cycle in advanced cancer patients: assessment of the most important laboratory indexes of cachexia and oxidative stress," *Journal of Molecular Medicine*, vol. 81, no. 10, pp. 664–673, 2003.
- [19] E. Agostinelli and N. Seiler, "Non-irradiation-derived reactive oxygen species (ROS) and cancer: therapeutic implications," *Amino Acids*, vol. 31, no. 3, pp. 341–355, 2006.
- [20] S. Kusmartsev, Y. Nefedova, D. Yoder, and D. I. Gabrilovich, "Antigen-specific inhibition of CD8⁺ T cell response by immature myeloid cells in cancer is mediated by reactive oxygen species," *Journal of Immunology*, vol. 172, p. 4647, 2004.
- [21] H. Sauer, M. Wartenberg, and J. Hescheler, "Reactive oxygen species as intracellular messengers during cell growth and differentiation," *Cellular Physiology and Biochemistry*, vol. 11, pp. 173–186, 2001.
- [22] C. S. Cobbs, T. R. Whisenhunt, D. R. Wesemann, L. E. Harkins, E. G. Van Meir, and M. Samanta, "Inactivation of wild-type p53 protein function by reactive oxygen and nitrogen species in malignant glioma cells," *Cancer Research*, vol. 63, no. 24, pp. 8670–8673, 2003.
- [23] B. G. Bentz, G. K. Haines 3rd, and J. A. Radosevich, "Increased protein nitrosylation in head and neck squamous cell carcinogenesis," *Head & Neck*, vol. 22, no. 1, pp. 64–70, 2000.
- [24] J. Dairou, J. M. Dupret, and F. Rodrigues-Lima, "Impairment of the activity of the xenobiotic-metabolizing enzymes

- arylamine *N*-acetyltransferases 1 and 2 (NAT1/NAT2) by peroxynitrite in mouse skeletal muscle cells," *FEBS Letters*, vol. 579, no. 21, pp. 4719–4723, 2005.
- [25] V. L. Kinnula, T. Torkkeli, P. Kristo et al., "Ultrastructural and chromosomal studies on manganese superoxide dismutase in malignant mesothelioma," *American Journal of Respiratory Cell and Molecular Biology*, vol. 31, no. 2, pp. 147–153, 2004.
- [26] Y. Nakamura, H. Yasuoka, M. Tsujimoto et al., "Nitric oxide in breast cancer: induction of vascular endothelial growth factor-C and correlation with metastasis and poor prognosis," *Clinical Cancer Research*, vol. 12, no. 4, pp. 1201–1207, 2006.
- [27] D. I. Gabrilovich, S. Ostrand-Rosenberg, and V. Bronte, "Coordinated regulation of myeloid cells by tumours," *Nature Reviews Immunology*, vol. 12, no. 4, pp. 253–268, 2012.
- [28] S. Nagaraj, K. Gupta, V. Pisarev et al., "Altered recognition of antigen is a mechanism of CD8⁺ T cell tolerance in cancer," *Nature Medicine*, vol. 13, no. 7, pp. 828–835, 2007.
- [29] E. Suzuki, V. Kapoor, A. S. Jassar, L. R. Kaiser, and S. M. Albelda, "Gemcitabine selectively eliminates splenic Gr-1⁺/CD11b⁺ myeloid suppressor cells in tumor-bearing animals and enhances antitumor immune activity," *Clinical Cancer Research*, vol. 11, no. 18, pp. 6713–6721, 2005.
- [30] C. Liu, S. Yu, J. Kappes et al., "Expansion of spleen myeloid suppressor cells represses NK cell cytotoxicity in tumor-bearing host," *Blood*, vol. 109, no. 10, pp. 4336–4342, 2007.
- [31] C. A. Crane, S. J. Han, J. J. Barry, B. J. Ahn, L. L. Lanier, and A. T. Parsa, "TGF- β downregulates the activating receptor NKG2D on NK cells and CD8⁺ T cells in glioma patients," *Neuro-Oncology*, vol. 12, no. 1, pp. 7–13, 2010.
- [32] B. Huang, P. Y. Pan, Q. Li et al., "Gr-1+CD115⁺ immature myeloid suppressor cells mediate the development of tumor-induced T regulatory cells and T-cell anergy in tumor-bearing host," *Cancer Research*, vol. 66, no. 2, pp. 1123–1131, 2006.
- [33] R. Yang, Z. Cai, Y. Zhang, W. H. Yutzy 4th, K. F. Roby, and R. B. Roden, "CD80 in immune suppression by mouse ovarian carcinoma-associated Gr-1⁺CD11b⁺ myeloid cells," *Cancer Research*, vol. 66, no. 13, pp. 6807–6815, 2006.
- [34] K. Movahedi, M. Guillemins, and J. Van den Bossche, "Identification of discrete tumor-induced myeloid-derived suppressor cell subpopulations with distinct T cell-suppressive activity," *Blood*, vol. 111, no. 8, pp. 4233–4244, 2008.
- [35] Z. Deng, Y. Rong, Y. Teng et al., "Exosomes miR-126a released from MDSC induced by DOX treatment promotes lung metastasis," *Oncogene*, vol. 36, no. 5, pp. 639–651, 2016.
- [36] T. Wu, Y. Zhao, H. Wang et al., "mTOR masters monocytic myeloid-derived suppressor cells in mice with allografts or tumors," *Scientific Reports*, vol. 6, no. 1, article 20250, 2016.
- [37] X. Chen, Z. Zhang, Y. Bi et al., "mTOR signaling disruption from myeloid-derived suppressive cells protects against immune-mediated hepatic injury through the HIF1 α -dependent glycolytic pathway," *Journal of Leukocyte Biology*, vol. 100, no. 6, pp. 1349–1362, 2016.
- [38] F. Hossain, A. A. Al-Khami, D. Wyczzechowska et al., "Inhibition of fatty acid oxidation modulates immunosuppressive functions of myeloid-derived suppressor cells and enhances cancer therapies," *Cancer Immunology Research*, vol. 3, no. 11, pp. 1236–1247, 2015.
- [39] Z. Husain, Y. Huang, P. Seth, and V. P. Sukhatme, "Tumor-derived lactate modifies antitumor immune response: effect on myeloid-derived suppressor cells and NK cells," *Journal of Immunology*, vol. 191, no. 3, pp. 1486–1495, 2013.
- [40] G. Liu, Y. Bi, B. Shen et al., "SIRT1 limits the function and fate of myeloid-derived suppressor cells in tumors by orchestrating HIF-1 α -dependent glycolysis," *Cancer Research*, vol. 74, no. 3, pp. 727–737, 2014.
- [41] T. Tian, X. Gu, B. Zhang et al., "Increased circulating CD14(+)/HLA-DR-/low myeloid-derived suppressor cells are associated with poor prognosis in patients with small-cell lung cancer," *Cancer Biomarkers*, vol. 15, no. 4, pp. 425–432, 2015.
- [42] A. Huang, B. Zhang, B. Wang, F. Zhang, K. X. Fan, and Y. J. Guo, "Increased CD14⁺HLA-DR^{-low} myeloid-derived suppressor cells correlate with extrathoracic metastasis and poor response to chemotherapy in non-small cell lung cancer patients," *Cancer Immunology, Immunotherapy*, vol. 62, no. 9, pp. 1439–1451, 2013.
- [43] P. L. de Goeje, K. Bezemer, M. E. Heuvers et al., "Immunoglobulin-like transcript 3 is expressed by myeloid-derived suppressor cells and correlates with survival in patients with non-small cell lung cancer," *OncoImmunology*, vol. 4, no. 7, article e1014242, 2015.
- [44] M. E. Heuvers, F. Muskens, K. Bezemer et al., "Arginase-1 mRNA expression correlates with myeloid-derived suppressor cell levels in peripheral blood of NSCLC patients," *Lung Cancer*, vol. 81, no. 3, pp. 468–474, 2013.
- [45] P. H. Feng, K. Y. Lee, Y. L. Chang et al., "CD14⁺S100A9⁺ monocytic myeloid-derived suppressor cells and their clinical relevance in non-small cell lung cancer," *American Journal of Respiratory and Critical Care Medicine*, vol. 186, no. 10, pp. 1025–1036, 2012.
- [46] G. Zhang, H. Huang, Y. Zhu et al., "A novel subset of B7-H3⁺CD14⁺HLA-DR^{-low} myeloid-derived suppressor cells are associated with progression of human NSCLC," *OncoImmunology*, vol. 4, no. 2, article e977164, 2015.
- [47] E. K. Vetsika, F. Koinis, M. Gioulbasani et al., "A circulating subpopulation of monocytic myeloid-derived suppressor cells as an independent prognostic/predictive factor in untreated non-small lung cancer patients," *Journal of Immunology Research*, vol. 2014, Article ID 659294, 12 pages, 2014.
- [48] S. Wang, Y. Fu, K. Ma et al., "The significant increase and dynamic changes of the myeloid-derived suppressor cells percentage with chemotherapy in advanced NSCLC patients," *Clinical and Translational Oncology*, vol. 16, no. 7, pp. 616–622, 2014.
- [49] F. Koinis, E. K. Vetsika, D. Aggouraki et al., "Effect of first-line treatment on myeloid-derived suppressor cells' subpopulations in the peripheral blood of patients with non-small cell lung cancer," *Journal of Thoracic Oncology*, vol. 11, no. 8, pp. 1263–1272, 2016.
- [50] T. Schneider, A. Sevko, C. P. Heussel et al., "Serum inflammatory factors and circulating immunosuppressive cells are predictive markers for efficacy of radiofrequency ablation in non-small-cell lung cancer," *Clinical & Experimental Immunology*, vol. 180, no. 3, pp. 467–474, 2015.
- [51] T. Wang, Z. Chu, H. Lin, J. Jiang, X. Zhou, and X. Liang, "Galectin-3 contributes to cisplatin-induced myeloid derived suppressor cells (MDSCs) recruitment in lewis lung cancer-bearing mice," *Molecular Biology Reports*, vol. 41, no. 6, pp. 4069–4076, 2014.
- [52] A. A. Shvedova, E. R. Kisin, N. Yanamala et al., "MDSC and TGF β are required for facilitation of tumor growth in the lungs

- of mice exposed to carbon nanotubes," *Cancer Research*, vol. 75, no. 8, pp. 1615–1623, 2015.
- [53] M. L. Ortiz, L. Lu, I. Ramachandran, and D. I. Gabrilovich, "Myeloid-derived suppressor cells in the development of lung cancer," *Cancer Immunology Research*, vol. 2, no. 1, pp. 50–58, 2014.
- [54] L. M. Hengesbach and K. A. Hoag, "Physiological concentrations of retinoic acid favor myeloid dendritic cell development over granulocyte development in cultures of bone marrow cells from mice," *The Journal of Nutrition*, vol. 134, no. 10, pp. 2653–2659, 2004.
- [55] J. Tian, K. Rui, X. Tang et al., "MicroRNA-9 regulates the differentiation and function of myeloid-derived suppressor cells via targeting Runx1," *Journal of Immunology*, vol. 195, no. 3, pp. 1301–1311, 2015.
- [56] J. D. Veltman, M. E. Lambers, M. van Nimwegen et al., "COX-2 inhibition improves immunotherapy and is associated with decreased numbers of myeloid-derived suppressor cells in mesothelioma. Celecoxib influences MDSC function," *BMC Cancer*, vol. 10, no. 1, p. 464, 2010.
- [57] S. J. Priceman, J. L. Sung, Z. Shaposhnik et al., "Targeting distinct tumor-infiltrating myeloid cells by inhibiting CSF-1 receptor: combating tumor evasion of antiangiogenic therapy," *Blood*, vol. 115, no. 7, pp. 1461–1471, 2010.
- [58] P. Y. Pan, G. X. Wang, B. Yin et al., "Reversion of immune tolerance in advanced malignancy: modulation of myeloid-derived suppressor cell development by blockade of stem-cell factor function," *Blood*, vol. 111, no. 1, pp. 219–228, 2008.
- [59] M. K. Srivastava, L. Zhu, M. Harris-White et al., "Myeloid suppressor cell depletion augments antitumor activity in lung cancer," *PLoS One*, vol. 7, no. 7, article e40677, 2012.
- [60] A. G. Blidner, M. Salatino, I. D. Mascanfroni et al., "Differential response of myeloid-derived suppressor cells to the nonsteroidal anti-inflammatory agent indomethacin in tumor-associated and tumor-free microenvironments," *Journal of Immunology*, vol. 194, no. 7, pp. 3452–3462, 2015.
- [61] C. Iclozan, S. Antonia, A. Chiappori, D. T. Chen, and D. Gabrilovich, "Therapeutic regulation of myeloid-derived suppressor cells and immune response to cancer vaccine in patients with extensive stage small cell lung cancer," *Cancer Immunology, Immunotherapy*, vol. 62, no. 5, pp. 909–918, 2013.
- [62] S. Nagaraj, J. I. Youn, H. Weber et al., "Anti-inflammatory triterpenoid blocks immune suppressive function of MDSCs and improves immune response in cancer," *Clinical Cancer Research*, vol. 16, no. 6, pp. 1812–1823, 2010.
- [63] Y. Nefedova, S. Nagaraj, A. Rosenbauer, C. Muro-Cacho, S. M. Sebt, and D. I. Gabrilovich, "Regulation of dendritic cell differentiation and antitumor immune response in cancer by pharmacologic-selective inhibition of the Janus-activated kinase 2/signal transducers and activators of transcription 3 pathway," *Cancer Research*, vol. 65, no. 20, pp. 9525–9535, 2005.
- [64] M. Kortylewski, M. Kujawski, T. Wang et al., "Inhibiting Stat3 signaling in the hematopoietic system elicits multicomponent antitumor immunity," *Nature Medicine*, vol. 11, no. 12, pp. 1314–1321, 2005.
- [65] D. Foell, H. Wittkowski, T. Vogl, and J. Roth, "S100 proteins expressed in phagocytes: a novel group of damage-associated molecular pattern molecules," *Journal of Leukocyte Biology*, vol. 81, no. 1, pp. 28–37, 2007.
- [66] L. Wu, H. Du, Y. Li, P. Qu, and C. Yan, "Signal transducer and activator of transcription 3 (Stat3c) promotes myeloid-derived suppressor cell expansion and immune suppression during lung tumorigenesis," *The American Journal of Pathology*, vol. 179, no. 4, pp. 2131–2141, 2011.
- [67] J. Zhou, Z. Qu, F. Sun et al., "Myeloid STAT3 promotes lung tumorigenesis by transforming tumor immunosurveillance into tumor-promoting inflammation," *Cancer Immunology Research*, vol. 5, no. 3, pp. 257–268, 2017.
- [68] V. Kumar, P. Cheng, T. Condamine et al., "CD45 phosphatase inhibits STAT3 transcription factor activity in myeloid cells and promotes tumor-associated macrophage differentiation," *Immunity*, vol. 44, no. 2, pp. 303–315, 2016.
- [69] A. Sawant, C. C. Schafer, T. H. Jin et al., "Enhancement of anti-tumor immunity in lung cancer by targeting myeloid-derived suppressor cell pathways," *Cancer Research*, vol. 73, no. 22, pp. 6609–6620, 2013.
- [70] M. K. Srivastava, S. Dubinett, and S. Sharma, "Targeting MDSCs enhance therapeutic vaccination responses against lung cancer," *Oncotarget*, vol. 1, no. 9, pp. 1650–1651, 2012.
- [71] H. J. Ko, Y. J. Kim, Y. S. Kim et al., "A combination of chemoimmunotherapies can efficiently break self-tolerance and induce antitumor immunity in a tolerogenic murine tumor model," *Cancer Research*, vol. 67, no. 15, pp. 7477–7486, 2007.
- [72] L. H. Hoepfner, Y. Wang, A. Sharma et al., "Dopamine D2 receptor agonists inhibit lung cancer progression by reducing angiogenesis and tumor infiltrating myeloid derived suppressor cells," *Molecular Oncology*, vol. 9, no. 1, pp. 270–281, 2015.
- [73] C. A. Dasanu, N. Sethi, and N. Ahmed, "Immune alterations and emerging immunotherapeutic approaches in lung cancer," *Expert Opinion on Biological Therapy*, vol. 12, no. 7, pp. 923–937, 2012.
- [74] J. P. Thiery, "Epithelial–mesenchymal transitions in tumour progression," *Nature Reviews. Cancer*, vol. 2, no. 6, pp. 442–454, 2002.
- [75] K. Hiramoto, H. Satoh, T. Suzuki et al., "Myeloid lineage-specific deletion of antioxidant system enhances tumor metastasis," *Cancer Prevention Research*, vol. 7, no. 8, pp. 835–844, 2014.
- [76] Y. Zheng, M. Xu, X. Li, J. Jia, K. Fan, and G. Lai, "Cimetidine suppresses lung tumor growth in mice through proapoptosis of myeloid-derived suppressor cells," *Molecular Immunology*, vol. 54, no. 1, pp. 74–83, 2013.
- [77] M. Gato-Canas, M. Zuazo, H. Arasanz et al., "PDL1 signals through conserved sequence motifs to overcome interferon-mediated cytotoxicity," *Cell Reports*, vol. 20, no. 8, pp. 1818–1829, 2017.
- [78] D. Ajona, S. Ortiz-Espinosa, H. Moreno et al., "A combined PD-1/C5a blockade synergistically protects against lung cancer growth and metastasis," *Cancer Discovery*, vol. 7, no. 7, pp. 694–703, 2017.
- [79] M. Ballbach, A. Dannert, A. Singh et al., "Expression of checkpoint molecules on myeloid-derived suppressor cells," *Immunology Letters*, vol. 192, pp. 1–6, 2017.
- [80] S. C. Katz, G. R. Point, M. Cunetta et al., "Regional CAR-T cell infusions for peritoneal carcinomatosis are superior to systemic delivery," *Cancer Gene Therapy*, vol. 23, no. 5, pp. 142–148, 2016.

- [81] A. Stiff, P. Trikha, R. Wesolowski et al., "Myeloid-derived suppressor cells express Bruton's tyrosine kinase and can be depleted in tumor-bearing hosts by ibrutinib treatment," *Cancer Research*, vol. 76, no. 8, pp. 2125–2136, 2016.
- [82] X. Tian, J. Ma, T. Wang et al., "LncRNA HOTAIRM1-HOXA1 axis down-regulates the immunosuppressive activity of myeloid-derived suppressor cells in lung cancer," *Frontiers in Immunology*, vol. 9, p. 473, 2018.



Hindawi

Submit your manuscripts at
www.hindawi.com

